

Supplement of Atmos. Chem. Phys., 19, 2527–2560, 2019
<https://doi.org/10.5194/acp-19-2527-2019-supplement>
© Author(s) 2019. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Overview paper: New insights into aerosol and climate in the Arctic

Jonathan P. D. Abbatt et al.

Correspondence to: Jonathan P. D. Abbatt (jonathan.abbatt@utoronto.ca) and W. Richard Leitch (leitchs@gmail.com)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Supplementary Information

Particle Measurements at Alert, Nunavut

The experimental methods for the particle microphysical and filter-based chemical measurements in Fig. 7 are discussed by Leaitch et al. (2013, 2018).

Monthly-averaged values of $N_{>50}$ (number concentration > 50 nm diameter), $N_{>100}$, N_{15-30} (number concentration in the size interval 15 nm to 30 nm), N_{30-50} , and N_{50-100} for the inclusive period of 2012–2014 are shown in Fig. 7a.

One estimate of the effect of growth of newly-formed particles on the concentrations of particles in the range of 15–100 nm is demonstrated using Fig. 2. Baseline concentrations are estimated by linearly interpolating each of the N_{15-30} , N_{30-50} and N_{50-100} from May to October. These are shown as the dashed lines in Fig. 7a. The increases in N_{15-30} , N_{30-50} and N_{50-100} during the summer months, relative to the estimated baseline concentrations, are shown in Fig. 7b. The progressive decrease in N_{15-30} to N_{30-50} to N_{50-100} is indicative of the average of growth from smaller to larger sizes. There is no obvious increase in particles larger than 100 nm during the summer, suggesting growth to such sizes is mostly undetectable by this approach. Despite the high frequency of NPF, the growth to 50 nm or larger during August is almost identical to that in July, suggesting that the levels of precursors for condensational growth are on average lower during August than July. The monthly-average OM/SO_4^{2-} , also shown in Figure 7b (from Leaitch et al. (2018)), increases substantially and coincidentally with the increases in 15–50 nm particle concentrations, suggesting that organic components are important for the growth of newly formed particles in the summer Arctic.

Two approaches were used to estimate the percentages of 30–50 nm particles resulting from naturally-formed new particles. The July and August total N_{30-50} in Fig. 7a are 48 cm^{-3} and 66 cm^{-3} , respectively. Two approaches are used to estimate the reference or baseline for the non-natural contributions. One approach is to use the baseline interpolated from May to October (dashed lines in Fig. 7a). The second approach assesses the non-natural baseline based on black carbon (BC) measurements, as discussed by Leaitch et al. (2013). The first approach yields natural number concentrations for the July and August months discussed in Leaitch et al. (2018) of 33 cm^{-3} and 54 cm^{-3} , respectively. The BC approach gives natural number concentrations of 9 cm^{-3} and 13 cm^{-3} for July and August respectively. The BC approach most likely underestimates the natural influence because remnant BC can remain during summer even while natural processes re-generate particles. The result based on the interpolation in Fig. 7 is an upper estimate of the natural contribution to the 30–50 nm particles because it assumes no anthropogenic contributions above the interpolated baseline. Therefore, at Alert, the ranges of natural contributions to the average July and August N_{30-50} are estimated to be 20–70% for July and 20–80% for August.

References

Leaitch, W. R., Sharma, S., Huang, L., Toom-Sauntry, D., Chivulescu, A., Macdonald, A. M., von Salzen, K., Pierce, J. R., Betram, A. K., Schroder, J. C., Shantz, N. C., Chang, R. and Norman, A. L.: Dimethyl sulfide control of the clean summertime Arctic aerosol and cloud, *Elementa*, 1, 1–000017, doi:10.12952/journal.elementa.000017, 2013.

Leaitch, W. R., Russell, L. M., Liu, J., Kolonjari, F., Toom, D., Huang, L., Sharma, S., Chivulescu, A., Veber, D. and Zhang, W.: Organic functional groups in the submicron aerosol at 82.5° N, 62.5° W from 2012 to 2014, *Atmos. Chem. Phys.*, 18(5), 3269–3287, doi:10.5194/acp-18-3269-2018, 2018.