

To cite this manuscript, use the following citation

St-Amour, S., Hains-Monfette, G., Dancause, K. N., Cailhol, L., & Bernard, P. (2021). Antidepressant medication use and objectively measured physical activity and sedentary behaviors in adults: A cross-sectional analysis of a nationally representative sample of Canadian adults. *Mental Health and Physical Activity*, 20, 100394. <https://doi.org/10.1016/j.mhpa.2021.100394>

Antidepressant medication use and objectively measured physical activity and sedentary behaviors in adults: a cross-sectional analysis of a nationally representative sample of Canadian adults.

St-Amour Samuel^{1,2}, Hains-Monfette Gabriel^{1,2}, Dancause Kelsey Needham¹, Cailhol Lionel², Bernard Paquito^{1,2}

¹Department of Physical Activity Sciences, Université du Québec à Montréal, Montréal, Québec, Canada

²Research Center, University Institute of Mental Health at Montreal, Montréal, Quebec, Canada

Correspondence concerning this article should be addressed to:

Paquito Bernard, PhD, Université du Québec à Montréal, Montréal, Canada
Faculté des sciences, UQÀM, Complexe des Sciences, Pavillon des sciences biologiques (SB), Local: SB-4445, 141, Avenue du Président Kennedy, Montréal, Québec, Canada, H2X 1Y4

bernard.paquito@uqam.ca

ORCID: <https://orcid.org/0000-0003-2180-9135>

Editorial disclosure:

Given his role as Editorial Board Member, Paquito Bernard had no involvement in the peer-review of this article and has no access to information regarding its peer-review. Full responsibility for the editorial process for this article was delegated to Adrian Taylor.

Introduction

Antidepressant medications represent one of the most prescribed medications in Canada, and are among the top five prescription medications used in Canadians from 15 to 64 years. Data from nationally representative samples show that antidepressant use was reported by 8% and 17% of adults aged 45 to 64 for men and women, respectively (Rotermann et al., 2014). Moreover, the prevalence of antidepressant prescriptions increased between 2006 and 2012 in primary care (from 9.2% to 12.8%). However, the incidence of prescription did not rise during the same period (Morkem et al., 2015). These different patterns suggested a more prolonged antidepressant use in Canadian adults.

Antidepressant users represent a heterogeneous population including individuals with depressive or other disorders. Antidepressants are an evidence-based pharmacological treatment primarily used for the first-line treatment of major depressive disorder (Kennedy et al., 2016). Different antidepressant classes are prescribed based on patient and medication characteristics. However, adults using antidepressants do not represent only patients with a depressive disorder. For instance, fluoxetine can be prescribed in adults with obsessive-compulsive disorder or panic disorder. Furthermore, recent investigations showed an important prevalence of antidepressant prescriptions for off-label uses (i.e., prescriptions for indications that have not been approved by health authorities; Carton et al., 2015) in non-psychiatric settings (Deferio et al., 2018; Skånland & Cieślár-Pobuda, 2019; Wong et al., 2017). Off-label prescriptions represented 15 to 30% of antidepressant prescriptions in primary care in a Canadian study (Wong et al., 2017). Tricyclic antidepressants were the most prescribed antidepressant class among antidepressant off-label prescriptions. The most common antidepressant off-label indications were anxiety, insomnia, chronic pain or migraine, eating disorders and smoking cessation (Deferio et al., 2018; Skånland & Cieślár-Pobuda, 2019; Wong et al., 2016). Overall, almost half the antidepressant prescriptions were for other disorders than depressive disorder (Wong et al., 2016).

Although antidepressant medications are effective to treat adults with major depressive disorders, antidepressant use could be (in)directly associated with lower and higher time spent in physical activity (PA) and sedentary behaviors, respectively. In a mice study, SSRI was related to a motor activity decrease (Marlatt et al., 2010). An increased risk of muscle injuries has been reported in adults with SSRI treatment (Labotz et al., 2006). Antidepressant use (i.e., antidepressants from tricyclic, SSRI and atypical categories) is associated with higher risk of cardiometabolic diseases and weight gain (Carvalho et al., 2016; Serretti & Mandelli, 2010), longitudinal risk of fractures, and gait disturbances in the elderly (Coupland et al., 2018). The weight gain associated with antidepressant use has been linked to a greater caloric intake and more time spent in sedentary behaviors (Jensen-Otsu & Austin, 2015). Moreover, adults with severe mental illness were less active compared to healthy adults. Among them, the antidepressant users were more inactive (Vancampfort et al., 2017). In this perspective, PA and sedentary behaviors could play a buffering effect between antidepressant use and adverse side effects (Bruins et al., 2014; Schuch et al., 2018).

PA and sedentary behaviors in adults using antidepressants have received little attention to date (Bernard & Carayol, 2015). Weekly self-reported time spent in PA was prospectively associated with a lower risk of antidepressant use in middle-aged Finnish adults (Lahti et al., 2013; Loponen et al., 2015; Stubbs et al., 2017; Waller et al., 2016). However, this association was not observed in adolescents (Kleppang et al., 2019). An increase of self-reported moderate-to-vigorous PA (MVPA) has been found as a behavioral marker of antidepressant treatment (Yun et al., 2020).

These studies suffered from several limitations. First, PA was self-reported although accelerometer data provide a more accurate quantification of the PA in daily life (Westerterp, 2009) and may exclude confounders such as recall bias commonly observed in adults with mood disorders (Morres et al., 2019). Second, previous findings about PA and sedentary behaviors in adults with major depressive disorder (Schuch et al., 2017) cannot be generalized to antidepressant users because 15 to 30% of antidepressant prescriptions may be for off-label indications and up to 45% of

antidepressant prescriptions may be for other disorders than depressive disorder (Wong et al., 2016). Third, no investigation examining PA in antidepressant users has been conducted in a nationally representative sample. Lastly, no information was available about PA and sedentary behavior patterns across antidepressant classes. In this perspective, Bernard and Carayol (2015) recommended the examination of PA and sedentary behaviors with objective measures in adults using antidepressant medications.

The aims of this study were: to describe PA and sedentary behaviors in a representative sample of Canadian adults using antidepressants, to compare objectively assessed daily time spent in PA and sedentary behaviors in a population-based sample of adults using antidepressant medication with adults without psychotropic use, and to investigate whether these behaviors differ according to four antidepressant classes. In the absence of any studies examining these accelerometer-measured behaviors, no a priori hypotheses were formulated.

Methods

Study setting and participants

This study used data collected by Statistics Canada during cycle 1, 2, and 3 (2007–2013) of the Canadian Health Measures Survey (CHMS). This national survey included people aged 6 to 79 years old and used a stratified three-stage sampling strategy (Tremblay & Gorber, 2007). Sociodemographic, general health and medication information were collected at participants' homes and direct physical measures were collected on a subsequent visit to a mobile examination center. All participants provided written informed consent. Ethics approval to conduct the survey was obtained from Health Canada's Research Ethics Board (Day et al., 2007). The current analyses included participants aged 18 to 79 years old with complete accelerometry and medication data. Pregnant women, participants with functional limitations and adults using any other psychotropic medication but not using antidepressants were excluded in our analyses. Controls did not use any psychotropic medication.

Measures

Sociodemographic and clinical characteristics

Age, level of education, household annual incomes, working status and body mass index were collected during the home interview. Self-reported diagnosis of mood disorders, level of mental health and subjective health were measured with three single-item questions. Smoking was established by urinary cotinine level measured during the mobile-examination clinic visit (Wong et al., 2013).

Antidepressant Medication

During the home visit, an interviewer asked participants to collect every medication container present in their house, collected all drug identification numbers from medication containers, then he asked when was the last time they used every medication. Current medication use was defined as any medication taken by the respondent during the previous seven days. Up to fifteen medications were recorded for each respondent. Each drug identification number was recorded according to Anatomical Therapeutic Chemical classification (ATC; Bosak et al., 2018; Bowring et al., 2017). For our analyses, psychotropic drugs from the following groups were considered: N05A antipsychotics, N05B anxiolytics, N05C hypnotics and sedatives, and N06A antidepressants. Four antidepressant medication classes were also coded: Serotonin-Norepinephrine Reuptake Inhibitors (SNRI; N06AX16/AA22, N06AX17/AA24, N06AX21, N06AX23), Selective Serotonin Reuptake Inhibitor (SSRI; N06AB), Tricyclic (N06AA [except N06AA22 AND N06AA24]) and other antidepressants (N06AF, N06AG, N06AX01-12 [or N07BA02], 13–15, 18, 19, 22, 24, 25; Laugesen et al., 2013).

Physical Activity and Sedentary Behaviors

PA and sedentary behaviors were objectively measured with Actical accelerometers (Phillips-Respironics). The Actical records time-stamped acceleration in three dimensions, thereby indicating physical activity intensity. This accelerometer has been validated to measure physical activity in adults (Heil, 2006). Participants wore the Actical over their right hip on an elasticized belt during their waking hours for seven consecutive days. To be included in the analysis, participants needed at least four

valid days of data defined as at least 10 hours of wear time (Colley et al., 2011). Time spent daily in PA at different intensity levels was categorized with validated count per minute (cpm) thresholds for adults: sedentary (<100 cpm), light (100 to 1,534 cpm) and moderate to vigorous intensity (≥ 1535 cpm; Colley & Tremblay, 2011). The Actical also measured steps per day. Participants with extreme counts (i.e. >20,000 cpm) were excluded from analyses (Colley et al., 2010).

Data Analysis

Student t tests and chi squares were performed to compare antidepressant users and controls on PA and sedentary behavior variables as well as sociodemographic, anthropometric and health variables. Four weighted analyses of covariance regression models (ANCOVA) incorporating age, sex, body mass index, accelerometer wear time, data collection season, work and marital status, smoking, education level, and income were carried out. The four dependent variables were: average time performing MVPA and light PA (LPA), average daily steps, and daily time spent in sedentary behaviors. To account for the complex, multistage probability sampling design, weights (i.e., activity monitor subsample weights combining cycles 1, 2, and 3) and bootstraps provided by Statistics Canada were used in analyses. Since the number of minutes of MVPA was not normally distributed, Poisson models were carried out for this variable. All analyses were performed during 2018 and 2019, using the “survey” package in R version 3.4.

Sensitivity Analyses

To examine the possible effect of antidepressants combined with another psychotropic medication, a set of sensitivity analyses was performed. The ANCOVAs were carried out including only adults using another psychotropic with antidepressants, then we reran models including participants using only an antidepressant medication. Based on reviewer suggestions, a second set of sensitivity analyses has been carried out. All models have been performed with self-reported levels of mental and physical health as covariates.

Results

Data from 7680 participants were included in analysis. The weighted prevalence of antidepressant medication use was 9.3% in adults 18- to 79-year-olds. Among these, 1.1% used tricyclic antidepressants, 1.7% used other antidepressants, 2.4% used SNRI, and 5.0% used SSRI. Moreover, 6.6% used only antidepressant medications and 2.9% used antidepressants along with at least another psychotropic medication. Unadjusted antidepressant users and control characteristics are provided in Table 1. Descriptive findings for adults reporting tricyclics, other antidepressants, SNRI or SSRI use; and antidepressant use combined with or without another psychotropic medication are presented in online supplements in Table S1, S2, and S3, respectively. Description of means, standard deviation and medians of LPA, MVPA, steps per day and time spent in sedentary behaviors are described in Table 2.

The weighted ANCOVAs revealed that antidepressant users spent significantly less time in MVPA ($d = -0.02$, $p = 0.03$) and performed a lower daily number of steps ($d = -0.01$, $p = 0.05$) in comparison to adults not using any psychotropic medication. Among antidepressant users, SSRI use was associated with significant lower time spent in MVPA ($d = -0.13$, $p = 0.01$) and daily steps ($d = -0.10$, $p = 0.05$). Adults classified as “other antidepressant users” had significantly lower daily steps ($d = -0.20$, $p = 0.03$) than the control group. No significant differences were found in adults using either SNRI or tricyclic antidepressants. No significant differences were observed for daily time spent in sedentary behaviors, except for adults using “other” antidepressant medications ($d = 0.18$, $p = 0.04$). All details about statistical results are available in online supplements.

The sensitivity analyses compared adults not using any psychotropic medication to adults using antidepressants and no other psychotropic medication, and adults using antidepressants and at least one other psychotropic medication. Similar findings were found for MVPA ($d = -0.10$, $p = 0.05$) and daily steps ($d = -0.10$, $p = 0.03$) in adults using only antidepressant medication. Otherwise, no significant differences were found between adults using antidepressants and at least one other psychotropic medication for PA and sedentary behavior outcomes. In multivariate models adjusted for

self-reported levels of mental and physical health, adults using SSRI had a significant lower daily time spent in MVPA ($d = -0.01$, $p = 0.01$). No other significant differences were found.

Discussion

This study compared objectively measured physical activity and sedentary behavior in adults using antidepressants with adults without psychotropic use in a nationally representative sample. To the best of our knowledge, this is the first study to examine this question with medication assessed during a home interview combined with accelerometer data in Canadians. Prevalence of antidepressant use in our sample of Canadian adults was in line with previous analyses of nationally representative data in Canada (Morkem et al., 2015) and the US (Pratt et al., 2017). Regarding sociodemographic characteristics, our antidepressant users group was comparable to the US adults using this medication (Jensen-Otsu & Austin, 2015). Our key findings were that adults with antidepressant treatment spent significantly fewer daily minutes in MVPA and performed fewer daily steps. However, it is very important to note that effect sizes were low. Thus, this difference may not be clinically significant. The differences were more marked for patients with SSRI or other antidepressant medications.

These results differ from Jensen-Otsu's investigation (2015) that did not identify a significant difference between antidepressant users and non-users for leisure PA. However, physical activity was assessed with a self-reported questionnaire. Even if no previous study assessed MVPA in antidepressant users, our findings can be compared to investigations including possible antidepressant users (e.g., adults with depressive disorders, anxiety, chronic pain or with sleep disorders; Wong et al., 2016). Thus, our findings were in accordance with previous studies that found lower objectively measured MVPA among adults with depressive disorders, chronic pain or other chronic diseases compared to healthy controls (Bernard et al., 2018; Farnsworth et al., 2015; Schuch et al., 2017). It has been suggested that antidepressant users reported more time in sedentary screen time activities (Jensen-Otsu & Austin, 2015). However, our results indicated that objectively measured time spent in sedentary behaviors was significantly higher only in adults using "other" antidepressant medications.

The fewer daily steps among antidepressant users are clinically relevant. The number of steps per day is a measure of total PA and differs from the measures of PA at defined levels of intensity (Tudor-Locke et al., 2011a). This outcome has been independently associated with numerous health outcomes (Newton et al., 2013; Tudor-Locke et al., 2018). Moreover, walking was the preferred PA in adults with depressive disorder (Burton et al., 2015; Busch et al., 2016), other mental disorders (Simonton et al., 2018), older adults and adults with chronic diseases (Wong et al., 2018).

The sensitivity analyses suggest that adults with "only" an antidepressant treatment had a significantly lower levels of PA. Nevertheless, no significant differences with controls were observed among adults combining an antidepressant with another psychotropic medication (i.e., antipsychotics, anxiolytics, hypnotics and sedatives). This suggests that other psychotropic treatments may have an antagonist effect with antidepressants and "compensate" the possible negative association between PA and antidepressant use (Bernard & Carayol, 2015). Moreover, in our sensitivity analysis controlling for self-reported mental and physical health, SSRI users performed significantly less daily minutes of MVPA than the control group. This is in line with the results from Marlatt et al. (2010) indicating a decrease in locomotor activity in mice using SSRI antidepressants. This result might suggest that the difference is attributed to a specific diagnosis for which the antidepressant was prescribed since this information were not available for this survey. It might also indicate a side effect from this specific type of medication. However, at this point causation cannot be inferred since the cross-sectional design of this study is not appropriate for this purpose and too little is known about the interaction between SSRIs and physical activity at a physiological level (Bernard & Carayol, 2015). These results and the fact that the difference appears even when our analysis is controlled for self-reported health reinforce the importance of the promotion of PA in SSRI users. It is important to note that in these sensitivity

analyses, all of the other differences observed became non-significant. As reported in supplementary files (Tables S20-S35) the difference in MVPA and daily steps observed in antidepressant users, and the difference in daily steps in SSRI and other antidepressant users would be better explained by the self-reported physical health. Considering the major prevalence of depressive symptoms in adults with physical diseases, antidepressant prescriptions are recommended to treat it (Olver & Hopwood, 2013). Moreover, it has been demonstrated that physical diseases were associated to less daily time spent in PA (Bernard et al., 2018; Bernard et al., 2018; Hains-Monfette et al., 2019).

This study set out with the aim of comparing physical activity and sedentary behavior outcomes between users of four classes of antidepressants and controls. Findings suggested that adults using SSRI or “other” antidepressants had significantly lower PA levels. Antidepressants belonging to the “other” category were mainly the Monoamine Oxidase Inhibitor (MOAI) class. These are recommended for adults with treatment-resistant major depressive disorder and atypical depression, but also adults with bipolar depression and older individuals with major depressive disorder as a second or third treatment attempt (Shulman et al., 2013). Consequently, these results may be partly explained by the fact that adults with MOAI treatment have more severe mental disorder symptoms and also higher risk of morbidity (Nelson & Spyker, 2017). Therefore, care providers may strengthen PA assessment and promotion in patients with SSRI and MOAI.

This is the first time that data from representative national samples of adults has been used to examine objectively measured PA and sedentary behaviors in antidepressant users. Thus, our findings are generalizable to the Canadian population. Additionally, the ATC information collected during a home interview decrease the risk of misclassification reported in previous studies using administrative data. Some limitations need to be considered when interpreting our results. First, the cross-sectional design precludes causal inferences. Therefore, it is impossible to establish a causality relation between PA and sedentary behaviors, and antidepressant use. Second, detailed information about mental disorder diagnostics was not available in CHMS data. Thus, our statistical models were not adjusted for these important variables. Third, the CHMS dataset contained no information about antidepressants dose, treatment duration or adherence (Bulloch & Patten, 2010). These parameters may affect PA and sedentary behaviors in CHMS participants. Fourth, our analyses were not controlled for non-psychotropic medication. Since use of prescribed medication (e.g., diuretics, anticoagulants, lipid-regulating agents) is linked to less PA (Fernandez-Navarro et al., 2018; Silva et al., 2012; Stamatakis et al., 2009) and antidepressant users are more at risk to use other medications (Sundbom & Hedborg, 2019), there might have been interactions between non-psychotropic medication and PA. Fifth, the time spent in sedentary behaviors or LPA could be underestimated because the accelerometer data were not collected with a 24h wear protocol (Matthews et al., 2012; Tudor-Locke et al., 2011b). Sixth, due to the large number of statistical analyses, the risk of type 1 is inflated. Finally, the Actical accelerometer might not be the best instrument to distinguish sedentary behaviors from LPA compared to other more recent devices (Duncan et al., 2018).

In conclusion, adults using SSRI have lower levels of MVPA than adults that do not use any psychotropics. First line care providers should consider assessing PA and sedentary behaviors (Romain et al., 2018), and particularly brisk walking behaviors, with their patients when they receive an antidepressant prescription and if relevant, help them make a plan to achieve an adequate number of steps and MVPA time daily. To help them achieve this goal, they may use guides such as *The Exercise and Depression Toolkit* (Exercise and Depression Toolkit - UBC, 2019; Glowacki et al., 2019), effective behavior change techniques (Bernard et al., 2015), or include kinesiologists in their interdisciplinary teams to specifically assess and suggest physical activity to patients (Busch et al., 2016). Future prospective cohort studies should monitor the device-measured PA and sedentary patterns in naive SSRI users before and after treatment begins. The assessment of these behaviors should be performed with more specific devices, such as the Genactive and activPAL accelerometers (Sellers et al., 2016).

References

- Bernard, P., & Carayol, M. (2015). A commentary on the importance of controlling for medication use within trials on the effects of exercise on depression and anxiety. *Mental Health and Physical Activity*, 9, 10–15. <https://doi.org/10.1016/j.mhpa.2015.07.001>
- Bernard, P., Hains-Monfette, G., Atoui, S., & Moullec, G. (2018). Daily Objective Physical Activity and Sedentary Time in Adults with COPD Using Spirometry Data from Canadian Measures Health Survey. *Canadian Respiratory Journal*, 2018. <https://doi.org/10.1155/2018/9107435>
- Bernard, Paquito, Hains-Monfette, G., Atoui, S., & Kingsbury, C. (2018). Differences in daily objective physical activity and sedentary time between women with self-reported fibromyalgia and controls: Results from the Canadian health measures survey. *Clinical Rheumatology*, 37(8), 2285–2290. <https://doi.org/10.1007/s10067-018-4139-6>
- Bernard, Paquito, Ninot, G., Cyprien, F., Courtet, P., Guillaume, S., Georgescu, V., Picot, M.-C., Taylor, A., & Quantin, X. (2015). Exercise and Counseling for Smoking Cessation in Smokers With Depressive Symptoms: A Randomized Controlled Pilot Trial. *Journal of Dual Diagnosis*, 11(3–4), 205–216. <https://doi.org/10.1080/15504263.2015.1113842>
- Bosak, M., Cyranka, K., Dudek, D., Kowalik, M., Molek, P., & Słowik, A. (2018). Psychiatric comedication in patients with epilepsy. *Epilepsy & Behavior*, 83, 207–211. <https://doi.org/10.1016/j.yebeh.2018.03.033>
- Bowring, D. L., Totsika, V., Hastings, R. P., Toogood, S., & McMahon, M. (2017). Prevalence of psychotropic medication use and association with challenging behaviour in adults with an intellectual disability. A total population study. *Journal of Intellectual Disability Research*, 61(6), 604–617. <https://doi.org/10.1111/jir.12359>
- Bruins, J., Jörg, F., Bruggeman, R., Slooff, C., Corpeleijn, E., & Pijnenborg, M. (2014). The Effects of Lifestyle Interventions on (Long-Term) Weight Management, Cardiometabolic Risk and Depressive Symptoms in People with Psychotic Disorders: A Meta-Analysis. *PLoS ONE*, 9(12), e112276. <https://doi.org/10.1371/journal.pone.0112276>
- Bulloch, A. G. M., & Patten, S. B. (2010). Non-adherence with psychotropic medications in the general population. *Social Psychiatry and Psychiatric Epidemiology*, 45(1), 47–56. <https://doi.org/10.1007/s00127-009-0041-5>
- Burton, E., Lewin, G., & Boldy, D. (2015). Physical activity preferences of older home care clients. *International Journal of Older People Nursing*, 10(3), 170–178. <https://doi.org/10.1111/opn.12065>
- Busch, A. M., Ciccolo, J. T., Puspitasari, A. J., Nosrat, S., Whitworth, J. W., & Stults-Kolehmainen, M. A. (2016). Preferences for exercise as a treatment for depression. *Mental Health and Physical Activity*, 10, 68–72. <https://doi.org/10.1016/j.mhpa.2015.12.004>
- Carton, L., Cottencin, O., Lapeyre-Mestre, M., A. Geoffroy, P., Favre, J., Simon, N., Bordet, R., & Rolland, B. (2015). Off-Label Prescribing of Antipsychotics in Adults, Children and Elderly Individuals: A Systematic Review of Recent Prescription Trends. *Current Pharmaceutical Design*, 21(23), 3280–3297.
- Carvalho, A. F., Sharma, M. S., Brunoni, A. R., Vieta, E., & Fava, G. A. (2016). The Safety, Tolerability and Risks Associated with the Use of Newer Generation Antidepressant Drugs: A Critical Review

- of the Literature. *Psychotherapy and Psychosomatics*, 85(5), 270–288. <https://doi.org/10.1159/000447034>
- Colley, R. C., Garriguet, D., Janssen, I., Craig, C. L., Clarke, J., & Tremblay, M. S. (2011). Physical activity of Canadian adults: Accelerometer results from the 2007 to 2009 Canadian Health Measures Survey. *Health Reports*, 22(1), 7–14.
- Colley, R., Connor Gorber, S., & Tremblay, M. S. (2010). Quality control and data reduction procedures for accelerometry-derived measures of physical activity. *Health Reports*, 21(1), 63–69.
- Colley, Rachel C., & Tremblay, M. S. (2011). Moderate and vigorous physical activity intensity cut-points for the Actical accelerometer. *Journal of Sports Sciences*, 29(8), 783–789. <https://doi.org/10.1080/02640414.2011.557744>
- Coupland, C., Hill, T., Morriss, R., Moore, M., Arthur, A., & Hippisley-Cox, J. (2018). Antidepressant use and risk of adverse outcomes in people aged 20–64 years: Cohort study using a primary care database. *BMC Medicine*, 16(1), 36. <https://doi.org/10.1186/s12916-018-1022-x>
- Day, B., Langlois, R., Tremblay, M., & Knoppers, B.-M. (2007). Canadian Health Measures Survey: Ethical, legal and social issues. *Health Reports*, 18 Suppl, 37–51.
- Deferio, J. J., Levin, T. T., Cukor, J., Banerjee, S., Abdulrahman, R., Sheth, A., Mehta, N., & Pathak, J. (2018). Using electronic health records to characterize prescription patterns: Focus on antidepressants in nonpsychiatric outpatient settings. *JAMIA Open*, 1(2), 233–245. <https://doi.org/10.1093/jamiaopen/ooy037>
- Duncan, S., Stewart, T., Schneller, M. B., Godbole, S., Cain, K., & Kerr, J. (2018). Convergent validity of ActiGraph and Actical accelerometers for estimating physical activity in adults. *PLOS ONE*, 13(6), e0198587. <https://doi.org/10.1371/journal.pone.0198587>
- Exercise and Depression Toolkit—UBC*. (2019). <https://exerciseanddepression.ca/>
- Farnsworth, J. L., Kim, Y., & Kang, M. (2015). Sleep Disorders, Physical Activity, and Sedentary Behavior Among U.S. Adults: National Health and Nutrition Examination Survey. *Journal of Physical Activity and Health*, 12(12), 1567–1575. <https://doi.org/10.1123/jpah.2014-0251>
- Fernandez-Navarro, P., Aragonés, M. T., & Ley, V. (2018). Leisure-time physical activity and prevalence of non-communicable pathologies and prescription medication in Spain. *PLoS ONE*, 13(1). <https://doi.org/10.1371/journal.pone.0191542>
- Glowacki, K., Arbour-Nicitopoulos, K., Burrows, M., Chesick, L., Heinemann, L., Irving, S., Lam, R. W., Macridis, S., Michalak, E., Scott, A., Taylor, A., & Faulkner, G. (2019). It's more than just a referral: Development of an evidence-informed exercise and depression toolkit. *Mental Health and Physical Activity*, 17, 100297. <https://doi.org/10.1016/j.mhpa.2019.100297>
- Hains-Monfette, G., Atoui, S., Needham Dancause, K., & Bernard, P. (2019). Device-Assessed Physical Activity and Sedentary Behaviors in Canadians with Chronic Disease(s): Findings from the Canadian Health Measures Survey. *Sports*, 7(5), 113. <https://doi.org/10.3390/sports7050113>
- Heil, D. P. (2006). Predicting Activity Energy Expenditure Using the Actical® Activity Monitor. *Research Quarterly for Exercise and Sport*, 77(1), 64–80. <https://doi.org/10.1080/02701367.2006.10599333>
- Jensen-Otsu, E., & Austin, G. (2015). Antidepressant Use is Associated with Increased Energy Intake and Similar Levels of Physical Activity. *Nutrients*, 7(11), 9662–9671. <https://doi.org/10.3390/nu7115489>
- Kennedy, S. H., Lam, R. W., McIntyre, R. S., Tourjman, S. V., Bhat, V., Blier, P., Hasnain, M., Jollant, F., Levitt, A. J.,

- MacQueen, G. M., McInerney, S. J., McIntosh, D., Milev, R. V., Müller, D. J., Parikh, S. V., Pearson, N. L., Ravindran, A. V., & Uher, R. (2016). Canadian Network for Mood and Anxiety Treatments (CANMAT) 2016 Clinical Guidelines for the Management of Adults with Major Depressive Disorder: Section 3. Pharmacological Treatments. *The Canadian Journal of Psychiatry*, *61*(9), 540–560. <https://doi.org/10.1177/0706743716659417>
- Kleppang, A. L., Hartz, I., Thurston, M., & Hagquist, C. (2019). Leisure-time physical activity among adolescents and subsequent use of antidepressant and hypnotic drugs: A prospective register linkage study. *European Child & Adolescent Psychiatry*, *28*(2), 177–188. <https://doi.org/10.1007/s00787-018-1160-x>
- Labotz, M., Wolff, T. K., Nakasone, K. T., Kimura, I. F., Hetzler, R. K., & Nichols, A. W. (2006). Selective Serotonin Reuptake Inhibitors and Rhabdomyolysis after Eccentric Exercise. *Medicine & Science in Sports & Exercise*, *38*(9), 1539–1542. <https://doi.org/10.1249/01.mss.0000227643.06478.c6>
- Lahti, J., Lallukka, T., Lahelma, E., & Rahkonen, O. (2013). Leisure-time physical activity and psychotropic medication: A prospective cohort study. *Preventive Medicine*, *57*(3), 173–177. <https://doi.org/10.1016/j.ypmed.2013.05.019>
- Laugesen, K., Olsen, M. S., Andersen, A. B. T., Frøslev, T., & Sørensen, H. T. (2013). In utero exposure to antidepressant drugs and risk of attention deficit hyperactivity disorder: A nationwide Danish cohort study. *BMJ Open*, *3*(9), e003507. <https://doi.org/10.1136/bmjopen-2013-003507>
- Loponen, T., Lallukka, T., Holstila, A., & Lahti, J. (2015). Joint association of physical activity and overweight with subsequent psychotropic medication: A register-linked follow-up study among employees. *BMC Public Health*, *15*(1), 1–7. <https://doi.org/10.1186/s12889-015-2346-5>
- Marlatt, M. W., Lucassen, P. J., & van Praag, H. (2010). Comparison of neurogenic effects of fluoxetine, duloxetine and running in mice. *Brain Research*, *1341*, 93–99. <https://doi.org/10.1016/j.brainres.2010.03.086>
- Matthews, C. E., Hagströmer, M., Pober, D. M., & Bowles, H. R. (2012). BEST PRACTICES FOR USING PHYSICAL ACTIVITY MONITORS IN POPULATION-BASED RESEARCH. *Medicine and Science in Sports and Exercise*, *44*(1 Suppl 1), S68–S76. <https://doi.org/10.1249/MSS.0b013e3182399e5b>
- Morkem, R., Barber, D., Williamson, T., & Patten, S. B. (2015). A Canadian Primary Care Sentinel Surveillance Network Study Evaluating Antidepressant Prescribing in Canada from 2006 to 2012. *The Canadian Journal of Psychiatry*, *60*(12), 564–570. <https://doi.org/10.1177/070674371506001207>
- Morres, I. D., Hatzigeorgiadis, A., Krommidas, C., Comoutos, N., Sideri, E., Ploumpidis, D., Economou, M., Papaioannou, A., & Theodorakis, Y. (2019). Objectively measured physical activity and depressive symptoms in adult outpatients diagnosed with major depression. Clinical perspectives. *Psychiatry Research*, *280*, 112489. <https://doi.org/10.1016/j.psychres.2019.112489>
- Nelson, J. C., & Spyker, D. A. (2017). Morbidity and Mortality Associated With Medications Used in the Treatment of Depression: An Analysis of Cases Reported to U.S. Poison Control Centers, 2000–2014. *American Journal of Psychiatry*, *174*(5), 438–450. <https://doi.org/10.1176/appi.ajp.2016.16050523>
- Newton, R. L., Han, H., Johnson, W. D., Hickson, D. A., Church, T. S., Taylor, H. A., Tudor-Locke, C., & Dubbert, P. M. (2013). Steps/day and metabolic

- syndrome in African American adults: The Jackson Heart Study. *Preventive Medicine*, 57(6), 855–859. <https://doi.org/10.1016/j.ypmed.2013.09.018>
- Olver, J. S., & Hopwood, M. J. (2013). Depression and physical illness. *Medical Journal of Australia*, 199(S6), S9–S12. <https://doi.org/10.5694/mja12.10597>
- Pratt, L. A., Brody, D. J., & Gu, Q. (2017). Antidepressant Use Among Persons Aged 12 and Over: United States, 2011–2014. *NCHS Data Brief*, 283, 1–8.
- Romain, A. J., Bernard, P., Stubbs, B., & Rosenbaum, S. (2018). Behavioral and Psychological Approaches in Exercise-Based Interventions in Severe Mental Illness. In *Exercise-Based Interventions for Mental Illness: Physical Activity as Part of Clinical Treatment* (pp. 189–217). Academic Press, Elsevier.
- Rotermann, M., Sanmartin, C., Hennessy, D., & Arthur, M. (2014). Prescription medication use by Canadians aged 6 to 79. *Health Reports*, 25(6), 3–9.
- Schuch, F. B., Vancampfort, D., Firth, J., Rosenbaum, S., Ward, P. B., Silva, E. S., Hallgren, M., Ponce De Leon, A., Dunn, A. L., Deslandes, A. C., Fleck, M. P., Carvalho, A. F., & Stubbs, B. (2018). Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. *American Journal of Psychiatry*, 175(7), 631–648. <https://doi.org/10.1176/appi.ajp.2018.17111194>
- Schuch, F., Vancampfort, D., Firth, J., Rosenbaum, S., Ward, P., Reichert, T., Bagatini, N. C., Bgeginski, R., & Stubbs, B. (2017). Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. *Journal of Affective Disorders*, 210, 139–150. <https://doi.org/10.1016/j.jad.2016.10.050>
- Sellers, C., Dall, P., Grant, M., & Stansfield, B. (2016). Validity and reliability of the activPAL3 for measuring posture and stepping in adults and young people. *Gait & Posture*, 43, 42–47. <https://doi.org/10.1016/j.gaitpost.2015.10.020>
- Serretti, A., & Mandelli, L. (2010). Antidepressants and Body Weight: A Comprehensive Review and Meta-Analysis. *The Journal of Clinical Psychiatry*, 71(10), 1259–1272. <https://doi.org/10.4088/JCP.09r05346blu>
- Shulman, K. I., Herrmann, N., & Walker, S. E. (2013). Current Place of Monoamine Oxidase Inhibitors in the Treatment of Depression. *CNS Drugs*, 27(10), 789–797. <https://doi.org/10.1007/s40263-013-0097-3>
- Silva, L. J. da, Azevedo, M. R., Matsudo, S., & Lopes, G. S. (2012). Association between levels of physical activity and use of medication among older women. *Cadernos de Saúde Pública*, 28(3), 463–471. <https://doi.org/10.1590/S0102-311X2012000300006>
- Simonton, A. J., Young, C. C., & Brown, R. A. (2018). Physical Activity Preferences and Attitudes of Individuals With Substance Use Disorders: A Review of the Literature. *Issues in Mental Health Nursing*, 39(8), 657–666. <https://doi.org/10.1080/01612840.2018.1429510>
- Skånland, S. S., & Cieślak-Pobuda, A. (2019). Off-label uses of drugs for depression. *European Journal of Pharmacology*, 865, 172732. <https://doi.org/10.1016/j.ejphar.2019.172732>
- Stamatakis, E., Hamer, M., & Primatesta, P. (2009). Cardiovascular medication, physical activity and mortality: Cross-sectional population study with ongoing mortality follow-up. *Heart*, 95(6), 448–453. <https://doi.org/10.1136/hrt.2008.152041>
- Stubbs, B., Vancampfort, D., Mänty, M., Svärd, A., Rahkonen, O., & Lahti, J. (2017). Bidirectional longitudinal relationship between leisure-time physical activity and psychotropic medication usage: A register linked follow-up study. *Psychiatry Research*, 247, 208–213. <https://doi.org/10.1016/j.psychres.2016.11.033>

- Sundbom, L. T., & Hedborg, K. (2019). Association between prescribed antidepressants and other prescribed drugs differ by gender: A nationwide register-based study in Sweden. *Nordic Journal of Psychiatry, 73*(1), 73–79. <https://doi.org/10.1080/08039488.2018.1536766>
- Tremblay, M. S., & Gorber, S. C. (2007). Canadian Health Measures Survey: Brief Overview. *Canadian Journal of Public Health / Revue Canadienne de Sante'e Publique, 98*(6), 453–456.
- Tudor-Locke, C., Han, H., Aguiar, E. J., Barreira, T. V., Schuna Jr, J. M., Kang, M., & Rowe, D. A. (2018). How fast is fast enough? Walking cadence (steps/min) as a practical estimate of intensity in adults: a narrative review. *British Journal of Sports Medicine, 52*(12), 776–788. <https://doi.org/10.1136/bjsports-2017-097628>
- Tudor-Locke, C., Johnson, W. D., & Katzmarzyk, P. T. (2011a). Relationship Between Accelerometer-Determined Steps/Day and Other Accelerometer Outputs in U.S. Adults. *Journal of Physical Activity and Health, 8*(3), 410–419. <https://doi.org/10.1123/jpah.8.3.410>
- Tudor-Locke, C., Johnson, W. D., & Katzmarzyk, P. T. (2011b). U.S. Population Profile of Time-Stamped Accelerometer Outputs: Impact of Wear Time. *Journal of Physical Activity and Health, 8*(5), 693–698. <https://doi.org/10.1123/jpah.8.5.693>
- Vancampfort, D., Firth, J., Schuch, F. B., Rosenbaum, S., Mugisha, J., Hallgren, M., Probst, M., Ward, P. B., Gaughran, F., De Hert, M., Carvalho, A. F., & Stubbs, B. (2017). Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: A global systematic review and meta-analysis. *World Psychiatry, 16*(3), 308–315. <https://doi.org/10.1002/wps.20458>
- Waller, K., Kaprio, J., Korhonen, T., Tuulio-Henriksson, A., & Kujala, U. M. (2016). Persistent leisure-time physical activity in adulthood and use of antidepressants: A follow-up study among twins. *Journal of Affective Disorders, 200*, 172–177. <https://doi.org/10.1016/j.jad.2016.04.036>
- Westerterp, K. R. (2009). Assessment of physical activity: A critical appraisal. *European Journal of Applied Physiology, 105*(6), 823–828. <https://doi.org/10.1007/s00421-009-1000-2>
- Wong, J., Motulsky, A., Abrahamowicz, M., Eguale, T., Buckeridge, D. L., & Tamblyn, R. (2017). Off-label indications for antidepressants in primary care: Descriptive study of prescriptions from an indication based electronic prescribing system. *BMJ, 356*. <https://doi.org/10.1136/bmj.j603>
- Wong, J., Motulsky, A., Eguale, T., Buckeridge, D. L., Abrahamowicz, M., & Tamblyn, R. (2016). Treatment Indications for Antidepressants Prescribed in Primary Care in Quebec, Canada, 2006-2015. *JAMA, 315*(20), 2230. <https://doi.org/10.1001/jama.2016.3445>
- Wong, J. N., McAuley, E., & Trinh, L. (2018). Physical activity programming and counseling preferences among cancer survivors: A systematic review. *International Journal of Behavioral Nutrition and Physical Activity, 15*(1), 48. <https://doi.org/10.1186/s12966-018-0680-6>
- Wong, S. L., Malaisson, E., Hammond, D., & Leatherdale, S. T. (2013). Secondhand Smoke Exposure Among Canadians: Cotinine and Self-Report Measures From the Canadian Health Measures Survey 2007–2009. *Nicotine & Tobacco Research, 15*(3), 693–700. <https://doi.org/10.1093/ntr/nts195>
- Yun, L., Fagan, M., Subramaniapillai, M., Lee, Y., Park, C., Mansur, R. B., McIntyre, R. S., & Faulkner, G. E. J. (2020). Are early increases in physical activity a behavioral marker for successful antidepressant treatment? *Journal of Affective Disorders, 260*, 287–291. <https://doi.org/10.1016/j.jad.2019.09.002>

Table 1

Descriptive statistics of adults of the Canadian Health Measures Survey (cycle 1,2,3) according to their antidepressant use

	Control (N≈22,046, 570)	Antidepressant (N≈2,261,180)	Univariate test	p
Sex (men) %	51.7	28.62	F = 39.39	7.548 ^{e-10}
Age (years) M(Se)	44.37 (0.20)	50.32 (0.92)	T = 6.12	1.924 ^{e-09}
BMI M(Se)	27.44(0.24)	28.58(0.45)	t = 2.38	0.04
Marital status (alone) %	32.88	42.22	F = 11.76	0.0007
Income %				
<\$15k	3.59	5.80	F = 1.78	0.10
\$15k-\$19,999k	2.07	4.18		
\$20k-\$29,999k	7.42	10.18		
\$30k-\$39,999k	10.00	8.04		
\$40k-\$49,999k	8.67	11.71		
\$50k-\$59,999k	8.83	9.47		
\$60k-\$79,999k	15.69	15.27		
\$80k-\$99,999k	12.70	10.29		
≥ \$100k	31.03	25.05		
Education %			F = 0.85	0.50
Lower than high school	7.72	4.90		
High school	10.04	11.16		
Work school	17.14	15.64		
College	8.06	10.43		
University <Bachelor	13.93	12.20		
University ≥ Bachelor	14.84	14.18		
Other	28.26	31.49		
Self-reported symptoms and disease				
Self-rated health compared to 1 year ago %			F = 2.80	0.095
Much better/somewhat/about the same	87.46	82.81		
Somewhat/much worse	12.54	17.19		
Self-reported mood disorder %	3.39	64.26	F = 1164.2	<2.2 ^{e-16}
Other physical or mental condition %	16.21	37.58	F = 82.97	<2.2 ^{e-16}
Psychosocial outcomes				
Self-rated quality of life %			F = 22.64	2.563 ^{e-06}
Fair/poor	4.42	14.15		
(Very)good/excellent	95.58	85.85		
Self-rated mental health %			F = 52.26	1.84 ^{e-12}
Fair/poor	3.54	19.35		
(Very)good/excellent	96.46	80.65		

Self-rated stress %			F = 18.20	2.384 ^{e-05}
Not at all/not very/a bit stressful	80.16	63.72		
Quite a bit/extremely stressful	19.84	36.28		
Smoking variables				
Smoking %	19.55	25.94	F = 2.99	0.08
Number of years smoked daily (former daily smokers) M(Se)	16.50(0.47)	19.18(1.24)	t = 2.14	0.03
Levels of cotinine M(Se)	235.29(14.67)	351.05(55.75)	t = 2.20	0.03
Characteristics of physical activity and sedentary behavior				
Acc wear time (h/day) M(Se)	13.15(0.05)	12.52(0.12)	T = -4.72	3.016 ^{e-06}
MVPA (min/day) M(Se)	22.45(0.80)	14.17(1.14)	t = -7.87	2.191 ^{e-14}
LPA (min/day) M(Se)	217.48(3.10)	194.47(5.76)	T = -3.68	0.0003
Steps (steps/day) M(Se)	8011.79(108.08)	6436.29(274.38)	T = -5.41	9.642 ^{e-08}
Sed (min/day) M(Se)	548.70(2.47)	542.38(5.66)	T = -1.03	0.30

Notes. BMI = Body Mass Index, Cig = Cigarette, Acc = Accelerometer, LPA = Light Physical Activity, MVPA = Moderate and Vigorous Physical Activity, Sed = Sedentary behavior. Group size are estimated from the weights and bootstraps provided by Statistics Canada.

Table 2

Physical activity and sedentary behaviors of Canadian adults according to their antidepressant use and the category of antidepressant used

	MVPA (min/day)			LPA (min/day)			Steps (/day)			SB (min/day)		
	M	Se	Md	M	Se	Md	M	Se	Md	M	Se	Md
HC	22.45	0.8 0	16.1	217.5	3.1	208.1	8012	108	7612	548.7	2.5	555.4
AD	14.17	1.1 4	9.6	194.5	5.8	180.7	6436	274	5714	542.4	5.7	545.1
SSRI	13.04	1.2 8	8.1	192.9	9.2	180.6	6216	375	5758	541.1	11.7	544.7
SNRI	15.76	2.8 1	10.4	207.8	13.6	202.2	6984	692	5776	531.2	11.0	518.9
Tri	16.07	2.9 0	8.5	203.1	11.9	182.9	7640	594	7816	572.2	16.3	560.4
Others	12.83	2.6 5	9.4	174.0	11.9	159.9	5749	555	5317	563.0	17.4	559.9

Notes. MVPA = Moderate and Vigorous Physical Activity, LPA = Light Physical Activity, SB = Sedentary behavior, HC = Healthy Controls, AD = Antidepressant Users, SSRI = Selective Serotonin Reuptake Inhibitor Users, SNRI = Serotonin-Norepinephrine Reuptake Inhibitor Users, Tri = Tricyclic antidepressant Users, Others = Other Antidepressant Users.

Supplementary files

1. Univariate analysis.....	17
1. Table S1—Descriptive statistics of adults in the Canadian Health Measures Survey (cycle 1,2,3) according to antidepressant classes used.....	17
2. Table S2- Descriptive statistics of adults in the Canadian Health Measures Survey (cycle 1,2,3) using only antidepressant medication.....	19
3. Table S3- Descriptive statistics of adults in the Canadian Health Measures Survey (cycle 1,2,3) using antidepressant medication and at least another psychotropic medication.....	21
2. Multivariate analysis.....	23
1 Table S4- Weighted ANCOVA table for use of antidepressant medication and MVPA.....	23
2 Table S5- Weighted ANCOVA table for use of antidepressant medication and LPA.....	24
3 Table S6- Weighted ANCOVA table for use of antidepressant medication and steps per day.....	25
4 Table S7- Weighted ANCOVA table for use of antidepressant medication and sedentary behaviors.....	26
5 Table S8- Weighted ANCOVA table for category of antidepressant used and MVPA.....	27
6 Table S9- Weighted ANCOVA table for category of antidepressant used and LPA.....	28
7 Table S10- Weighted ANCOVA table for category of antidepressant used and steps per day.....	29
8 Table S11- Weighted ANCOVA table for category of antidepressant used and sedentary behaviors.....	30
9 Table S12- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and MVPA.....	31
10 Table S13- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and LPA.....	32
11 Table S14- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and steps per day.....	33
12 Table S15- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and sedentary behaviors.....	34
13 Table S16- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and MVPA.....	35
14 Table S17- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and LPA.....	36
15 Table S18- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and steps per day.....	37
16 Table S19- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and sedentary behaviors.....	38

Univariate analysis

Table S1—Descriptive statistics of adults in the Canadian Health Measures Survey (cycle 1,2,3) according to antidepressant classes used

In this table, some data had to be omitted (e.g., income) to respect Statistics Canada's confidentiality requirements.

	Control (N≈20,754,500)	SSRI (N≈1,045,490)	SNRI (N≈536,460)	Tricycliques (N≈192,120)	Others (N≈274,160)
Sex (men) %	51.67	23.75	40.00	26.19	34.71
Age (years) M(Se)	44.37(0.20)	48.91(1.64)	48.57(1.87)	55.88(2.34)	56.40(1.51)
BMI M(Se)	27.44(0.24)	28.89(0.86)	29.06(1.12)	28.14(1.73)	27.46(0.97)
Marital status (alone) %	32.89	46.41	39.41	46.43	25.83
Education %					
High school and lower	17.76 25.21	14.88 25.62	10.39 29.44	20.37 21.97	19.05 19.05
Work school or college	13.94 14.84 28.25	17.36 19.83 22.31	11.69 22.08 26.41	11.90 9.84 35.93	13.10 15.48 33.33
University <Bachelor					
University ≥ Bachelor					
Other					
Self-reported symptoms and disease					
Self-rated health compared to 1 year ago %					
Much better/somewhat/ about the same	87.46 12.54	81.66 18.34	83.83 16.17	89.29 10.71	80.83 19.17
Somewhat/much worse					
Self-reported mood disorder %	3.39	66.67	80.85	36.47	35.00
Psychosocial outcomes					
Self-rated stress %					
Not at all/not very/a bit stressful	80.16 19.84	74.17 25.83	66.81 33.19	78.57 21.43	74.17 25.83
Quite a bit/extremely					

stressful

Smoking variables

Smoking %	19.55	22.50	20.00	28.54	24.71
Number of years smoked daily (former daily smokers)					
M(Se)	16.50(0.47)	19.21(2.55)	17.55(3.00)	20.33(2.44)	19.45(1.83)
Levels of cotinine M(Se)	235.29 (14.67)	387.61(94.47)	307.04(96.61)	299.80(99.27)	232.97(56.26)

Characteristics of physical activity and sedentary behavior

Acc wearing time (h/day)					
M(Se)	13.15(0.05)	12.45(0.18)	12.58(0.23)	13.19(0.18)	12.52(0.39)
MVPA (min/day) M(Se)	22.45(0.80)	13.04(1.28)	15.76(2.81)	16.07(2.90)	12.83(2.65)
t(p)		-5.57(4.29e-08)	-2.03(0.04)	-1.67(0.09)	-2.81(0.005)
LPA (min/day) M(Se)	217.48(3.10)	192.86(9.18)	207.84(13.55)	203.06(11.91)	174.02(11.94)
t(p)		-2.66(0.008)	-0.69(0.49)	-1.26(0.21)	-3.56(0.0004)
Steps (steps/day) M(Se)	8011.79 (108.08)	6216.15 (375.02)	6983.59 (692.39)	7640.42 (594.04)	5749.05 (554.80)
t(p)		-4.77(2.46e-06)	-1.42(0.16)	-0.62(0.54)	-4.15(3.96e-05)
Sed (min/day) M(Se)	548.70(2.47)	541.07(11.66)	531.16(11.01)	572.19(16.28)	563.02(17.40)
t(p)		-0.67(0.50)	-1.46(0.15)	1.47(0.14)	0.78(0.43)

Notes. BMI = Body Mass Index, Cig = Cigarette, Acc = Accelerometer, LPA = Light Physical Activity, MVPA = Moderate and Vigorous Physical Activity, Sed = Sedentary behavior. Group size are estimated from the weights and bootstraps provided by Statistics Canada.

Table S2- Descriptive statistics of adults in the Canadian Health Measures Survey (cycle 1,2,3) using only antidepressant medication

	Control (N≈22,573,400)	AD— (N≈1,734,352)	Univariate test	P
Sex (men) %	51.67	28.01	F = 20.40	7.832e-06
Age (years) M(Se)	44.37(0.20)	49.05(1.27)	T = 6.12	1.924e-09
BMI M(Se)	27.44(0.24)	28.68(0.55)	T = 3.58	0.0004
Worked at job last year %				
Yes	79.01	77.67	F = 0.10	0.90
No	18.87	20.22		
Study/retired	2.12	2.11		
Marital status (alone) %	32.88	41.46	F = 7.15	0.008
Income %				
<\$15k	3.59	3.78	F = 0.73	0.61
\$15k-\$19,999k	2.07	2.52		
\$20k-\$29,999k	7.42	8.54		
\$30k-\$39,999k	9.99	7.56		
\$40k-\$49,999k	8.67	13.03		
\$50k-\$59,999k	8.83	10.22		
\$60k-\$79,999k	15.69	17.37		
\$80k-\$99,999k	12.70	11.06		
≥ \$100k	31.05	25.91		
Education %			F = 0.60	0.67
Lower than high school	7.72	5.04		
High school	10.03	12.66		
Work school	17.14	13.81		
College	8.07	10.36		
University <Bachelor	13.94	12.52		
University ≥ Bachelor	14.84	16.40		
Other	28.26	29.21		
Self-reported symptoms and disease				
Self-rated health compared to 1 year ago %			F = 0.71	0.40
Much better/somewhat/about the same	87.47	84.15		
Somewhat/much worse	12.53	15.85		
Self-reported mood disorder %	3.39	60.31	F = 770.86	<2.2e-16
Other physical or mental condition %	16.22	37.03	F = 48.76	9.267e-12
Psychosocial outcomes				
Self-rated quality of life %			F = 6.53	>0.01
Fair/poor	4.42	11.24		
(Very)good/excellent	95.58	88.76		

Self-rated mental health %			F = 15.58	9.044e-05
Fair/poor	3.53	14.49		
(Very)good/excellent	96.47	85.51		
Self-rated stress %			F = 9.69	0.002
Not at all/not very/a bit stressful	80.16	66.01		
Quite a bit/extremely stressful	19.84	33.99		
Smoking variables				
Smoking %	19.55	25.21	F = 1.70	0.19
Number of years smoked daily (former daily smokers) M(Se)	16.50(0.47)	18.78(1.63)	T = 1.36	0.18
Levels of cotinine M(Se)	235.29(14.67)	346.40(62.31)	T = 1.80	0.07
Characteristics of physical activity and sedentary				
Acc wear time (h/day) M(Se)	13.15(0.05)	12.62(0.15)	T = -3.26	0.001
MVPA (min/day) M(Se)	22.45(0.80)	14.66(1.33)	T = -6.37	4.353e-10
LPA (min/day) M(Se)	217.48(3.10)	199.14(7.75)	T = -2.28	0.02
Steps (steps/day) M(Se)	8011.79(108.08)	6590.53(348.70)	T = -3.88	0.0001
Sed (min/day) M(Se)	548.70(2.47)	543.01(7.16)	T = -0.73	0.47

Notes. BMI = Body Mass Index, Cig = Cigarette, Acc = Accelerometer, LPA = Light Physical Activity, MVPA = Moderate and Vigorous Physical Activity, Sed = Sedentary behavior. Group size are estimated from the weights and bootstraps provided by Statistics Canada.

Table S3- Descriptive statistics of adults in the Canadian Health Measures Survey (cycle 1,2,3) using antidepressant medication and at least another psychotropic medication

	Control (N≈23,551,350)	AD+ (N≈756,410)	Univariate test	p
Sex (men) %	51.68	30.23	F = 19.20	1.435e-05
Age (years) M(Se)	44.37(0.20)	53.35(1.00)	T = 8.49	2.331e-16
BMI M(Se)	27.44(0.24)	28.34(0.83)	T = 1.25	0.21
Marital status (alone) %	32.88	44.05	F = 6.02	0.014
Income %			F =	
<\$15k	3.59	10.61		
\$15k-\$19,999k	2.07	8.04		
\$20k-\$29,999k	7.42	14.15		
\$30k-\$39,999k	10.00	9.32		
\$40k-\$49,999k	8.67	8.36		
\$50k-\$59,999k	8.82	7.72		
\$60k-\$79,999k	15.69	10.29		
\$80k-\$99,999k	12.69	8.36		
≥ \$100k	31.04	23.15		
Education %			F = 1.78	0.11
Lower than high school	7.72	4.59		
High school	10.04	7.54		
Work school	17.14	20.00		
College	8.07	10.82		
University <Bachelor	13.94	11.15		
University ≥ Bachelor	14.84	8.85		
Other	28.26	37.05		
Self-reported symptoms and disease				
Self-rated health compared to 1 year ago %			F = 5.24	0.02
Much better/somewhat/about the same	87.46	79.74		
Somewhat/much worse	12.54	20.26		
Self-reported mood disorder %	3.39	73.72	F = 708.74	<2.2e-16
Other physical or mental condition %	16.21	38.71	F = 47.24)	1.883e-11
Psychosocial outcomes				
Self-rated quality of life %			F = 53.43	1.072e-12
Fair/poor	4.42	21.15		
(Very)good/excellent	95.58	78.84		
Self-rated mental health %			F = 122.74	<2.2e-16
Fair/poor	3.53	30.77		
(Very)good/excellent	96.47	69.23		

Self-rated stress %			F =22.34	2.979e-06
Not at all/not very/a bit stressful	80.16	58.20		
Quite a bit/extremely stressful	19.84	41.80		
Smoking variables				
Smoking %	19.55	27.56	F = 2.21	0.14
Number of years smoked daily (former daily smokers) M(Se)	16.50(0.47)	20.18(1.81)	t = 2.10	0.04
Levels of cotinine M(Se)	235.29(14.67)	362.11(95.04)	T = 1.41	0.16
Characteristics of physical activity and sedentary				
Acc wear time (h/day) M(Se)	13.15(0.05)	12.29(0.16)	T = -4.98	8.954e-07
MVPA (min/day) M(Se)	22.45(0.80)	12.99(1.55)	T = -6.08	2.399e-09
LPA (min/day) M(Se)	217.48(3.10)	183.29(5.98)	T = -5.05	6.209e-07
Steps (steps/day) M(Se)	8011.79(108.08)	6067.33(291.56)	T = -6.49	2.086e-10
Sed (min/day) M(Se)	548.70(2.47)	540.87(7.81)	T = -1.04	0.30

Notes. BMI = Body Mass Index, Cig = Cigarette, Acc = Accelerometer, LPA = Light Physical Activity, MVPA = Moderate and Vigorous Physical Activity, Sed = Sedentary behavior. Group size are estimated from the weights and bootstraps provided by Statistics Canada.

Multivariate analysis

Table S4- Weighted ANCOVA table for use of antidepressant medication and MVPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
Antidepressant use	-0.14	0.07	-0.27 – -0.01	-2.16	0.03
Age	-0.01	0.001	-0.01 – -0.007	-8.35	7.59 ^{e-16}
BMI	-0.03	0.004	-0.04 – -0.03	-9.24	<2 ^{e-16}
Sex (women)	-0.25	0.04	-0.33 – -0.18	-6.55	1.55 ^{e-10}
Accelerometer wear time	0.08	0.009	0.06—0.10	8.66	<2 ^{e-16}
Worked last year					
Study/retired	-0.07	0.16	-0.38 – 0.24	-0.43	0.67
Working	0.21	0.06		3.51	0.0005
Seasons					
Spring	-0.02	0.07	-0.17 – 0.12	-0.32	0.75
Summer	-0.07	0.06	-0.19 – 0.05	-1.12	0.26
Winter	-0.08	0.10	-0.27 – 0.11	-0.86	0.39
Marital status (couple)	-0.20	0.04	-0.28 – -0.11	-4.41	0.00001
Levels of cotinine	-0.0002	0.00003	-0.0002 – -0.0001	-4.96	9.96 ^{e-07}
Education					
High school	0.06	0.07	-0.08 – 0.21	0.85	0.40
Work school	0.03	0.07	-0.11 – 0.17	0.46	0.65
College	0.13	0.08	-0.03 – 0.28	1.63	0.10
University <Bachelor	0.14	0.07	-0.001 – 0.29	1.95	0.052
Bachelor	0.33	0.10	0.13—0.52	3.32	0.0001
University> Bachelor	0.44	0.15	0.16—0.73	3.03	0.003
Other	-0.01	0.06	-0.13 – 0.10	-0.24	0.81
Household income					
\$15k-\$19,999k	-0.15	0.14	-0.43 – 0.13	-1.06	0.29
\$20k-\$29,999k	-0.20	0.10	-0.39 – -0.004	-2.00	0.046
\$30k-\$39,999k	-0.13	0.11	-0.35 – 0.08	-1.25	0.21
\$40k-\$49,999k	-0.16	0.09	-0.35 – 0.02	-1.76	0.08
\$50k-\$59,999k	-0.16	0.10	-0.37 – 0.04	-1.57	0.12
\$60k-\$79,999k	-0.13	0.09	-0.31 – 0.05	-1.43	0.15
\$80k-\$99,999k	-0.14	0.11	-0.35 – 0.07	-1.27	0.20
≥ \$100k	-0.07	0.09	-0.25 – 0.12	-0.72	0.47

Table S5- Weighted ANCOVA table for use of antidepressant medication and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
Antidepressant use	2.19	3.95	-5.55 – 9.92	0.55	0.58
Age	-0.66	0.10	-0.85 – -0.47	-6.85	2.35 ^{e-11}
BMI	-0.07	0.25	-0.55 – 0.41	-0.29	0.77
Sex (women)	-6.00	2.50	-10.91 – -1.09	-2.40	0.02
Accelerometer wear time	22.02	0.91	20.24—23.79	24.33	<2 ^{e-16}
Worked last year					
Study/retired	-11.57	4.85	-21.09 – -2.06	-2.39	0.02
Working	24.17	3.12	18.05—30.28	7.75	5.70 ^{e-14}
Seasons					
Spring	3.62	4.48	-5.17 – 12.41	0.81	0.42
Summer	13.25	5.22	3.03—23.48	2.54	0.01
Winter	-10.00	4.56	-18.95 – -1.06	-2.19	0.03
Marital status (couple)	15.89	2.52	10.95—20.84	6.30	6.91 ^{e-10}
Levels of cotinine	0.0005	0.003	-0.005 – 0.006	0.18	0.85
Education					
High school	20.97	7.47	6.32—35.62	2.81	0.006
Work school	5.13	7.43	-9.42 – 19.68	0.69	0.49
College	4.23	6.93	-9.36 – 17.83	0.61	0.54
University <Bachelor	-11.96	6.07	-23.86 – -0.07	-1.97	0.049
Bachelor	-16.00	6.20	-28.14 – -3.86	-2.58	>0.01
University> Bachelor	-17.84	7.36	-32.26 – -3.42	-2.43	0.02
Other	18.79	4.69	9.60—27.97	4.01	7.08 ^{e-05}
Household income					
\$15k-\$19,999k	3.65	7.95	-11.92 – 19.22	0.46	0.65
\$20k-\$29,999k	6.36	8.73	-10.75 – 23.47	0.73	0.47
\$30k-\$39,999k	10.71	8.35	-5.65 – 27.07	1.28	0.20
\$40k-\$49,999k	10.74	8.78	-6.47 – 27.96	1.22	0.22
\$50k-\$59,999k	9.62	8.09	-6.24 – 25.47	1.19	0.24
\$60k-\$79,999k	10.74	8.46	-5.85 – 27.32	1.27	0.21
\$80k-\$99,999k	7.46	8.36	-8.94 – 23.85	0.89	0.37
≥ \$100k	0.08	7.65	-14.90 – 15.07	0.01	0.99

Table S6- Weighted ANCOVA table for use of antidepressant medication and steps per day

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
Antidepressant use	-521.47	219.48	-951.65 – -91.29	-2.38	0.02
Age	-16.97	3.02	-22.89 – -11.05	-5.62	3.29 ^{e-08}
BMI	-78.91	9.86	-98.25 – -59.59	-8.00	9.49 ^{e-15}
Sex (women)	-850.13	104.19	-1054.35 – -645.92	-8.16	3.08 ^{e-15}
Accelerometer wear time	716.18	37.39	642.90—789.47	19.15	<2 ^{e-16}
Worked last year					
Study/retired	7.09	284.64	-550.79 – 564.96	0.03	0.98
Working	1036.76	179.57	684.81—1388.71	5.77	1.41 ^{e-08}
Seasons					
Spring	54.85	247.74	-430.71 – 540.41	0.22	0.82
Summer	207.59	218.47	-220.62 – 635.79	0.95	0.34
Winter	-625.58	206.11	-1029.55 – -221.61	-3.04	0.003
Marital status (couple)	-170.33	146.64	-457.74 – 117.07	-1.16	0.25
Levels of cotinine	-0.42	0.10	-0.62 – -0.22	-4.08	5.35 ^{e-05}
Education					
High school	666.20	286.45	104.76—1227.63	2.33	0.02
Work school	111.47	275.78	-429.05 – 652.00	0.40	0.69
College	-119.14	279.40	-666.75 – 428.47	-0.43	0.67
University <Bachelor	144.43	280.30	-404.94 – 693.80	0.52	0.61
Bachelor	395.12	341.59	-274.37 – 1064.62	1.16	0.25
University> Bachelor	463.30	434.78	-388.84 – 1315.45	1.07	0.29
Other	454.73	224.27	15.16—894.30	2.03	0.04
Household income					
\$15k-\$19,999k	-491.28	338.85	-1155.41 – 172.84	-1.45	0.15
\$20k-\$29,999k	-626.87	354.60	-1321.87 – 68.13	-1.77	0.08
\$30k-\$39,999k	48.58	324.97	-588.35 – 685.52	0.15	0.88
\$40k-\$49,999k	-358.17	348.62	-1041.45 – 325.10	-1.03	0.30
\$50k-\$59,999k	-83.37	314.41	-699.62 – 532.87	-0.27	0.79
\$60k-\$79,999k	-246.73	285.84	-806.97 – 313.50	-0.86	0.39
\$80k-\$99,999k	-113.15	275.50	-653.12 – 426.82	-0.41	0.68
≥ \$100k	-102.87	270.61	-633.26 – 427.52	-0.38	0.70

Table S7- Weighted ANCOVA table for use of antidepressant medication and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
Antidepressant use	-2.85	3.84	-10.37 – 4.68	-0.74	0.46
Age	0.82	0.10	0.62—1.02	8.14	3.59 ^{e-15}
BMI	0.82	0.24	0.34—1.29	3.39	0.0008
Sex (women)	11.62	2.52	6.68—16.56	4.61	5.27 ^{e-06}
Accelerometer wear time	35.65	0.79	34.10—37.19	45.18	<2 ^{e-16}
Worked last year					
Study/retired	12.35	4.72	3.09—21.61	2.61	0.009
Working	-27.51	3.75	-34.86 – -20.16	-7.34	9.65 ^{e-13}
Seasons					
Spring	-1.86	4.19	-10.08 – 6.36	-0.44	0.66
Summer	-11.19	5.41	-21.80 – -0.58	-2.07	0.04
Winter	13.98	4.59	4.98—22.98	3.04	0.002
Marital status (couple)	-10.83	2.78	-16.28 – -5.37	-3.89	0.0001
Levels of cotinine	0.003	0.003	-0.003 – 0.008	0.97	0.68
Education					
High school	-22.69	7.46	-37.31 – -8.07	-3.04	0.002
Work school	-3.89	7.53	-18.66 – 10.87	-0.52	0.61
College	-3.11	7.64	-18.09 – 11.87	-0.41	0.68
University <Bachelor	10.65	6.45	-1.99 – 23.29	1.65	0.10
Bachelor	8.21	6.50	-4.53 – 20.94	1.26	0.21
University> Bachelor	9.07	7.90	-6.42 – 24.56	1.15	0.25
Other	-16.45	4.99	-26.24 – -6.66	-3.29	0.001
Household income					
\$15k-\$19,999k	-2.95	8.59	-19.79 – 13.90	-0.34	0.73
\$20k-\$29,999k	-2.65	8.72	-19.74 – 14.44	-0.30	0.76
\$30k-\$39,999k	-7.42	8.95	-24.97 – 10.13	-0.83	0.41
\$40k-\$49,999k	-8.90	8.84	-26.22 – 8.42	-1.01	0.31
\$50k-\$59,999k	-6.98	7.95	-22.56 – 8.61	-0.88	0.38
\$60k-\$79,999k	-10.20	8.65	-27.16 – 6.76	-1.18	0.24
\$80k-\$99,999k	-4.27	8.29	-20.51 – 11.97	-0.52	0.61
≥ \$100k	0.71	7.62	-14.23 – 15.64	0.09	0.93

Table S8- Weighted ANCOVA table for category of antidepressant used and MVPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	-0.11	0.14	-0.38 – 0.16	-0.78	0.44
SSRI	-0.20	0.08	-0.38 – -0.05	-2.53	>0.01
Tri	-0.11	0.19	-0.48 – 0.25	-0.60	0.55
Other	-0.29	0.19	-0.66 – 0.09	-1.50	0.13
Age	-0.0095	0.001	-0.01 – -0.007	-8.47	3.17 ^{e-16}
BMI	-0.03	0.004	-0.04 – -0.03	-9.57	<2 ^{e-16}
Sex (women)	-0.25	0.04	-0.33 – -0.18	-6.62	1.00 ^{e-10}
Accelerometer wear time	0.08	0.009	0.06—0.10	8.56	<2 ^{e-16}
Worked last year					
Study/retired	-0.07	0.16	-0.38 – 0.24	-0.44	0.66
Working	0.21	0.06	0.10—0.32	3.73	0.0002
Seasons					
Spring	-0.02	0.07	-0.16 – 0.13	-0.25	0.80
Summer	-0.07	0.06	-0.19 – 0.06	-1.06	0.29
Winter	-0.08	0.10	-0.27 – 0.11	-0.84	0.40
Marital status (couple)	-0.20	0.05	-0.28 – -0.11	-4.64	4.46 ^{e-06}
Levels of cotinine	-0.0002	0.00003	-0.0002 – -0.00009	-4.88	1.43 ^{e-06}
Education					
High school	0.06	0.07	-0.09 – 0.20	0.73	0.46
Work school	0.03	0.07	-0.11 – 0.17	0.42	0.67
College	0.13	0.08	-0.02 – 0.29	1.65	0.10
University <Bachelor	0.14	0.07	-0.003 – 0.29	1.93	>0.05
Bachelor	0.32	0.10	0.14—0.51	3.38	0.0008
University> Bachelor	0.43	0.15	0.15—0.72	2.98	0.003
Other	0.02	0.06	-0.14 – 0.10	-0.36	0.72
Household income					
\$30k-\$59,999k	-0.02	0.06	-0.13 – 0.10	-0.27	0.79
\$60k-\$99,999k	0.003	0.07	-0.13 – 0.14	0.05	0.96
≥ \$100k	0.07	0.07	-0.06 – 0.20	1.02	0.31

Table S9- Weighted ANCOVA table for category of antidepressant used and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	4.24	9.82	-15.00 – 23.48	0.43	0.67
SSRI	6.22	5.67	-4.89 – 17.34	1.10	0.27
Tri	-5.71	8.67	-22.71 – 11.29	-0.66	0.51
Other	-13.08	6.99	-26.77 – 0.62	-1.87	0.06
Age	-0.65	0.10	-0.85 – -0.46	-6.46	2.57 ^{e-10}
BMI	-0.08	0.24	-0.56 – 0.40	-0.33	0.75
Sex (women)	-6.20	2.61	-11.31 – -1.09	-2.38	0.02
Accelerometer wear time	22.00	0.91	20.22—23.78	24.21	<2 ^{e-16}
Worked last year					
Study/retired	-11.62	4.68	-20.79 – -2.46	-2.49	0.01
Working	24.07	3.30	17.60—30.54	7.29	1.32 ^{e-12}
Seasons					
Spring	3.44	4.58	-5.53 – 12.41	0.75	0.45
Summer	13.26	5.31	2.85—23.67	2.50	0.01
Winter	-9.66	4.57	-18.61 – -0.71	-2.12	0.03
Marital status (couple)	16.33	2.49	11.44—21.21	6.55	1.50 ^{e-10}
Levels of cotinine	0.0004	0.003	-0.005 – 0.006	0.16	0.88
Education					
High school	20.98	7.47	6.34—35.62	2.81	0.005
Work school	5.04	7.39	-9.44 – 19.52	0.68	0.50
College	3.37	7.07	-10.49 – 17.23	0.48	0.63
University <Bachelor	-12.10	6.11	-24.09 – -0.12	-1.98	0.048
Bachelor	-16.51	6.36	-28.98 – -4.05	-2.60	<0.01
University> Bachelor	-18.26	7.31	-32.58 – -3.94	-2.50	>0.01
Other	18.68	4.78	9.31—28.05	3.91	0.0001
Household income					
\$30k-\$59,999k	6.84	4.76	-2.50 – 16.17	1.44	0.15
\$60k-\$99,999k	5.76	5.39	-12.63 – 5.88	1.07	0.29
≥ \$100k	-3.37	4.72	-12.63 – 5.88	-0.71	0.48

Table S10- Weighted ANCOVA table for category of antidepressant used and steps per day

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	-433.62	455.57	-1326.52 – 459.27	-0.95	0.34
SSRI	-678.29	346.20	-1356.82 – 0.24	-1.96	0.05
Tri	234.66	598.33	-938.04 – 1407.36	0.39	0.70
Other	-1086.45	486.14	-2039.27 – -133.63	-2.24	0.03
Age	-17.22	2.89	-22.89 – -11.55	-5.96	5.08 ^{e-09}
BMI	-79.66	9.59	-98.46 – -60.87	-8.31	1.04 ^{e-15}
Sex (women)	-854.71	104.76	-1060.03 – -649.39	-8.16	3.06 ^{e-15}
Accelerometer wear time	718.16	37.12	645.40—790.92	19.35	<2 ^{e-16}
Worked last year					
Study/retired	-28.89	285.88	-589.21 – 531.43	-0.10	0.92
Working	1023.70	176.48	677.81—1369.60	5.80	1.21 ^{e-08}
Seasons					
Spring	54.16	254.24	-444.14 – 552.46	0.21	0.83
Summer	205.86	222.00	-229.26 – 640.97	0.93	0.35
Winter	-622.99	207.07	-1028.83 – -217.14	-3.01	0.003
Marital status (couple)	-176.31	148.32	-467.01 – 114.39	-1.19	0.24
Levels of cotinine	-0.42	0.11	-0.62 – -0.21	-3.95	9.08 ^{e-05}
Education					
High school	643.63	289.46	76.29—1210.97	2.22	0.03
Work school	114.40	273.74	-422.12 – 650.92	0.42	0.68
College	-110.57	280.16	-659.66 – 438.53	-0.40	0.69
University <Bachelor	152.37	280.27	-396.95 – 701.69	0.54	0.59
Bachelor	388.59	338.38	-274.63 – 1051.82	1.15	0.25
University> Bachelor	457.88	432.87	-390.53 – 1306.30	1.06	0.29
Other	441.29	228.68	-6.93 – 889.50	1.93	0.05
Household income					
\$30k-\$59,999k	321.52	201.10	-72.62 – 715.66	1.60	0.11
\$60k-\$99,999k	250.03	181.09	-104.89 – 604.95	1.38	0.17
≥ \$100k	336.36	203.30	-62.09 – 734.82	1.66	0.10
Tukey				Z	
SNRI — Other	138.09	646.83		0.21	>0.99
SSRI — Other	194.13	572.82		0.34	0.99
Tri – Other	516.53	636.52		0.81	0.84
SSRI—SNRI	56.03	447.11		0.13	>0.99
Tri – SNRI	378.44	705.88		0.54	0.95
Tri – SSRI	322.41	559.03		0.58	0.94

Table S11- Weighted ANCOVA table for category of antidepressant used and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	-2.71	10.61	-23.50 – 18.08	-0.26	0.80
SSRI	-4.32	5.84	-15.76 – 7.13	-0.74	0.46
Tri	8.29	9.96	-11.23 – 27.81	0.83	0.41
Other	16.53	8.01	0.84 – 32.23	2.06	0.04
Age	0.83	0.10	0.63—1.04	7.96	1.31 ^{e-14}
BMI	0.81	0.24	0.35—1.27	3.44	0.0006
Sex (women)	10.72	2.74	5.35—16.09	3.91	0.0001
Accelerometer wear time	36.13	0.92	34.33—37.94	39.22	<2 ^{e-16}
Worked last year					
Study/retired	12.00	4.55	3.09—20.91	2.64	0.009
Working	-27.30	3.91	-34.96 – -19.64	-6.99	9.61 ^{e-12}
Seasons					
Spring	-2.38	4.40	-11.01 – 6.24	-0.54	0.58
Summer	-11.89	5.58	-22.82 – -0.96	-2.13	0.03
Winter	11.95	4.29	3.55—20.35	2.79	0.006
Marital status (couple)	-11.51	2.80	-17.01 – -6.02	-4.11	4.72 ^{e-05}
Levels of cotinine	0.003	0.003	-0.003 – 0.008	0.91	0.36
Education					
High school	-22.97	7.66	-37.99 – -7.96	-3.00	0.003
Work school	-4.14	7.59	-19.02 – 10.74	-0.55	0.59
College	-3.42	7.88	-18.86 – 12.02	-0.43	0.66
University <Bachelor	10.54	6.51	-2.22 – 23.30	1.62	0.11
Bachelor	8.41	6.64	-4.60 – 21.42	1.27	0.21
University > Bachelor	9.43	8.08	-6.41 – 25.26	1.17	0.24
Other	-16.74	5.15	-26.82 – -6.65	-3.25	0.001
Household income					
\$30k-\$59,999k	-6.47	4.93	-16.12 – 3.19	-1.31	0.19
\$60k-\$99,999k	-6.44	5.54	-17.30 – 4.43	-1.16	0.25
≥ \$100k	1.57	4.73	-7.69 – 10.84	0.33	0.74
Tukey				Z	
SNRI — Other	-20.74	13.52		-1.54	0.41
SSRI — Other	-19.06	10.78		-1.77	0.28
Tri – Other	-2.52	14.30		-0.18	>0.99
SSRI—SNRI	1.68	11.78		0.14	>0.99
Tri – SNRI	18.22	13.59		1.34	0.53
Tri – SSRI	16.55	10.02		1.65	0.34

Table S12- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and MVPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD—	-0.15	0.07	-0.29 – -0.002	-1.99	0.05
Age	-0.009	0.001	-0.01 – -0.007	-8.20	2.31 ^{e-15}
BMI	-0.03	0.004	-0.04 – -0.03	-8.98	<2 ^{e-16}
Sex (women)	-0.25	0.04	-0.33 – -0.18	-6.49	2.14 ^{e-10}
Accelerometer wear time	0.08	0.009	0.06—0.10	8.62	<2 ^{e-16}
Worked last year					
Study/retired	-0.04	0.16	-0.35 – 0.27	-0.25	0.80
Working	0.22	0.06	0.11—0.33	3.83	0.0001
Seasons					
Spring	-0.02	0.07	-0.16 – 0.12	-0.28	0.78
Summer	-0.07	0.06	-0.19 – 0.06	-1.08	0.28
Winter	-0.08	0.10	-0.27 – 1.09	-0.82	0.41
Marital status (couple)	-0.20	0.04	-0.29 – -0.12	-4.59	5.77 ^{e-06}
Levels of cotinine	-0.0002	0.00003	-0.0002 – -0.00009	-4.80	2.18 ^{e-06}
Education					
High school	0.06	0.07	-0.09 – 0.20	0.79	0.43
Work school	0.04	0.07	-0.10 – 0.17	0.51	0.61
College	0.14	0.08	-0.02 – 0.29	1.75	0.08
University <Bachelor	0.15	0.07	0.0009—0.29	1.97	<0.05
Bachelor	0.32	0.10	0.13—0.51	3.35	0.0009
University> Bachelor	0.44	0.15	0.15—0.73	2.97	0.003
Other	-0.01	0.06	-0.13 – 0.10	-0.20	0.84
Household income					
\$15k-\$19,999k	-0.16	0.13	-0.40 – 0.09	-1.27	0.21
\$20k-\$29,999k	-0.18	0.10	-0.38 – 0.02	-1.81	0.07
\$30k-\$39,999k	-0.12	0.11	-0.33 – 0.09	-1.10	0.27
\$40k-\$49,999k	-0.15	0.10	-0.34 – 0.05	-1.48	0.14
\$50k-\$59,999k	-0.15	0.11	-0.35 – 0.06	-1.39	0.16
\$60k-\$79,999k	-0.11	0.09	-0.30 – 0.07	-1.19	0.24
\$80k-\$99,999k	-0.12	0.11	-0.33 – 0.09	-1.10	0.27
≥ \$100k	-0.05	0.10	-0.24 – 0.14	-0.53	0.59

Table S13- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD—	3.18	4.88	-6.39 – 12.74	0.65	0.52
Age	-0.65	0.10	-8.49 – -0.46	-6.54	1.61 ^{e-10}
BMI	-0.11	0.25	-6.06 – 0.38	-0.44	0.66
Sex (women)	-6.35	2.63	-11.50 – -1.20	-2.42	0.02
Accelerometer wear time	22.09	0.93	20.27—23.92	23.72	<2 ^{e-16}
Worked last year					
Study/retired	-12.70	4.88	-22.26 – -3.14	-2.60	0.01
Working	24.09	3.22	17.78—30.40	7.48	3.61 ^{e-13}
Seasons					
Spring	3.72	4.58	-5.26 – 12.71	0.81	0.42
Summer	13.62	5.28	3.26—23.97	2.58	0.01
Winter	-9.51	4.57	-18.47 – -0.55	-2.08	0.04
Marital status (couple)	16.38	2.59	11.30—21.45	6.32	5.96 ^{e-10}
Levels of cotinine	0.0006	0.003	-0.005 – 0.006	0.22	0.82
Education					
High school	21.47	7.73	6.32—36.62	2.78	0.006
Work school	4.93	7.67	-10.10 – 19.96	0.64	0.52
College	3.78	7.29	-10.50 – 18.06	0.52	0.60
University <Bachelor	-11.95	6.30	-24.30 – 0.41	-1.90	0.06
Bachelor	-16.34	6.25	-28.58 – -4.09	-2.62	<0.01
University> Bachelor	-18.08	7.50	-32.79 – -3.37	-2.41	0.02
Other	18.72	4.94	9.04—28.40	3.79	0.0002
Household income					
\$15k-\$19,999k	4.19	8.21	-11.91 – 20.29	0.51	0.61
\$20k-\$29,999k	7.78	9.26	-10.37 – 25.93	0.84	0.40
\$30k-\$39,999k	11.86	9.00	-5.78 – 29.50	1.32	0.19
\$40k-\$49,999k	11.40	9.55	-7.32 – 30.11	1.19	0.23
\$50k-\$59,999k	10.72	8.79	-6.51 – 27.95	1.22	0.22
\$60k-\$79,999k	11.88	9.16	-6.06 – 29.83	1.30	0.19
\$80k-\$99,999k	8.28	9.02	-9.39 – 25.95	0.92	0.36
≥ \$100k	0.38	8.16	-15.61 – 16.37	0.05	0.96

Table S14- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and steps per day

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD—	-598.00	266.66	-1120.64 – -75.37	-2.24	0.03
Age	-16.11	2.99	-21.97 – -10.25	-5.39	1.14 ^{e-07}
BMI	-78.79	10.09	-98.57 – -59.01	-7.81	3.79 ^{e-14}
Sex (women)	-857.32	109.14	-1071.22 – -643.42	-7.86	2.71 ^{e-14}
Accelerometer wear time	720.83	37.77	646.80—794.86	19.09	<2 ^{e-16}
Worked last year					
Study/retired	6.12	299.63	-581.13 – 593.38	0.02	0.98
Working	1054.01	177.08	706.93—1401.09	5.95	5.17 ^{e-09}
Seasons					
Spring	43.05	252.63	-452.10 – 538.20	0.17	0.86
Summer	212.09	231.28	-241.22 – 665.40	0.92	0.36
Winter	-612.23	209.93	-1023.68 – -200.77	-2.92	0.004
Marital status (couple)	-187.33	150.28	-481.87 – 107.22	-1.25	0.21
Levels of cotinine	-0.41	0.10	-0.61 – -0.20	-3.92	0.0001
Education					
High school	686.81	289.25	119.89—1253.74	2.37	0.02
Work school	126.03	280.31	-423.36 – 675.42	0.45	0.65
College	-106.25	278.20	-651.51 – 439.01	-0.38	0.70
University <Bachelor	167.19	287.80	-396.90 – 731.28	0.58	0.56
Bachelor	398.86	337.38	-262.40 – 1060.11	1.18	0.24
University > Bachelor	474.73	430.57	-369.17 – 1318.63	1.10	0.27
Other	458.63	231.04	5.81—911.46	1.99	0.05
Household income					
\$15k-\$19,999k	-400.01	349.00	-1084.04 – 284.02	-1.15	0.25
\$20k-\$29,999k	-601.35	367.50	-1321.64 – 118.95	-1.64	0.10
\$30k-\$39,999k	121.54	332.50	-530.15 – 773.22	0.37	0.71
\$40k-\$49,999k	-306.35	373.59	-1038.56 – 425.87	-0.82	0.41
\$50k-\$59,999k	-15.10	330.20	-662.27 – 632.08	-0.05	0.96
\$60k-\$79,999k	-181.17	301.64	-772.37 – 410.04	-0.60	0.55
\$80k-\$99,999k	-58.07	283.50	-613.71 – 497.57	-0.21	0.84
≥ \$100k	-57.03	275.39	-596.80 – 482.73	-0.21	0.84

Table S15- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD—	-4.35	4.88	-13.92 – 5.22	-0.89	0.37
Age	0.81	0.10	0.61—1.01	7.85	2.80 ^{e-14}
BMI	0.85	0.25	0.36—1.34	3.45	0.0006
Sex (women)	12.27	2.63	7.12—17.42	4.67	3.92 ^{e-06}
Accelerometer wear time	35.52	0.81	33.93—37.10	43.94	<2 ^{e-16}
Worked last year					
Study/retired	13.30	4.77	3.95—22.66	2.79	0.006
Working	-27.97	3.89	-35.59 – -20.35	-7.20	2.48 ^{e-12}
Seasons					
Spring	-1.30	4.38	-9.88 – 7.28	-0.30	0.77
Summer	-10.83	5.71	-22.01 – 0.36	-1.90	0.06
Winter	14.08	4.64	4.99—23.18	3.04	0.003
Marital status (couple)	-11.46	2.91	-17.16 – -5.76	-3.94	9.33 ^{e-05}
Levels of cotinine	0.002	0.003	-0.004 – 0.007	0.51	0.61
Education					
High school	-22.83	7.67	-37.85 – -7.80	-2.98	0.003
Work school	-3.55	7.76	-18.76 – 11.67	-0.46	0.65
College	-2.68	7.83	-18.02 – 12.66	-0.34	0.73
University <Bachelor	10.62	6.70	-2.51 – 23.75	1.59	0.11
Bachelor	8.56	6.50	-4.18 – 21.31	1.32	0.19
University> Bachelor	9.49	7.93	-6.06 – 25.04	1.20	0.23
Other	-16.77	5.20	-26.96 – -6.57	-3.22	0.001
Household income					
\$15k-\$19,999k	-3.17	8.56	-19.94 – 13.61	-0.37	0.71
\$20k-\$29,999k	-4.12	9.20	-22.14 – 13.90	-0.45	0.65
\$30k-\$39,999k	-8.69	9.46	-27.24 – 9.86	-0.92	0.36
\$40k-\$49,999k	-9.77	9.47	-28.32 – 8.78	-1.03	0.30
\$50k-\$59,999k	-10.31	8.63	-27.24 – 6.61	-1.20	0.23
\$60k-\$79,999k	-11.60	9.24	-29.72 – 6.51	-1.26	0.21
\$80k-\$99,999k	-5.22	8.75	-22.37 – 11.93	-0.60	0.55
≥ \$100k	0.21	7.93	-15.33 – 15.76	0.03	0.98

Table S16- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and MVPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD+	-0.15	0.10	-0.34 – 0.04	-1.52	0.13
Age	-0.009	0.001	-0.01 – -0.007	-8.62	<2 ^{e-16}
BMI	-0.03	0.004	-0.04 – -0.03	-9.19	<2 ^{e-16}
Sex (women)	-0.25	0.04	-0.33 – -0.17	-6.19	1.30 ^{e-09}
Accelerometer wear time	0.08	0.01	0.06—0.10	8.73	<2 ^{e-16}
Worked last year					
Study/retired	-0.10	0.17	-0.43 – 0.23	-0.59	0.56
Working	0.20	0.06	0.08—0.32	3.32	0.001
Seasons					
Spring	-0.01	0.07	-0.16 – 0.14	-0.16	0.88
Summer	-0.05	0.06	-0.18 – 0.07	-0.86	0.39
Winter	-0.06	0.10	-0.25 – 0.12	-0.67	0.50
Marital status (couple)	-0.20	0.04	-0.29 – -0.11	-4.53	7.61 ^{e-06}
Levels of cotinine	-0.0002	0.00003	-0.0002 – -0.00009	-4.58	6.07 ^{e-06}
Education					
High school	0.06	0.08	-0.09 – 0.21	0.76	0.45
Work school	0.02	0.07	-0.12 – 0.16	0.30	0.77
College	0.12	0.08	-0.04 – 0.27	1.47	0.14
University <Bachelor	0.13	0.08	-0.02 – 0.28	1.71	0.09
Bachelor	0.31	0.10	0.12—0.51	3.18	0.002
University> Bachelor	0.44	0.14	0.16—0.71	3.10	0.002
Other	-0.04	0.06	-0.15 – 0.08	-0.63	0.53
Household income					
\$15k-\$19,999k	-0.11	0.14	-0.38 – 0.16	-0.82	0.41
\$20k-\$29,999k	-0.20	0.10	-0.40 – -0.01	-2.06	0.04
\$30k-\$39,999k	-0.13	0.11	-0.35 – 0.09	-1.12	0.26
\$40k-\$49,999k	-0.15	0.10	-0.34 – 0.04	-1.55	0.12
\$50k-\$59,999k	-0.15	0.10	-0.34 – 0.05	-1.43	0.15
\$60k-\$79,999k	-0.14	0.09	-0.32 – 0.04	-1.56	0.11
\$80k-\$99,999k	-0.14	0.11	-0.35 – 0.08	-1.26	0.21
≥ \$100k	-0.06	0.09	-0.24 – 0.13	-0.61	0.54

Table S17- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD+	1.38	4.92	-8.26 – 11.01	0.28	0.78
Age	-0.67	0.10	-0.86 – -0.48	-6.90	1.70 ^{e-11}
BMI	0.07	0.30	-0.51 – 0.65	-0.23	0.82
Sex (women)	-6.16	2.41	-10.88 – -1.43	-2.55	0.01
Accelerometer wear time	22.38	0.74	20.93—23.83	30.29	<2 ^{e-16}
Worked last year					
Study/retired	-13.40	5.09	-23.38 – -3.41	-2.63	0.009
Working	24.25	3.21	17.96—30.54	7.56	2.14 ^{e-13}
Seasons					
Spring	3.03	5.05	-6.87 – 12.93	0.60	0.55
Summer	13.32	5.65	2.25—24.38	2.36	0.02
Winter	-11.34	5.06	-21.25 – -1.42	-2.24	0.03
Marital status (couple)	17.93	2.75	12.55—23.32	6.52	1.77 ^{e-10}
Levels of cotinine	0.0008	0.003	-0.005 – 0.006	0.30	0.76
Education					
High school	19.20	6.32	6.81—31.59	3.04	0.003
Work school	5.52	7.30	-8.79 – 19.83	0.76	0.45
College	3.35	6.14	-8.68 – 15.38	0.55	0.59
University <Bachelor	-9.93	6.17	-22.01 – 2.16	-1.61	0.11
Bachelor	-15.67	6.58	-28.57 – -2.77	-2.38	0.02
University > Bachelor	-15.61	7.56	-30.43 – -0.78	-2.06	0.04
Other	19.16	4.64	10.07—28.25	4.13	4.29 ^{e-05}
Household income					
\$15k-\$19,999k	4.59	8.16	-11.41 – 20.58	0.56	0.57
\$20k-\$29,999k	6.79	9.00	-10.86 – 24.44	0.75	0.45
\$30k-\$39,999k	12.40	8.99	-5.23 – 30.03	1.38	0.17
\$40k-\$49,999k	11.91	9.65	-7.01 – 30.83	1.23	0.22
\$50k-\$59,999k	11.62	8.32	-4.69 – 27.94	1.40	0.16
\$60k-\$79,999k	10.81	8.95	-6.73 – 28.36	1.21	0.23
\$80k-\$99,999k	8.01	8.93	-9.49 – 25.51	0.90	0.37
≥ \$100k	-0.92	7.92	-16.35 – 14.60	-0.12	0.91

Table S18- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and steps per day

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD+	-317.44	219.70	-748.04 – 113.15	-1.45	0.15
Age	-18.15	3.11	-24.23 – -12.06	-5.84	9.66 ^{e-09}
BMI	-70.44	10.24	-90.50 – -50.37	-6.88	1.90 ^{e-11}
Sex (women)	-843.08	106.32	-1051.46 – -634.70	-7.93	1.60 ^{e-14}
Accelerometer wear time	727.49	37.75	653.50—801.48	19.27	<2 ^{e-16}
Worked last year					
Study/retired	-29.07	275.45	-568.93 – 510.80	-0.11	0.92
Working	1040.11	189.80	668.10—1412.12	5.48	6.94 ^{e-08}
Seasons					
Spring	-38.95	256.50	-541.67 – 463.78	-0.15	0.88
Summer	287.16	231.94	-167.43 – 741.74	1.24	0.22
Winter	-544.15	210.67	-957.06 – -131.24	-2.58	0.01
Marital status (couple)	-111.65	152.46	-410.46 – 187.17	-0.73	0.46
Levels of cotinine	-0.39	-0.11	-0.61 – -0.16	-3.40	0.0007
Education					
High school	661.41	281.96	108.77—1214.04	2.35	0.02
Work school	24.01	282.83	-530.33 – 578.35	0.09	0.93
College	-178.82	288.68	-744.63 – 386.99	-0.62	0.54
University <Bachelor	110.99	281.23	-440.22 – 662.19	0.40	0.69
Bachelor	342.00	332.00	-308.70 – 992.70	1.03	0.30
University> Bachelor	587.57	393.12	-182.92 – 1358.06	1.50	0.14
Other	327.97	232.18	-127.10 – 783.03	1.41	0.16
Household income					
\$15k-\$19,999k	-466.63	344.21	-1141.27 – 208.02	-1.36	0.18
\$20k-\$29,999k	-493.87	365.26	-1209.75 – 222.02	-1.35	0.18
\$30k-\$39,999k	75.97	348.45	-606.99 – 758.92	0.22	0.83
\$40k-\$49,999k	-254.57	328.33	-898.09 – 388.95	-0.78	0.44
\$50k-\$59,999k	73.42	289.70	-494.38 – 641.22	0.25	0.80
\$60k-\$79,999k	-311.56	300.22	-899.97 – 276.85	-1.04	0.30
\$80k-\$99,999k	-117.24	291.16	-687.90 – 453.42	-0.40	0.69
≥ \$100k	-73.78	266.63	-596.35 – 448.80	-0.28	0.78

Table S19- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education and total household income)

	Estimate	SE	95% CI	t	p
AD+	0.65	5.46	-10.06 – 11.36	0.12	0.91
Age	0.83	0.10	0.63—1.03	8.03	7.70 ^{e-15}
BMI	0.66	0.28	0.11—1.21	2.37	0.02
Sex (women)	11.28	2.49	6.39—16.16	4.53	7.61 ^{e-06}
Accelerometer wear time	35.76	0.79	34.21—37.31	45.22	<2 ^{e-16}
Worked last year					
Study/retired	14.57	4.94	4.89—24.26	2.95	0.003
Working	-27.91	3.95	-35.65 – -20.18	-7.07	5.49 ^{e-12}
Seasons					
Spring	-1.45	5.01	-11.27 – 8.38	-0.29	0.77
Summer	-11.54	5.99	-23.29 – 0.21	-1.92	>0.05
Winter	13.16	4.68	4.00—22.33	2.81	0.005
Marital status (couple)	-12.52	2.97	-18.35 – -6.69	-4.21	3.08 ^{e-05}
Levels of cotinine	0.002	0.003	-0.004 – 0.008	0.56	0.57
Education					
High school	-23.45	4.54	-38.23 – -8.67	-3.11	0.002
Work school	-4.54	7.55	-19.35 – 10.26	-0.60	0.55
College	-2.81	6.78	-16.09 – 10.47	-0.42	0.68
University <Bachelor	8.64	6.49	-4.08 – 21.35	1.33	0.18
Bachelor	7.50	6.65	-5.53 – 20.53	1.13	0.26
University > Bachelor	6.34	7.98	-9.29 – 21.98	0.80	0.43
Other	-17.47	4.99	-27.26 – -7.69	-3.50	0.0005
Household income					
\$15k-\$19,999k	-4.27	8.69	-21.30 – 12.75	-0.49	0.62
\$20k-\$29,999k	-3.03	8.84	-20.35 – 14.30	-0.34	0.73
\$30k-\$39,999k	-8.99	9.46	-27.54 – 9.56	-0.95	0.34
\$40k-\$49,999k	-9.87	9.57	-28.62 – 8.88	-1.03	0.30
\$50k-\$59,999k	-11.47	8.10	-27.54 – 4.40	-1.42	0.16
\$60k-\$79,999k	-10.31	9.14	-28.23 – 7.61	-1.13	0.26
\$80k-\$99,999k	-4.97	8.85	-22.32 – 12.38	-0.56	0.58
≥ \$100k	0.40	7.81	-14.91 – 15.72	0.05	0.96

Table S20- Weighted ANCOVA table for use of antidepressant medication and MVPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
Antidepressant use	-0.12	0.06	-0.24 – 0.01	-1.78	0.08
Age	-0.01	0.001	-0.01 - -0.007	-8.40	5.31 ^{e-16}
BMI	-0.03	0.004	-0.4 - -0.02	-8.57	<2 ^{e-16}
Sex (women)	-0.25	0.04	-0.33 - -0.18	-6.65	8.31 ^{e-11}
Accelerometer wear time	0.08	0.009	0.06 – 0.29	8.71	<2 ^{e-16}
Worked last year					
Study/retired	-0.07	0.16	-0.38 – 0.29	-0.43	0.67
Working	0.18	0.06	0.07 – 0.10	3.14	0.002
Seasons					
Spring	-0.02	0.07	-0.17 – 0.12	-0.33	0.74
Summer	-0.07	0.06	-0.19 – 0.06	-1.04	0.30
Winter	-0.08	0.10	-0.27 – 0.11	-0.84	0.40
Marital status (couple)	-0.20	0.04	-0.28 - -0.11	-4.54	0.00001
Levels of cotinine	-0.0001	0.00003	-0.0002 - -0.00009	-4.74	2.83 ^{e-06}
Education					
High school	0.06	0.07	-0.09 - 0.20	0.76	0.45
Work school	0.03	0.07	-0.11 – 0.17	0.46	0.65
College	0.12	0.08	-0.4 – 0.28	1.51	0.13
University <Bachelor	0.14	0.07	-0.01 – 0.28	1.82	0.07
Bachelor	0.32	0.10	0.13 – 0.51	3.26	0.001
University> Bachelor	0.43	0.15	0.15 – 0.72	2.96	0.003
Other	-0.01	0.06	-0.13 – 0.11	-0.16	0.88
Household income					
\$30k-\$59,999k	-0.04	0.06	-0.15 – 0.08	-0.59	0.56
\$60k-\$99,999k	-0.02	0.07	-0.15 – 0.12	-0.26	0.79
≥ \$100k	0.04	0.07	-0.09 – 0.17	0.67	0.51
Self-reported mental health	0.02	0.11	-0.20 – 0.24	0.19	0.85
Self-reported physical health	-0.35	0.07	-0.48 - -0.21	-5.14	4.10 ^{e-07}

Table S21- Weighted ANCOVA table for use of antidepressant medication and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
Antidepressant use	5.36	3.37	-1.24 – 11.96	1.59	0.11
Age	-0.66	0.10	-0.85 - -0.47	-6.86	2.23 ^{e-11}
BMI	-0.04	0.24	-0.43 – 0.50	0.15	0.88
Sex (women)	-6.06	2.49	-10.94 - -1.17	-2.43	0.02
Accelerometer wear time	21.86	0.90	20.10 – 23.63	24.31	<2 ^{e-16}
Worked last year					
Study/retired	-10.46	4.32	-18.92 - -2.00	-2.42	0.02
Working	22.84	3.11	10.84 – 20.63	7.34	9.07 ^{e-13}
Seasons					
Spring	3.79	4.50	-5.02 – 12.60	0.84	0.40
Summer	13.70	5.22	3.47 – 23.93	2.63	0.009
Winter	-9.16	4.66	-18.30 - -0.02	-1.97	0.049
Marital status (couple)	15.73	2.50	10.84 – 20.63	6.30	6.72 ^{e-10}
Levels of cotinine	0.001	0.003	-0.004 – 0.007	0.49	0.62
Education					
High school	21.38	7.33	7.01 – 35.74	2.92	0.004
Work school	5.85	7.33	-8.51 – 20.21	0.80	0.43
College	4.26	6.76	-8.99 – 17.51	0.63	0.53
University <Bachelor	-12.18	6.00	-23.94 - -0.42	-20.3	0.04
Bachelor	-16.05	6.20	-28.19 - -3.90	-2.59	0.099
University> Bachelor	-17.13	7.36	-31.56 - -2.70	-2.33	0.02
Other	20.03	4.79	10.63 – 29.42	4.18	3.52 ^{e-05}
Household income					
\$30k-\$59,999k	4.76	4.45	-3.96 – 13.49	1.07	0.29
\$60k-\$99,999k	3.40	5.07	-6.54 – 13.33	0.67	0.50
≥ \$100k	-6.23	4.44	-14.93 – 2.46	-1.41	0.16
Self-reported mental health	-10.45	6.17	-22.54 – 1.63	-1.70	0.09
Self-reported physical health	-16.18	4.16	-24.33 – 8.02	-3.89	0.0001

Table S22- Weighted ANCOVA table for use of antidepressant medication and steps per day (controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
Antidepressant use	-356.18	217.73	-782.93 – 70.57	-1.64	0.10
Age	-17.70	2.91	-23.40 – 12.00	-6.09	2.37 ^{e-09}
BMI	-73.87	9.97	-93.41 - -54.33	-7.41	5.84 ^{e-13}
Sex (women)	-852.06	102.46	-1052.88 - -651.24	-8.32	9.68 ^{e-16}
Accelerometer wear time	711.32	37.03	638.74 – 783.91	19.21	<2 ^{e-16}
Worked last year					
Study/retired	-2.87	280.20	-552.05 – 546.32	-0.01	0.99
Working	953.98	174.69	611.60 – 1296.36	5.46	7.64 ^{e-08}
Seasons					
Spring	55.20	249.90	-434.59 – 544.99	0.22	0.83
Summer	232.46	210.89	-180.88 – 645.80	1.10	0.27
Winter	-609.28	203.93	-1008.97 - -209.59	-2.99	0.003
Marital status (couple)	-192.71	151.80	-490.23 – 104.81	-1.27	0.20
Levels of cotinine	-0.37	0.10	-0.28 - -0.17	-3.59	0.0004
Education					
High school	669.47	287.94	105.12 – 1233.81	2.33	0.02
Work school	139.28	268.24	-386.45 – 665.02	0.52	0.60
College	-104.95	274.25	-642.48 – 432.58	-0.38	0.70
University <Bachelor	129.42	275.40	-410.35 – 669.18	0.47	0.64
Bachelor	395.18	332.64	-256.77 – 1047.13	1.19	0.24
University> Bachelor	494.06	415.65	-320.59 – 1308.72	1.19	0.24
Other	507.91	228.94	59.21 – 956.61	2.22	0.03
Household income					
\$30k-\$59,999k	219.83	197.57	-167.39 – 607.06	1.11	0.27
\$60k-\$99,999k	139.57	170.45	-194.51 – 473.66	0.82	0.41
≥ \$100k	199.35	190.38	-173.79 – 572.49	1.05	0.30
Self-reported mental health	-377.95	385.42	-1133.36 – 377.45	-0.98	0.33
Self-reported physical health	-952.45	178.07	-1301.46 - -603.43	-5.35	1.38 ^{e-07}

Table S23- Weighted ANCOVA table for use of antidepressant medication and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
Antidepressant use	-6.51	3.42	-13.21 – 0.18	-1.91	0.06
Age	0.83	0.10	0.63 – 1.02	8.24	1.66 ^{e-15}
BMI	0.68	0.23	0.22 – 1.14	2.91	0.004
Sex (women)	11.59	2.50	6.68 – 16.50	4.63	4.76 ^{e-06}
Accelerometer wear time	35.86	0.77	34.36 – 37.36	46.82	<2 ^{e-16}
Worked last year					
Study/retired	11.08	4.20	2.84 – 19.32	2.64	0.009
Working	-25.75	3.70	-33.00 – -18.50	-6.96	1.15 ^{e-11}
Seasons					
Spring	-2.06	4.18	-10.25 – 6.12	-0.49	0.62
Summer	-11.69	5.37	-22.23 – -1.16	-2.18	0.03
Winter	13.02	4.62	3.96 – 22.07	2.82	0.005
Marital status (couple)	-10.55	2.78	-16.01 – -5.10	-3.79	0.0002
Levels of cotinine	0.002	0.003	-0.004 – 0.007	0.60	0.55
Education					
High school	-23.19	7.37	-37.63 – -8.75	-3.15	0.002
Work school	-4.86	7.46	-19.48 – 9.75	-0.65	0.51
College	-3.24	7.47	-17.88 – 11.40	-0.43	0.66
University <Bachelor	10.75	6.40	-1.78 – 23.29	1.68	0.09
Bachelor	8.16	6.44	-4.46 – 20.78	1.27	0.21
University > Bachelor	8.22	7.85	-7.16 – 23.59	1.05	0.30
Other	-18.01	5.13	-28.07 – -7.96	-3.51	0.0005
Household income					
\$30k-\$59,999k	-4.15	4.62	-13.21 – 4.91	-0.90	0.37
\$60k-\$99,999k	-3.68	5.15	-13.77 – 6.42	-0.71	0.48
≥ \$100k	5.15	4.38	-3.44 – 13.74	1.18	0.24
Self-reported mental health	9.87	7.15	-4.15 – 23.89	1.38	0.17
Self-reported physical health	20.93	4.46	12.20 – 29.67	4.70	3.43 ^{e-06}

Table S24- Weighted ANCOVA table for category of antidepressant used and MVPA
 (controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	-0.07	0.14	-0.34 – 0.20	-0.52	0.60
SSRI	-0.19	0.08	-0.33 – -0.04	-2.47	0.01
Tri	-0.09	0.18	-0.45 – 0.27	-0.49	0.63
Other	-0.24	0.19	-0.61 – 0.13	-1.28	0.20
Age	-0.009	0.001	-0.01 – -0.007	-8.40	5.31 ^{e-16}
BMI	-0.03	0.004	-0.04 – -0.03	-8.90	<2 ^{e-16}
Sex (women)	-0.25	0.04	-0.33 – -0.18	-6.65	7.99 ^{e-11}
Accelerometer wear time	0.08	0.009	0.06 – 0.10	8.89	<2 ^{e-16}
Worked last year					
Study/retired	-0.06	0.16	-0.37 – 0.26	-0.35	0.72
Working	0.19	0.06	0.08 – 0.30	3.48	0.0005
Seasons					
Spring	-0.02	0.07	-0.16 – 0.13	-0.22	0.83
Summer	-0.06	0.06	-0.19 – 0.06	-0.97	0.33
Winter	-0.07	0.09	-0.26 – 0.11	-0.77	0.44
Marital status (couple)	-0.20	0.04	-0.28 – -0.12	-4.65	4.36 ^{e-06}
Levels of cotinine	-0.0001	0.00003	-0.0002 – -0.00008	-4.64	4.57 ^{e-06}
Education					
High school	0.05	0.08	-0.10 – 0.20	0.64	0.52
Work school	0.03	0.07	-0.11 – 0.17	0.44	0.66
College	0.12	0.08	-0.04 – 0.28	1.52	0.13
University <Bachelor	0.13	0.07	-0.01 – 0.28	1.81	0.07
Bachelor	0.31	0.09	0.13 – 0.50	3.30	0.001
University> Bachelor	0.43	0.15	0.14 – 0.71	2.95	0.003
Other	-0.01	0.06	-0.13 – 0.11	-0.22	0.83
Household income					
\$30k-\$59,999k	-0.03	0.06	-0.15 – 0.09	-0.44	0.66
\$60k-\$99,999k	-0.01	0.07	-0.15 – 0.13	-0.17	0.87
≥ \$100k	0.05	0.07	-0.09 – 0.19	0.72	0.47
Self-reported mental health	0.008	0.12	-0.22 – 0.23	0.07	0.94
Self-reported physical health	-0.35	0.07	-0.49 – -0.21	-4.87	1.51 ^{e-06}

Table S25- Weighted ANCOVA table for category of antidepressant used and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	8.28	9.33	-10.01 – 26.57	0.89	0.38
SSRI	8.34	5.12	-1.69 – 18.37	1.63	0.10
Tri	-4.64	8.40	-21.10 – 11.82	-0.55	0.58
Other	-8.84	7.44	-23.42 – 5.74	-1.19	0.24
Age	-0.66	0.10	-0.85 - -0.46	-6.59	1.18 ^{e-10}
BMI	0.005	0.24	-0.47 – 0.48	0.02	0.98
Sex (women)	-6.25	2.57	-11.29 - -1.20	-2.43	0.02
Accelerometer wear time	21.86	0.91	20.08 – 23.64	24.07	<2 ^{e-16}
Worked last year					
Study/retired	-10.77	4.28	-19.16 - -2.38	-2.52	0.01
Working	22.82	3.19	16.57 – 29.07	7.16	3.14 ^{e-12}
Seasons					
Spring	3.63	4.60	-5.39 – 12.65	0.79	0.43
Summer	13.62	5.27	3.28 – 23.95	2.58	0.01
Winter	-9.00	4.66	-18.13 – 0.13	-1.93	0.05
Marital status (couple)	16.11	2.48	11.26 – 20.97	6.51	1.94 ^{e-10}
Levels of cotinine	0.001	0.003	-0.004 – 0.007	0.43	0.67
Education					
High school	21.26	7.33	6.90 – 35.62	2.90	0.004
Work school	5.55	7.33	-8.82 – 19.92	0.76	0.45
College	3.46	6.89	-10.04 – 16.96	0.50	0.62
University <Bachelor	-12.45	6.11	-24.21 - -0.69	-2.08	0.04
Bachelor	-16.63	6.39	-29.14 - -4.11	-2.60	0.01
University> Bachelor	-17.59	7.36	-32.02 - -3.16	-2.39	0.02
Other	19.80	4.84	10.32 – 29.29	4.09	0.00005
Household income					
\$30k-\$59,999k	5.38	4.54	-3.51 – 14.27	1.19	0.24
\$60k-\$99,999k	4.02	5.20	-6.18 – 14.21	0.77	0.44
≥ \$100k	-5.54	4.56	-14.49 – 3.40	-1.21	0.23
Self-reported mental health	-11.09	6.65	-24.12 – 1.94	-1.67	0.10
Self-reported physical health	-15.87	4.27	-24.23 - -7.51	-3.72	0.0002

Table S26- Weighted ANCOVA table for category of antidepressant used and steps per day
 (controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	-220.66	471.14	-1144.07 – 702.75	-0.47	0.64
SSRI	-544.34	298.00	-1128.41 – 39.74	-1.83	0.07
Tri	298.65	582.90	-843.81 – 1441.11	0.51	0.61
Other	-869.35	508.52	-1866.03 – 127.32	-1.71	0.09
Age	-17.36	2.84	-22.92 - -11.80	-6.12	1.95 ^{e-09}
BMI	-74.08	9.72	-93.13 - -55.03	-7.62	1.39 ^{e-13}
Sex (women)	-856.81	104.00	-1060.65 - -652.97	-8.24	1.73 ^{e-15}
Accelerometer wear time	711.41	36.26	640.35 – 782.46	19.62	<2 ^{e-16}
Worked last year					
Study/retired	7.59	277.27	-535.84 – 551.02	0.03	0.98
Working	961.28	172.84	622.52 – 1300.05	5.56	4.48 ^{e-08}
Seasons					
Spring	64.08	255.15	-436.02 – 564.17	0.25	0.80
Summer	231.62	214.52	-188.83 – 652.08	1.08	0.28
Winter	-589.08	201.69	-984.39 - -193.77	-2.92	0.004
Marital status (couple)	-187.00	152.43	-485.75 – 111.76	-1.23	0.22
Levels of cotinine	-0.37	0.11	-0.58 - -0.17	-3.53	0.0004
Education					
High school	651.06	285.71	91.09 – 1211.04	2.28	0.02
Work school	133.60	270.59	-396.74 – 663.94	0.49	0.62
College	-121.43	277.82	-665.95 – 423.08	-0.44	0.66
University <Bachelor	125.04	275.13	-414.21 – 664.29	0.45	0.65
Bachelor	373.35	332.75	-278.82 – 1025.52	1.12	0.26
University> Bachelor	486.24	415.14	-327.41 – 1299.89	1.17	0.24
Other	493.00	232.95	36.44 – 949.57	2.12	0.03
Household income					
\$30k-\$59,999k	240.47	195.21	-142.13 – 623.06	1.23	0.22
\$60k-\$99,999k	150.10	173.46	-189.88 – 490.09	0.87	0.39
≥ \$100k	210.29	193.21	-168.41 – 588.98	1.09	0.28
Self-reported mental health	-410.68	393.94	-1182.79 – 361.42	-1.043	0.30
Self-reported physical health	-961.31	183.40	-1320.77 - -601.84	-5.24	2.41 ^{e-07}

Table S27- Weighted ANCOVA table for category of antidepressant used and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
ADCat					
SNRI	-7.33	10.10	-27.12 – 12.45	-0.73	0.47
SSRI	-6.69	5.21	-16.91 – 3.52	-1.28	0.20
Tri	6.87	9.63	-12.02 – 25.75	0.71	0.48
Other	11.65	8.38	-4.76 – 28.07	1.39	0.16
Age	0.84	0.10	0.63 – 1.04	8.13	3.93 ^{e-15}
BMI	0.70	0.23	0.24 – 1.15	2.98	0.003
Sex (women)	10.76	2.67	5.54 – 15.99	4.04	0.0001
Accelerometer wear time	36.30	0.92	34.49 – 38.10	39.41	<2 ^{e-16}
Worked last year					
Study/retired	10.87	4.14	2.75 – 18.99	2.62	0.009
Working	-25.79	3.79	-33.23 – -18.35	-6.80	3.25 ^{e-11}
Seasons					
Spring	-2.62	4.44	-11.33 – 6.08	-0.59	0.55
Summer	-12.30	5.49	-23.05 – 1.54	-2.24	0.03
Winter	11.22	4.39	2.62 – 19.82	2.56	0.01
Marital status (couple)	-11.28	2.77	-16.70 – -5.85	-4.08	5.39 ^{e-05}
Levels of cotinine	0.002	0.003	-0.004 – 0.007	0.60	0.55
Education					
High school	-23.28	7.52	-38.02 – -8.54	-3.10	0.002
Work school	-4.77	7.55	-19.57 – 10.02	-0.63	0.53
College	-3.52	7.68	-18.57 – 11.53	-0.46	0.65
University <Bachelor	10.95	6.35	-1.51 – 23.40	1.72	0.09
Bachelor	8.55	6.58	-4.34 – 21.45	1.30	0.19
University > Bachelor	8.68	8.09	-7.18 – 24.54	1.07	0.28
Other	-18.09	5.23	-28.33 – -7.84	-3.46	0.0005
Household income					
\$30k-\$59,999k	-4.78	4.66	-13.91 – 4.34	-1.03	0.30
\$60k-\$99,999k	-4.38	5.33	-14.83 – 6.06	-0.82	0.41
≥ \$100k	4.17	4.54	-4.73 – 13.07	0.92	0.36
Self-reported mental health	10.83	7.63	-4.12 – 25.78	1.42	0.16
Self-reported physical health	20.58	4.63	11.51 – 29.66	4.45	1.10 ^{e-05}

Table S28- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and MVPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD—	-0.13	0.07	-0.27 – 0.01	-1.82	0.07
Age	-0.009	0.001	-0.01 - -0.007	-8.14	3.61 ^{e-15}
BMI	-0.03	0.004	-0.04 – -0.02	-8.36	6.83 ^{e-16}
Sex (women)	-0.25	0.04	-0.32 - -0.18	-6.58	1.27 ^{e-10}
Accelerometer wear time	0.08	0.009	0.06 – 0.10	8.80	<2 ^{e-16}
Worked last year					
Study/retired	-0.04	0.16	-0.35 – 0.27	-0.25	0.81
Working	0.20	0.06	0.09 – 0.31	3.55	0.0004
Seasons					
Spring	-0.02	0.07	-0.16 – 0.12	-0.26	0.79
Summer	-0.06	0.06	-0.19 – 0.06	-1.00	0.32
Winter	-0.08	0.09	-0.26 – 0.11	-0.80	0.42
Marital status (couple)	-0.21	0.04	-0.29 - -0.12	-4.69	3.52 ^{e-06}
Levels of cotinine	-0.0002	0.00003	-0.0002 - -0.00008	-4.64	4.48 ^{e-06}
Education					
High school	0.05	0.08	-0.09 – 0.20	0.70	0.49
Work school	0.04	0.07	-0.11 – 0.18	0.49	0.63
College	0.13	0.08	-0.03 – 0.28	1.61	0.11
University <Bachelor	0.14	0.08	-0.009 – 0.29	1.84	0.07
Bachelor	0.31	0.10	0.12 – 0.50	3.26	0.001
University> Bachelor	0.43	0.15	0.14 – 0.72	2.89	0.004
Other	-0.009	0.06	-0.13 – 0.11	-0.14	0.89
Household income					
\$30k-\$59,999k	-0.02	0.06	-0.14 – 0.10	-0.34	0.74
\$60k-\$99,999k	-0.004	0.07	-0.15 – 0.14	-0.06	0.95
≥ \$100k	0.06	0.07	-0.08 – 0.20	0.80	0.42
Self-reported mental health	0.03	0.12	-0.21 – 0.27	0.27	0.79
Self-reported physical health	0.04	0.07	-0.49 - -0.21	-4.87	1.52 ^{e-06}

Table S29- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD—	5.05	4.52	-3.80 – 13.91	1.12	0.26
Age	-0.65	0.10	-0.85 - -0.46	-6.49	2.18 ^{e-10}
BMI	0.009	0.24	-0.47 – 0.48	0.04	0.97
Sex (women)	-6.36	2.59	-11.44 - -1.28	-2.45	0.01
Accelerometer wear time	21.96	0.93	20.14 – 23.78	23.62	<2 ^{e-16}
Worked last year					
Study/retired	-11.30	4.35	-19.83 - -2.77	-2.60	0.01
Working	23.12	3.19	16.86 – 29.38	7.24	1.83 ^{e-12}
Seasons					
Spring	3.94	4.59	-5.06 – 12.93	0.86	0.39
Summer	14.02	5.27	3.70 – 24.34	2.66	0.008
Winter	-8.77	4.66	-17.91 – 0.36	-1.88	0.06
Marital status (couple)	16.29	2.60	11.19 – 21.38	6.26	8.58 ^{e-10}
Levels of cotinine	0.001	0.003	-0.004 – 0.007	0.47	0.64
Education					
High school	21.82	7.57	6.98 – 36.67	2.88	0.004
Work school	5.51	7.61	-9.40 – 20.42	0.72	0.47
College	3.60	7.14	-10.38 – 17.59	0.51	0.61
University <Bachelor	-12.18	6.25	-24.42 – 0.07	-1.95	>0.05
Bachelor	-16.39	6.28	-28.70 - -4.07	-2.61	0.009
University> Bachelor	-17.41	7.55	-32.21 - -2.61	-2.31	0.02
Other	19.92	5.03	10.06 – 29.79	3.96	0.00009
Household income					
\$30k-\$59,999k	5.00	4.52	-3.86 – 13.86	1.11	0.27
\$60k-\$99,999k	3.62	5.24	-6.66 – 13.90	0.69	0.49
≥ \$100k	-6.76	4.62	-15.81 – 2.30	-1.46	0.14
Self-reported mental health	-8.51	6.73	-21.71 – 4.69	-1.26	0.21
Self-reported physical health	-17.29	4.17	-25.46 - -9.11	-4.15	0.00004

Table S30- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and steps per day

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD—	-500.64	266.74	-1023.44 – 22.17	-1.88	0.06
Age	-16.52	2.89	-22.19 - -10.86	-5.71	1.95 ^{e-08}
BMI	-73.15	10.18	-93.10 - -53.20	-7.19	2.58 ^{e-12}
Sex (women)	-856.99	106.84	-1066.39 - -647.59	-8.02	8.23 ^{e-15}
Accelerometer wear time	716.14	37.32	642.99 – 789.26	19.19	<2 ^{e-16}
Worked last year					
Study/retired	13.62	291.13	-556.98 – 584.22	0.05	0.96
Working	991.36	171.58	655.07 – 1327.64	5.78	1.37 ^{e-08}
Seasons					
Spring	48.56	255.75	-452.71 – 549.83	0.19	0.85
Summer	235.45	224.17	-203.91 – 674.81	1.05	0.29
Winter	-599.16	206.76	-1004.40 - -193.92	-2.90	0.004
Marital status (couple)	-208.43	155.75	-513.69 – 96.82	-1.34	0.18
Levels of cotinine	-0.37	0.10	-0.57 - -0.16	-3.49	0.0005
Education					
High school	679.78	291.31	108.83 – 1250.72	2.33	0.02
Work school	142.69	274.41	-395.14 – 680.51	0.52	0.60
College	-109.22	274.54	-647.31 – 428.87	-0.40	0.69
University <Bachelor	145.54	283.45	-410.01 – 701.10	0.51	0.61
Bachelor	388.36	330.13	-258.69 – 1035.41	1.18	0.24
University> Bachelor	498.45	412.00	-309.06 – 1305.96	1.21	0.23
Other	503.02	236.01	40.44 – 965.59	2.13	0.03
Household income					
\$30k-\$59,999k	266.65	197.47	-120.38 – 653.69	1.35	0.18
\$60k-\$99,999k	178.31	176.47	168.52 – 525.13	1.01	0.31
≥ \$100k	220.99	194.96	-161.13 – 603.11	1.13	0.26
Self-reported mental health	-329.45	421.53	-1155.63 – 496.73	-0.78	0.43
Self-reported physical health	-1023.22	186.87	-1389.48 - -656.95	-5.48	7.08 ^{e-08}

Table S31- Weighted ANCOVA table for use of no other psychotropic medication than antidepressants and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD—	-6.54	4.71	-15.76 – 2.69	-1.39	0.17
Age	0.81	0.10	0.61 – 1.01	7.86	2.58 ^{e-14}
BMI	0.70	0.24	0.24 – 1.16	2.97	0.003
Sex (women)	12.20	2.60	7.10 – 17.30	4.69	3.53 ^{e-06}
Accelerometer wear time	35.69	0.79	34.14 – 37.24	45.17	<2 ^{e-16}
Worked last year					
Study/retired	11.83	4.20	3.60 – 20.06	2.82	0.005
Working	-26.60	3.78	-34.02 - -19.18	-7.03	7.26 ^{e-12}
Seasons					
Spring	-1.65	4.35	-10.17 – 6.87	-0.38	0.70
Summer	-11.36	5.63	-22.39 - -0.33	-2.02	0.04
Winter	13.23	4.66	4.10 – 22.36	2.84	0.005
Marital status (couple)	-11.28	2.95	-17.07 - -5.50	-3.82	0.0002
Levels of cotinine	0.0007	0.003	-0.005 – 0.007	0.23	0.82
Education					
High school	-23.22	7.55	-38.01 - -8.42	-3.08	0.002
Work school	-4.40	7.73	-19.55 – 10.74	-0.57	0.57
College	-2.57	7.67	-17.61 – 12.47	-0.34	0.74
University <Bachelor	10.71	6.66	-2.33 – 23.76	1.61	0.11
Bachelor	8.52	6.48	-4.18 – 21.22	1.32	0.19
University> Bachelor	8.58	7.93	-6.97 – 24.12	1.08	0.28
Other	-18.27	5.32	-28.69 - -7.85	-3.44	0.0006
Household income					
\$30k-\$59,999k	-5.27	4.63	-14.35 – 3.80	-1.14	0.26
\$60k-\$99,999k	-4.13	5.26	-14.44 – 6.18	-0.79	0.43
≥ \$100k	5.47	4.50	-3.35 – 14.30	1.22	0.22
Self-reported mental health	7.57	7.85	-7.81 – 22.95	0.96	0.34
Self-reported physical health	22.67	4.46	13.92 – 31.42	5.08	5.52 ^{e-07}

Table S32- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and MVPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD+	-0.09	0.10	-0.29 – 0.11	-0.91	0.36
Age	-0.009	0.001	-0.01 - -0.007	-8.68	<2 ^{e-16}
BMI	-0.03	0.004	-0.04 - -0.02	-8.36	7.11 ^{e-16}
Sex (women)	-0.24	0.04	-0.32 - -0.17	-6.28	7.90 ^{e-09}
Accelerometer wear time	0.08	0.009	0.06 – 0.10	8.85	<2 ^{e-16}
Worked last year					
Study/retired	-0.10	0.17	-0.43 – 0.24	-0.56	0.58
Working	0.18	0.06	0.06 – 0.29	3.01	0.003
Seasons					
Spring	-0.01	0.08	-0.16 – 0.14	-0.13	0.90
Summer	-0.05	0.06	-0.18 – 0.07	-0.82	0.41
Winter	-0.06	0.10	-0.25 – 0.13	-0.64	0.52
Marital status (couple)	-0.20	0.04	-0.29 - -0.12	-4.62	4.87 ^{e-06}
Levels of cotinine	-0.0001	0.00003	-0.0002 - -0.00008	-4.43	1.20 ^{e-06}
Education					
High school	0.06	0.08	-0.09 – 0.21	0.73	0.46
Work school	0.02	0.07	-0.12 – 0.17	0.33	0.74
College	0.11	0.08	-0.04 – 0.27	1.40	0.16
University <Bachelor	0.12	0.08	-0.03 – 0.27	1.62	0.11
Bachelor	0.30	0.10	0.11 – 0.50	3.12	0.002
University> Bachelor	0.43	0.14	0.14 – 0.71	2.96	0.003
Other	-0.03	0.06	-0.15 – 0.09	-0.51	0.61
Household income					
\$30k-\$59,999k	-0.02	0.06	-0.15 – 0.10	-0.37	0.71
\$60k-\$99,999k	-0.03	0.07	-0.17 – 0.11	-0.39	0.70
≥ \$100k	0.05	0.07	-0.08 – 0.19	0.79	0.43
Self-reported mental health	0.03	0.13	-0.22 – 0.28	0.26	0.79
Self-reported physical health	-0.34	0.07	-0.48 - -0.20	-4.78	2.40 ^{e-06}

Table S33- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and LPA

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD+	6.55	4.35	-1.97 – 15.07	1.51	0.13
Age	-0.67	0.10	-0.86 - -0.48	-6.77	3.78 ^{e-11}
BMI	0.16	0.29	-0.41 - 0.74	0.55	0.58
Sex (women)	-6.07	2.44	-10.86 - -1.28	-2.48	0.01
Accelerometer wear time	22.24	0.74	20.80 – 23.68	30.23	<2 ^{e-16}
Worked last year					
Study/retired	-12.11	4.65	-21.22 - -2.99	-2.60	0.01
Working	23.27	3.17	17.04 – 29.50	7.32	1.04 ^{e-12}
Seasons					
Spring	3.38	5.04	-6.50 – 13.26	0.67	0.50
Summer	13.72	5.66	2.62 – 24.81	2.42	0.02
Winter	-10.61	5.06	-20.52 - -0.70	-2.10	0.04
Marital status (couple)	17.74	2.67	12.50 – 22.98	6.64	8.61 ^{e-11}
Levels of cotinine	0.002	0.003	-0.004 – 0.007	0.54	0.59
Education					
High school	19.57	6.26	7.30 – 31.84	3.13	0.002
Work school	6.12	7.19	-7.97 – 20.21	0.85	0.40
College	3.34	6.08	-8.58 – 15.26	0.55	0.58
University <Bachelor	-10.14	6.07	-22.03 – 1.75	-1.67	0.10
Bachelor	-16.05	6.46	-28.73 - -3.38	-2.48	0.01
University> Bachelor	-15.77	7.44	-30.36 - -1.18	-2.12	0.03
Other	20.33	4.76	10.99 – 29.66	4.27	2.39 ^{e-05}
Household income					
\$30k-\$59,999k	6.17	4.67	-2.99 – 15.33	1.32	0.19
\$60k-\$99,999k	3.46	4.92	-6.17 – 13.10	0.70	0.48
≥ \$100k	-7.28	4.35	-15.81 – 1.25	-1.67	0.10
Self-reported mental health	-7.87	7.05	-21.70 – 5.95	-1.12	0.26
Self-reported physical health	-15.85	4.80	-25.26 - -6.43	-3.30	0.001

Table S34- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and steps per day

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD+	-48.16	236.69	-512.07 – 415.75	-0.20	0.84
Age	-18.51	3.09	-24.57 - -12.45	-5.99	4.19 ^{e-09}
BMI	-66.35	10.69	-87.31 - -45.40	-6.21	1.18 ^{e-09}
Sex (women)	-841.83	105.39	-1048.39 - -635.28	-7.99	1.05 ^{e-14}
Accelerometer wear time	724.48	37.64	650.71 – 798.25	19.25	<2 ^{e-16}
Worked last year					
Study/retired	-32.51	272.47	-566.55 – 501.53	-0.12	0.91
Working	976.01	184.59	614.22 – 1337.79	5.29	1.90 ^{e-07}
Seasons					
Spring	-25.86	256.71	-529.01 – 477.29	-0.10	0.92
Summer	305.24	225.06	-135.87 – 749.35	1.36	0.18
Winter	-534.02	211.07	-947.72 - -120.33	-2.53	0.01
Marital status (couple)	-125.56	157.27	-433.80 – 182.68	-0.80	0.43
Levels of cotinine	-0.35	0.11	-0.57 - -0.12	-3.05	0.002
Education					
High school	675.17	278.46	129.40 – 1220.94	2.43	0.02
Work school	57.39	276.12	-483.80 – 598.58	0.21	0.84
College	-153.28	288.42	-718.57 – 412.01	-0.53	0.60
University <Bachelor	107.20	276.52	-434.78 – 649.17	0.39	0.70
Bachelor	340.12	322.70	-292.36 – 972.60	1.05	0.29
University> Bachelor	584.71	392.24	-184.07 – 1353.50	1.49	0.14
Other	383.16	234.32	-76.10 – 842.42	1.64	0.10
Household income					
\$30k-\$59,999k	248.98	194.96	-133.13 – 631.09	1.28	0.20
\$60k-\$99,999k	40.23	172.53	-297.92 – 378.38	0.23	0.82
≥ \$100k	172.80	191.82	-203.16 – 548.75	0.90	0.37
Self-reported mental health	-224.37	418.40	-1044.41 – 595.68	-0.54	0.59
Self-reported physical health	-898.39	203.78	-1297.80 - -498.98	-4.41	1.29 ^{e-05}

Table S35- Weighted ANCOVA table for use of antidepressants and at least one other psychotropic medication and sedentary behaviors

(controlling for age, body mass index, sex, time wearing the accelerometer, working status, season, marital status, smoking status [blood cotinine levels], education, total household income, and self-reported mental and physical health)

	Estimate	SE	95% CI	t	p
AD+	-5.79	5.00	-15.59 – 4.01	-1.16	0.25
Age	0.83	0.10	0.63 – 1.03	8.006	9.21 ^{e-15}
BMI	0.54	0.28	-0.004 – 1.09	1.95	0.05
Sex (women)	11.10	2.52	6.17 – 16.03	4.41	1.27 ^{e-05}
Accelerometer wear time	35.93	0.79	34.39 – 37.47	45.77	<2 ^{e-16}
Worked last year					
Study/retired	13.26	4.52	-11.73 – 7.77	2.93	0.004
Working	-26.48	3.81	-23.76 - -0.35	-6.95	1.22 ^{e-11}
Seasons					
Spring	-1.98	4.97	3.00 – 21.63	-0.40	0.69
Summer	-12.06	5.97	-18.03 - -6.40	-2.02	0.04
Winter	12.31	4.75	-0.005 – 0.007	2.59	0.01
Marital status (couple)	-12.21	2.97	-18.03 - -6.40	-4.12	4.50 ^{e-05}
Levels of cotinine	0.0008	0.003	-0.005 – 0.007	0.27	0.79
Education					
High school	-23.85	7.41	-38.37 - -9.32	-3.22	0.001
Work school	-5.45	7.45	-20.05 – 9.14	-0.73	0.46
College	-2.99	6.70	-16.11 – 10.14	-0.45	0.66
University <Bachelor	8.70	6.40	-3.84 – 21.24	1.36	0.17
Bachelor	7.79	6.52	-5.00 – 20.57	1.19	0.23
University> Bachelor	6.28	7.95	-9.30 – 21.86	0.79	0.43
Other	-19.01	5.13	-29.07 - -8.96	-3.71	0.0002
Household income					
\$30k-\$59,999k	-6.14	4.82	-15.59 – 3.30	-1.27	0.20
\$60k-\$99,999k	-3.66	5.04	-13.54 – 6.22	-0.73	0.47
≥ \$100k	5.03	4.23	-3.27 – 13.32	1.19	0.24
Self-reported mental health	8.13	8.23	-7.99 – 24.25	0.99	0.32
Self-reported physical health	20.81	5.16	10.71 – 30.92	4.04	6.29 ^{e-05}