

ASYMPTOTICAL BEHAVIOUR OF ROOTS OF INFINITE COXETER GROUPS

CHRISTOPHE HOHLWEG, JEAN-PHILIPPE LABBÉ, AND VIVIEN RIPOLL

ABSTRACT. Let W be an infinite Coxeter group. We initiate the study of the set E of limit points of “normalized” roots (representing the directions of the roots) of W . We show that E is contained in the isotropic cone Q of the bilinear form B associated to a geometric representation, and illustrate this property with numerous examples and pictures in rank 3 and 4. We also define a natural geometric action of W on E , and then we exhibit a countable subset of E , formed by limit points for the dihedral reflection subgroups of W . We explain how this subset is built from the intersection with Q of the lines passing through two positive roots, and finally we establish that it is dense in E .

INTRODUCTION

When dealing with Coxeter groups, one of the most powerful tools we have at our disposal is the notion of root systems. In the case of a finite Coxeter group W , i.e., a finite reflection group, roots are representatives of normal vectors for the Euclidean reflections in W . Thinking about finite Coxeter groups and their associated *finite* root systems allows the use of arguments from Euclidean geometry and finite group theory, which makes finite root systems well studied, see for instance [Hum90, Ch.1], and the references therein.

To deal with root systems of infinite Coxeter groups, we usually distinguish two classes: affine reflection groups, and the other infinite but not affine Coxeter groups. The root systems associated to affine Coxeter groups are also well-studied: an affine root system can be realized in an affine Euclidean space as a finite root system up to translations, see for instance [Hum90, Ch.4]. For the other *infinite* (not affine) Coxeter groups, in comparison, very little is known.

While investigating a conjecture on biclosed sets of positive roots in an infinite root system (Conjecture 2.5 in [Dye11]), we came across a difficulty: we do not know much how the roots of an infinite root system are geometrically distributed over the space.

First, observe that even the term (*infinite*) *root system* seems to designate different objects, depending on whether associated to Lie algebras (see [Kac90, LN04]), Kac-Moody Lie algebras (see [MP89]) or Coxeter groups via their geometric representations (see [Hum90, Ch.5 & 6]). While all definitions of root systems are


2010 *Mathematics Subject Classification.* Primary 17B22, Secondary 20F55.

Key words and phrases. Coxeter group, root system, roots, limit point, accumulation set.

The first author is supported by a NSERC grant, the second author is supported by a FQRNT doctoral scholarship and the third author is supported by a postdoctoral fellowship from CRM-ISM and LaCIM.

related to a given bilinear form, the bilinear forms considered in the case of Kac-Moody algebras or Lie algebras are different from the one in the classical definition of a root system for infinite Coxeter groups. In particular, a difference lies in the ability to change the value of the bilinear form on a pair of reflections whose product has infinite order. In this vein, more general geometric representations of a Coxeter group and of root systems (that we take up in §1) have been introduced. These more general geometric representations were recently presented in [Kra09] and [BD10] (see also Howlett [How96]) but seem to go back to E. B. Vinberg [Vin71], as stated in D. Krammer’s thesis. They have been the framework of several recent studies about infinite root systems of Coxeter groups (see for instance [BD10, Dye10, Dye11, Fu12, Fu13]).

Taking up this framework and using the computer algebra system Sage, we obtain the following pictures (Figures 1(a) and 1(b)), which suggests that roots have a very interesting asymptotical behaviour. In this article, we initiate the study of this behaviour.



(a) The first 100 normalized roots, around the isotropic cone Q , for the rank 3 Coxeter group with the depicted graph.

(b) The first 1665 normalized roots, around the isotropic cone Q , for the rank 4 Coxeter group with the depicted graph.

FIGURE 1. Root systems for two infinite Coxeter groups computed via the computer algebra system Sage.

Let us explain what we see in these pictures. First, we fix a geometric action of W on a finite dimensional real vector space V , which implies the data of a symmetric bilinear form B , and a simple system Δ , which is a basis for V (the framework we use is introduced in detail in §1). In §2, we first show that the norm of an (injective) sequence of roots diverges to infinity. So, in order to visualize “limits” of roots (actually the limits of their directions), we cut the picture by an affine hyperplane. Define V_1 to be the affine hyperplane spanned by the points corresponding to the simple roots: Figures 1(a) and (b) live in V_1 and the triangle (resp. tetrahedron) is the convex hull of the simple roots. The blue dots are the

intersection of V_1 with the rays spanned by the roots, and we call them *normalized roots*. The red part depicts the isotropic cone $Q = \{v \in V \mid B(v, v) = 0\}$ of the quadratic form associated to B . We see on the pictures that the normalized roots tend to converge to points on Q , and that the set of limit points has an interesting structure: it seems to be equal to Q in Figure 1(a), whereas in Figure 1(b) it is similar to an Apollonian gasket.

Let E be the set of accumulation points of the normalized roots, and call *limit roots* the points of E . After having explained our framework in §1, we state our first result (Theorem 2.7 in §2.2): the set E is indeed always contained in the isotropic cone Q . M. Dyer discovered independently this property in his research on the imaginary cone of Coxeter groups, see [Dye12] and Remark 2.8. However, we state this result and its proof in an affine context, which is slightly different from M. Dyer's framework, and allows us to describe many examples and pictures in rank 2, 3 and 4. Through them, we see new geometric properties emerging; in §3 and §4 we describe two of these properties of E which we feel should motivate further works on the subject:

- (i) The geometric action of W on V induces an action on E . This action is simply given by the following process: for $\alpha \in \Delta$ and $x \in E$, the image $s_\alpha \cdot x$ of x is the intersection point (other than x , if possible) of Q with the line passing through the (normalized) root α and the point x (see Prop. 3.1 and 3.5). Contrary to the usual action of W on the roots, the action of W on E stays in the positive cone on Δ ; indeed E lies in the convex hull $\text{conv}(\Delta)$ of the simple roots. Using this action, we give in §3.2 some ways to understand the fractal phenomenon.
- (ii) The set E is the closure of the set of accumulation points obtained from the dihedral reflection subgroups of W only. Equivalently, E is the closure of the set of all points you obtain by intersecting Q with the lines in V_1 passing through two normalized roots (see Theorem 4.2).

A classical question when dealing with a property of Coxeter groups is how it passes down to subgroups (parabolic, or more generally reflection subgroups). In the last section (§5), we define the set of limit roots for a reflection subgroup, and discuss how it compares to the set of limit roots for the whole group.

Along the text, we also present possible future directions and open problems.

In a forthcoming paper ([DHR13]), the first and third authors, together with M. Dyer, show that E is the closure of the orbit of a finite set of accumulation points, and make some connections with the notions of root posets and of dominance order via the imaginary cone for Coxeter groups (cf. [Dye12, Fu13]).

Figures. The pictures were realized using the $\text{T}_\text{E}\text{X}$ -package TikZ, and computed by dint of the computer algebra system Sage [S⁺13].

1. GEOMETRIC REPRESENTATIONS OF A COXETER GROUP

In this section, we recall some properties of Coxeter groups, the construction of their associated root systems, and we fix notations. The theory of Coxeter groups is a rich one, and we recall here only what is necessary for the purpose of this paper. For more details, see for instance [Bou68, Hum90, Kan01, BB05], and the references therein.

We consider a Coxeter system (W, S) . Recall that $S \subseteq W$ is a set of generators for W , subject only to relations of the form $(st)^{m_{s,t}} = 1$, where $m_{s,t} \in \mathbb{N}^* \cup \{\infty\}$ is attached to each pair of generators $s, t \in S$, with $m_{s,s} = 1$ and $m_{s,t} \geq 2$ for $s \neq t$. We write $m_{s,t} = \infty$ if the product st has infinite order. In the following, we always suppose that S is finite, and denote by $n = |S|$ the rank of W .

1.1. The classical geometric representation of a Coxeter group.

Coxeter groups are modeled to be the abstract combinatorial counterpart of reflection groups, i.e., groups generated by reflections. It is well known that any finite Coxeter group can be represented geometrically as a (finite) reflection group. This property still holds for infinite Coxeter groups, for some adapted definition of reflection that we first recall below.

For B a symmetric bilinear form on a real vector space V (of finite dimension), and $\alpha \in V$ such that $B(\alpha, \alpha) \neq 0$, we denote by s_α the following map:

$$(1.1) \quad s_\alpha(v) = v - 2 \frac{B(\alpha, v)}{B(\alpha, \alpha)} \alpha, \quad \text{for any } v \in V.$$

We denote by $H_\alpha := \{v \in V \mid B(\alpha, v) = 0\}$ the orthogonal of the line $\mathbb{R}\alpha$ for the form B . Since $B(\alpha, \alpha) \neq 0$, we have $H_\alpha \oplus \mathbb{R}\alpha = V$. It is straightforward to check that s_α fixes H_α , that $s_\alpha(\alpha) = -\alpha$, and s_α also preserves the form B , so it lies in the associated orthogonal group $O_B(V)$. We call s_α the *B-reflection associated to α* (or simply reflection whenever B is clear). When B is a scalar product, this is of course the usual formula for a Euclidean reflection.

Let us now recall this *classical geometric representation* (following [Hum90, §5.3-5.4]). Consider a real vector space V of dimension n , with basis $\Delta = \{\alpha_s \mid s \in S\}$. We define a symmetric bilinear form B by:

$$B(\alpha_s, \alpha_t) = \begin{cases} -\cos\left(\frac{\pi}{m_{s,t}}\right) & \text{if } m_{s,t} < \infty, \\ -1 & \text{if } m_{s,t} = \infty. \end{cases}$$

Then any element s of S acts on V as the B -reflection associated to α_s (as defined in Equation (1.1)), i.e., $s(v) = v - 2B(\alpha_s, v) \alpha_s$ for $v \in V$. This induces a *faithful* action of W on V , which preserves the form B ; thus we denote by the same letter an element of W and its associated element of $O_B(V)$.

1.2. Root system and reflection subgroups of a Coxeter group.

The root system of W is a way to encode the reflections of the Coxeter group, i.e., the conjugates of elements of S , which are called simple reflections. The elements of $\Delta = \{\alpha_s \mid s \in S\}$ are called *simple roots* of W , and the *root system* Φ of W is defined to be the orbit of Δ under the action of W . By construction, any root $\rho \in \Phi$ gives rise to the reflection s_ρ of W , which is conjugate to some $s_\alpha \in S$.

A *reflection subgroup* of W is a subgroup of W generated by reflections; so it can be built from a subset of Φ . It turns out that any such reflection subgroup is again a Coxeter group, with some canonical generators ([Deo89, Dye90]). So it is natural to desire to apply results valid for W to a reflection subgroup simply by restriction. A major drawback of the classical geometric representation described above is that it is not “functorial” with respect to the reflection subgroups: it can happen that the representation of some reflection subgroups W' of W , induced (by restriction) by the geometric representation of W , is not the same as the geometric

representation of W' as a Coxeter group, see Example 1.1 below. Note that it does also happen that a Coxeter group of rank n contains a reflection subgroup of higher rank, as shown in Example 5.1.

Example 1.1 (Reflection subgroups of rank 2). Let us consider the Coxeter group of rank 3 with $S = \{s_\alpha, s_\beta, s_\gamma\}$ and $m_{s_\alpha, s_\beta} = 5$, $m_{s_\beta, s_\gamma} = m_{s_\alpha, s_\gamma} = 3$ (whose Coxeter diagram is on Figure 1(a)). Take the root $\rho = s_\alpha s_\beta(\alpha) = s_\beta s_\alpha(\beta)$, so that s_ρ corresponds to the longest element in the subgroup $\langle s_\alpha, s_\beta \rangle$: $s_\rho = s_\alpha s_\beta s_\alpha s_\beta s_\alpha = s_\beta s_\alpha s_\beta s_\alpha s_\beta$. We compute $\rho = \frac{1+\sqrt{5}}{2}(\alpha + \beta)$. Consider the reflection subgroup W' generated by s_γ and s_ρ . The product $s_\gamma s_\rho$ has infinite order, so W' is an infinite dihedral group, with generators s_γ and s_ρ . But, if B denotes the bilinear form associated to the Coxeter group W , we get: $B(\gamma, \rho) = -\frac{1+\sqrt{5}}{2} \neq -1$. So, the restriction to W' of the geometric representation of W does not correspond to the classical geometric representation of W' as an infinite dihedral group. In Example 2.4 we give a geometric interpretation of this fact, which is visible in Figure 1(a).

1.3. Coxeter groups from based root systems.

In order to solve the problem in Example 1.1, we relax the requirements on the bilinear form B used to represent the group W : we allow the values of some $B(\alpha, \beta)$ to be any real numbers less than or equal to -1 (when the associated product of reflections $s_\alpha s_\beta$ has infinite order). Actually, an even more general setting is better adapted here: the notion of a *based root system* (used for instance in [How96, Kra09, BD10]).

Definition 1.2. Let V be a real vector space, equipped with a bilinear form B . Consider a finite subset Δ of V such that

- (i) Δ is positively independent¹: if $\sum_{\alpha \in \Delta} \lambda_\alpha \alpha = 0$ with all $\lambda_\alpha \geq 0$, then all $\lambda_\alpha = 0$;
- (ii) for all $\alpha, \beta \in \Delta$, with $\alpha \neq \beta$, $B(\alpha, \beta) \in (-\infty, -1] \cup \{-\cos(\frac{\pi}{k}), k \in \mathbb{Z}_{\geq 2}\}$;
- (iii) for all $\alpha \in \Delta$, $B(\alpha, \alpha) = 1$.

Such a set Δ is called a *simple system*. Denote by $S := \{s_\alpha \mid \alpha \in \Delta\}$ the set of B -reflections associated to elements in Δ (see Equation (1.1)). Let W be the subgroup of $O_B(V)$ generated by S , and Φ be the orbit of Δ under the action of W .

The pair (Φ, Δ) is a *based root system in (V, B)* ; its *rank* is the cardinality of Δ , i.e., the cardinality of S . We call the pair (V, B) a *geometric W -module*². We equivalently refer to a *geometric representation of W* instead of a geometric W -module.

Remark 1.3.

- Condition (ii) is natural to ensure that subrepresentations are again geometric representation in the sense of this new definition (we saw in Example 1.1 that this does not work for the usual definition).
- In Condition (i), the relaxation is more subtle, but also necessary if we want a nice functorial behaviour on the subrepresentations. For instance, for some Coxeter group W there exists a reflection subgroup of rank (as a Coxeter group) strictly higher than that of W , see Example 5.1.

¹Geometrically, this means we require that 0 does not lie in the convex hull of the points of Δ .

²The triplet (V, Δ, B) is sometimes called a *Coxeter datum* in the literature, see for instance [Fu12, Fu13].

- Even if Δ is not anymore required to be a basis, the condition that it is positively independent is *necessary* to keep the usual properties of root systems, in particular the distinction between a set of *positive* roots and a set of *negative* roots. Indeed, it is not difficult to prove that a set is positively independent if and only if its is included in an open half-space supported by a linear hyperplane, see [How96, p.4 note (b)].

This generalization of root system enjoys the following expected properties (see for instance [Kra09, BD10]):

- (W, S) is a Coxeter system³, where the order of $s_\alpha s_\beta$ is k whenever $B(\alpha, \beta) = -\cos(\frac{\pi}{k})$, and ∞ if $B(\alpha, \beta) \leq -1$.
- The convex set $\text{cone}(\Delta)$ consisting of all positive linear combinations of elements of Δ allows us to define the set of *positive roots* $\Phi^+ := \Phi \cap \text{cone}(\Delta)$, and then $\Phi = \Phi^+ \sqcup (-\Phi^+)$ and $\mathbb{R}\rho \cap \Phi = \{\rho, -\rho\}$, for $\rho \in \Phi$.

The classical geometric representation (that we recalled in §1.1) is an example of such a geometric W -module. If all $m_{s,t}$ (called the *labels* of the group) are finite, then the only possible representation (supposing that Δ is a basis) is the classical one. In particular, when the form B is positive definite, then Φ is a *finite* root system and contains no more information than its associated finite Coxeter group.

We say that (Φ, Δ) is an *affine based root system* when the form B is positive semidefinite, but not definite. Note that traditionally, the Coxeter group itself is said to be affine if the root system of its classical geometric representation is affine.

Example 1.4 (Irreducible affine root systems). An infinite dihedral group W has non-affine geometric representations as well as the classical affine representation. If Φ is an infinite root system of rank 2, with simple roots α, β , then $B(\alpha, \beta) \leq -1$, and Φ is affine if and only if $B(\alpha, \beta) = -1$ (i.e., when Φ corresponds to the classical geometric representation of W). We give a geometric description of these two cases in Figure 2. However, note that if W is irreducible⁴ of rank ≥ 3 , then Φ is affine if and only if W is affine (because there is no label ∞ in an irreducible affine Coxeter graph of rank ≥ 3).

All the desired properties of the root system and of positive and negative roots still hold for a based root system. In particular, the following statements, which are essential in the next sections, are still valid in this new framework.

Proposition 1.5. *Let (Φ, Δ) be a based root system in (V, B) , with associated Coxeter system (W, S) .*

- The set $\{B(\alpha, \rho) \mid \alpha \in \Delta, \rho \in \Phi^+ \text{ and } |B(\alpha, \rho)| < 1\}$ is finite.
- Denote by Q the isotropic cone:

$$Q := \{v \in V \mid q(v) = 0\}, \text{ where } q(v) = B(v, v).$$

Let $\rho_1 \neq \rho_2$ be two roots in Φ^+ . Denote by W' the dihedral reflection subgroup of W generated by the two reflections s_{ρ_1} and s_{ρ_2} , and

$$\Phi' := \{\rho \in \Phi \mid s_\rho \in W'\}.$$

Then there exists $\Delta' \subseteq \Phi^+ \cap \Phi'$ of cardinality 2 such that (Φ', Δ') is a based root system of rank 2, with associated Coxeter group W' . Moreover:

³This result is [Vin71, Theorem 1.2].

⁴See §2.3 for the definition.

- (a) Φ' is infinite if and only if the plane $\text{span}(\rho_1, \rho_2)$ intersects $Q \setminus \{0\}$, if and only if $|B(\rho_1, \rho_2)| \geq 1$;
- (b) Φ' is affine if and only if $\text{span}(\rho_1, \rho_2) \cap Q$ is a line, if and only if $B(\rho_1, \rho_2) = \pm 1$;
- (c) when Φ' is infinite, $\Delta' = \{\rho_1, \rho_2\}$ if and only if $B(\rho_1, \rho_2) \leq -1$.

Proof. The nontrivial parts of the proofs of these statements in the context of a based root system are word for word the same as the proofs in the case of the root system of the classical geometric representation, proofs that the reader may find for instance in [BB05, §4.5]. The last statement (ii)(c) is a consequence of [Dye90, Theorem 4.4], see also [Fu12, Theorem 1.8 (ii)]. \square

1.4. Other geometric representations.

Let (W, S) be a Coxeter group. Fix a matrix $A = (a_{s,t})_{s,t \in S}$ such that

$$(1.2) \quad \begin{cases} a_{s,t} = -\cos\left(\frac{\pi}{m_{s,t}}\right) & \text{if } m_{s,t} < \infty, \\ a_{s,t} \leq -1 & \text{if } m_{s,t} = \infty. \end{cases}$$

We associate to the matrix A a *canonical geometric W -module* (V_A, B_A) as follows.

- V_A is a real vector space with basis $\Delta_A = \{\alpha_s \mid s \in S\}$ and B_A is the symmetric bilinear form defined by $B_A(\alpha_s, \alpha_t) = a_{s,t}$ for $s, t \in S$.
- Any element s of S acts on V as the B -reflection associated to α_s , i.e., $s(v) = v - 2B(\alpha_s, v) \alpha_s$ for $v \in V_A$.

Since Δ_A satisfies the requirement of Definition 1.2, W acts faithfully on V_A as the subgroup of $O_{B_A}(V_A)$ spanned by the B -reflections associated to the α_s . Moreover, (Φ_A, Δ_A) is a based root system of (V_A, B_A) , where Φ_A is the W -orbit of Δ_A . Note that giving a matrix A , as we did, is equivalent to fix the values in Conditions (ii) and (iii) in Definition 1.2.

Example 1.6 (Continuation of Example 1.1). In the case of Example 1.1, the restriction of the classical geometric representation of W to the reflection subgroup W' generated by s_γ and s_ρ gives the geometric representation that is associated to the canonical geometric W' -module given by the matrix

$$A = \begin{pmatrix} 1 & -\frac{1+\sqrt{5}}{2} \\ -\frac{1+\sqrt{5}}{2} & 1 \end{pmatrix}.$$

Remark 1.7. By construction, in the based root system (Φ_A, Δ_A) associated to the canonical geometric W -module (V_A, B_A) defined above, the set Δ_A is a basis. This setting will actually be the one used throughout the three next subsections:

In §§2-3-4, we assume that the set of simple roots is a basis for the vector space.

Note that analogous results remain true in all generality, but this assumption highly simplifies the constructions and the statements, and allows us to get into the main subject more quickly. However, for the sake of completeness, we added in §5 a discussion of the case where Δ is not a basis. Moreover, this turns out to be necessary in order to deal thoroughly with the behaviour of restriction to reflection subgroups.

2. LIMIT POINTS OF NORMALIZED ROOTS AND ISOTROPIC CONE

Let (Φ, Δ) be a based root system in (V, B) , with associated Coxeter group W (as defined in Definition 1.2). We suppose that Δ is a basis for V ; analogous results remain true in full generality (see §5.2 for details).

When W is finite, Φ is also finite and the distribution of the roots in the space V is well studied. However, when W is infinite, the root system is infinite and we have, as far as we know, not many tools to study the distribution of the roots over V . The asymptotical behaviour of roots is one of them. This section deals with a first step of this study: we show that the “lengths” of the roots tend to infinity, and that the limit points of the “directions” of the roots are included in the isotropic cone of the bilinear form associated to Φ . In order to get a first grip of what happens, we begin with some enlightening examples.

2.1. Roots and normalized roots in rank 2, 3, 4, and general setting.

Since $\Phi = \Phi^+ \sqcup (-\Phi^+)$, it is enough to look at the positive roots, which are inside the simplicial cone $\text{cone}(\Delta)$.

Example 2.1. (Rank 2: affine and non-affine representations of infinite dihedral groups). Let (Φ, Δ) be a based root system of rank 2, as defined in §1.3. We get a Coxeter group W of rank 2, geometrically represented in a 2-dimensional vector space V (together with a bilinear form B), where V is generated by two simple roots α, β . Assume that W is an infinite dihedral group, so $B(\alpha, \beta) \leq -1$. Suppose first that $B(\alpha, \beta) = -1$, i.e., that Φ is affine and with the classical geometric representation. Then any positive root has the following form:

$$\rho_n = (n+1)\alpha + n\beta, \text{ or } \rho'_n = n\alpha + (n+1)\beta, \text{ for } n \in \mathbb{N}.$$

If we fix a (Euclidean) norm on V (e.g., such that $\{\alpha, \beta\}$ is an orthonormal basis), then the norms of the roots tend to infinity, but their directions tend to the line generated by $\alpha + \beta$ as depicted in Figure 2 (a). Note that this line is precisely the isotropic cone of the bilinear form B , i.e., the set

$$Q := \{v \in V \mid q(v) = 0\}, \text{ where } q(v) = B(v, v).$$

In a general geometric representation of W , Φ can be non-affine, i.e., $B(\alpha, \beta) = k$ with $k < -1$. Then the isotropic cone Q consists of two lines (generated by $(-k \pm \sqrt{k^2 - 1})\alpha + \beta$). If we draw the roots, we note that, again, their norms diverge to infinity and their directions tend to the two directions of the lines of Q ; see Figure 2 (b), and [How96, p.3] for a detailed computation.

Let us go back to the general case of an infinite based root system of rank n . In the simple example of dihedral groups, we saw that the roots themselves have no limit points; this phenomenon is actually general, so we are rather interested in the asymptotical behaviour of their directions. In order to talk properly about limits of directions, we want to “normalize” the roots and construct “unit vectors” representing each root. One simple way to do so is to intersect the line $\mathbb{R}\beta$ generated by a root β in Φ with the affine hyperplane V_1 spanned by the simple roots (seen as points), i.e., the affine hyperplane

$$V_1 := \{v \in V \mid \sum_{\alpha \in \Delta} v_\alpha = 1\},$$




FIGURE 2. The isotropic cone Q and the first positive roots of an infinite based root system of rank 2. (a): in the (classical) affine representation. (b): in a non-affine representation (the red part Q^- denotes the set $\{v \in V \mid q(v) < 0\}$).

where the v_α 's are the coordinates of v in the basis Δ of simple roots. That way, we obtain what we call the set of *normalized roots*, denoted by $\widehat{\Phi}$:

$$\widehat{\Phi} := \bigcup_{\beta \in \Phi} \mathbb{R}\beta \cap V_1.$$

Let us describe this set more precisely. Set

$$\begin{aligned} V_0 &:= \{v \in V \mid |v|_1 = 0\}, \quad \text{and} \\ V_0^+ &:= \{v \in V \mid |v|_1 > 0\}, \quad \text{where } |v|_1 := \sum_{\alpha \in \Delta} v_\alpha. \end{aligned}$$

Note that $|\cdot|_1$ is not a norm on V . Since all the positive roots are in the half-space V_0^+ , the entire root system Φ is contained in $V \setminus V_0$. So the following normalization map can be applied to Φ :

$$\begin{aligned} V \setminus V_0 &\rightarrow V_1 \\ v &\mapsto \widehat{v} := \frac{v}{|v|_1}. \end{aligned}$$

For any subset A of $V \setminus V_0$, write \widehat{A} for the set $\{\widehat{a} \mid a \in A\}$. Because for $\rho \in \Phi$, $\mathbb{R}\rho \cap \Phi = \{\rho, -\rho\}$, it is then obvious that Φ^+ is in bijection with

$$\widehat{\Phi} = \widehat{\Phi}^+ = \widehat{-\Phi^+} = \{\widehat{\rho} \mid \rho \in \Phi^+\}.$$

Remark 2.2. Obviously, we could also have considered other affine hyperplanes to “cut” the rays of Φ ; it suffices that the chosen hyperplane be “transverse to Φ^+ ”, and this is discussed in §5.2. We could also have considered the roots abstractly, in the projective space $\mathbb{P}V$. The principal advantage to consider an affine hyperplane explicitly, such as V_1 , is to visualize positive roots in an affine subspace of dimension $n-1$, inside an n -simplex (here $n = \dim V$). Indeed, the simple roots are in V_1 , so $\widehat{\Phi}$ lies in the convex hull $\text{conv}(\Delta)$, which is an n -simplex in V_1 . Note that as a

convex polytope, $\text{conv}(\Delta)$ is closed and compact, which is practical when studying sequences of roots. From now on, in examples, we only draw the normalized roots inside the n -simplex $\text{conv}(\Delta)$.

The aim of this work is to study the accumulation points of $\widehat{\Phi}$, i.e., the set of limit points of normalized roots. We first examine its relation with the isotropic cone Q .

Example 2.3 (Normalized roots in the dihedral case). In the infinite dihedral case, the “normalized” version of Figure 2 is Figure 3. Here $\widehat{\Phi}$ is contained in the segment $[\alpha, \beta]$ and there are one or two limit points of normalized roots (depending on whether $B(\alpha, \beta) = -1$ or not), and the set of limit points is always equal to the intersection $Q \cap V_1 = \widehat{Q}$.




FIGURE 3. The normalized isotropic cone \widehat{Q} and the first normalized roots of an infinite based root system of rank 2. (a): in the (classical) affine representation. (b): in a non-affine representation.

Notation. The graph we draw to define a based root system is the same as the classical Coxeter graph, except that, when the label of the edge $s_\alpha - s_\beta$ is ∞ , we specify in parenthesis the value of $B(\alpha, \beta)$ if it is not -1 (i.e., when we do not consider the classical representation).

We give now some examples and pictures in rank 3 and 4.

Example 2.4 (Rank 3). In Figures 1(a) (in the introduction) and 4 through 7, we draw the normalized isotropic cone \widehat{Q} (in red), the 3-simplex cone (Δ) (in green), and the first normalized roots (in blue), for five different based root systems of rank 3. Note that the notion of depth used in the captions is a measure of the “complexity” of a root, which is defined in §2.2.

The normalized roots seem again to tend quickly towards \widehat{Q} . In the affine cases, \widehat{Q} contains only one point, which is the intersection of the line V^\perp (the radical of B) with V_1 . In rank 3, there are three different types: \widetilde{A}_2 , \widetilde{B}_2 , and \widetilde{G}_2 . The latter is drawn in Figure 4. Otherwise, \widehat{Q} is always a conic (because the signature of B is $(2, 1)$), and moreover it is always an ellipse in the classical geometric representation (see §5.2 for more details).

Some rank 2 root subsystems appear in the pictures; they correspond to dihedral reflection subgroups. The normalized roots corresponding to such a reflection subgroup, generated by two reflections s_{ρ_1} and s_{ρ_2} , lie in the line containing the normalized roots $\widehat{\rho}_1$ and $\widehat{\rho}_2$. Because of Proposition 1.5 (ii), the subgroup is infinite if and only if \widehat{Q} intersects this line. In Figure 1(a), for the group with labels 5, 3, 3,




FIGURE 4. The normalized isotropic cone \widehat{Q} and the first normalized roots (with depth ≤ 12) for the based root system of type \widehat{G}_2 (affine).




FIGURE 5. The normalized isotropic cone \widehat{Q} and the first normalized roots (with depth ≤ 10) for the based root system with labels 2, 3, 7.




FIGURE 6. The normalized isotropic cone \widehat{Q} and the first normalized roots (with depth ≤ 10) for the based root system with labels $2, \infty(-1.1), \infty(-1.1)$.




FIGURE 7. The normalized isotropic cone \widehat{Q} and the first normalized roots (with depth ≤ 8) for the based root system with labels $\infty, \infty(-1.5), 4$.

the line joining γ and $\widehat{\rho} = \frac{\alpha+\beta}{2}$ intersects the ellipse in two points, as predicted by Example 1.1.

In general, the behaviour for standard parabolic dihedral subgroups is seen on the faces of the simplex, where three situations can occur. The ellipse \widehat{Q} can either cut an edge $[\alpha, \beta]$ in two points, or be tangent, or not intersect it, depending on whether $B(\alpha, \beta) < -1, = -1, \text{ or } > -1$ respectively; see in particular Figures 6 and 7.

Remark 2.5. When \widehat{Q} is included in the simplex, it seems that the limit points of normalized roots cover the whole ellipse, whereas in the other cases the behaviour is more complicated. We discuss this phenomenon in §3.2.

Example 2.6 (Rank 4). Figures 1(b) (in the introduction), and 8-9 illustrate some based root systems of rank 4, together with the tetrahedron $\text{conv}(\Delta)$. Analogous




FIGURE 8. The normalized isotropic cone \widehat{Q} and the first normalized roots (with depth ≤ 8) for the based root system with diagram the complete graph with labels 3.




FIGURE 9. The normalized isotropic cone \widehat{Q} and the first normalized roots (with depth ≤ 8) for the based root system with diagram the complete graph with labels ∞ .

properties seem to be true: the limit points are in \widehat{Q} , and the way how \widehat{Q} cuts a facet depends on whether the associated standard parabolic subgroup of rank 3 is infinite non affine, affine, or finite. Moreover, Remark 2.5 still holds: in Figure 8 the limit points seem to cover the whole of \widehat{Q} , whereas in Figures 1(b) and 9, some Apollonian gasket shapes appear. This fractal behaviour is discussed in §3.2.

2.2. The limit points of normalized roots lie in the isotropic cone.

Recall that we denote by q the quadratic form associated to B , and by Q the isotropic cone:

$$Q := \{v \in V \mid q(v) = 0\}, \text{ where } q(v) = B(v, v).$$

The following theorem summarizes our first observations.

Theorem 2.7. *Consider an injective sequence of positive roots $(\rho_n)_{n \in \mathbb{N}}$, and suppose that $(\widehat{\rho}_n)$ converges to a limit ℓ . Then:*

- (i) *the norm $\|\rho_n\|$ tends to infinity (for any norm on V);*
- (ii) *the limit ℓ lies in $\widehat{Q} = Q \cap V_1$.*

In other words, the set \widehat{Q} of accumulation points of normalized roots $\widehat{\Psi}$ is contained in the isotropic cone.

Remark 2.8. M. Dyer proved independently this property in the context of his work on imaginary cone [Dye12]⁵, extending a study of V. Kac (in the framework of Weyl groups of Lie algebras), who states that the convex hull of the limit points correspond to the closure of the imaginary cone (see [Kac90, Lemma 5.8 and Exercise 5.12]).

⁵M. Dyer, personal communication, September 2011.

Note that this theorem has the following consequence (which can of course be proved more directly using the fact that W is discrete in $\mathrm{GL}(V)$, see [Kra09] or [Hum90, Prop. 6.2]):

Corollary 2.9. *The set of roots of a Coxeter group is discrete.*

Proof. Suppose ρ_n is an injective sequence converging to $\rho \in \Phi^+$. Then $\widehat{\rho_n}$ converges to $\widehat{\rho}$, so by Theorem 2.7, $\widehat{\rho} \in Q$. Therefore $q(\widehat{\rho}) = 0$ which gives a contradiction since $q(\rho) = 1$. \square

The remainder of this subsection is devoted to the proof of Theorem 2.7. We first need to recall the notion of depth of a root. The depth of a positive root is a natural way to measure the ‘‘complexity’’ of this root in regard to the simple root it is obtained from (see [Sau91, §4] or [BB05, §4.6] for details): for $\rho \in \Phi^+$,

$$\mathrm{dp}(\rho) = 1 + \min\{k \mid \rho = s_{\alpha_1} s_{\alpha_2} \dots s_{\alpha_k}(\alpha_{k+1}), \text{ for } \alpha_1, \dots, \alpha_k, \alpha_{k+1} \in \Delta\}.$$

The depth is a very useful tool that allows inductive proof in infinite root systems: if γ is a root of depth $r \geq 2$, then there is a root γ' of depth $r - 1$ and a simple root $\alpha \in \Delta$ such that $\gamma = s_\alpha(\gamma')$, and moreover $B(\alpha, \gamma') < 0$, see [BB05, Lemma 4.6.2].

Since S is assumed to be finite, it follows that the number of positive roots of bounded depth is finite. Consider an injective sequence $(\rho_n)_{n \in \mathbb{N}}$ of positive roots, as in Theorem 2.7. Then we obtain easily that $\mathrm{dp}(\rho_n)$ diverges to infinity as $n \rightarrow \infty$. So, to prove the first item of the theorem, it is sufficient to show that when the depth of a sequence of roots tends to infinity, so does the norm of the roots. This is done using the following lemma, which clarifies the relation between norm and depth.

Lemma 2.10. *Let (Φ, Δ) be a based root system, as defined in §1.3. We take for the norm $\|\cdot\|$ the Euclidean norm for which Δ is an orthonormal basis for V .*

Then, with the above notations, we have the following properties:

- (i) $\exists \kappa > 0, \forall \alpha \in \Delta, \forall \rho \in \Phi^+, B(\alpha, \rho) \neq 0 \Rightarrow |B(\alpha, \rho)| \geq \kappa$;
- (ii) $\exists \lambda > 0, \forall \rho \in \Phi^+, \|\rho\|^2 \geq 1 + \lambda(\mathrm{dp}(\rho) - 1)$.

Proof. The first point is a direct consequence of Proposition 1.5 (i). Let us now prove, by induction on $\mathrm{dp}(\rho)$, that

$$\forall \rho \in \Phi^+, \|\rho\|^2 \geq 1 + \lambda(\mathrm{dp}(\rho) - 1),$$

where $\lambda = 4\kappa^2$ with κ given by (i). If $\mathrm{dp}(\rho) = 1$, $\rho \in \Delta$ so $\|\rho\| = 1 = 1 + \lambda(\mathrm{dp}(\rho) - 1)$ by the choice of the norm $\|\cdot\|$. If $\mathrm{dp}(\rho) = r \geq 2$, then we can write $\rho = s_\alpha(\rho')$, with $\rho' \in \Phi^+$ and $\alpha \in \Delta$ such that $\mathrm{dp}(\rho') = r - 1$ and $B(\alpha, \rho') < 0$, by [BB05, Lemma 4.6.2]. We get

$$\begin{aligned} \|\rho\|^2 &= \|\rho' - 2B(\alpha, \rho')\alpha\|^2 \\ &= \|\rho'\|^2 + 4B(\alpha, \rho')^2 - 4B(\alpha, \rho') \langle \alpha, \rho' \rangle, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ is the Euclidean product of $\|\cdot\|$. But we know that $B(\alpha, \rho') < 0$, and $\langle \alpha, \rho' \rangle \geq 0$ since $\rho' \in \mathrm{cone}(\Delta)$ and Δ is an orthonormal basis for $\|\cdot\|$. So we obtain by induction hypothesis on ρ' and by (i):

$$\|\rho\|^2 \geq \|\rho'\|^2 + 4B(\alpha, \rho')^2 \geq 1 + (r - 2)\lambda + 4\kappa^2.$$

Since $\lambda = 4\kappa^2$, we have $\|\rho\|^2 \geq 1 + (r - 2)\lambda + \lambda = 1 + (r - 1)\lambda$, which concludes the proof of (ii). \square

We can now finish the proof of Theorem 2.7.

Proof of Theorem 2.7. As explained before Lemma 2.10, and by (ii) of this same lemma, the norm $\|\rho_n\|$ of any injective sequence $(\rho_n)_{n \in \mathbb{N}}$ in Φ^+ tends to infinity. Recall from §2.1 that for $v \in V_0^+$, $\widehat{v} = \frac{v}{|v|_1}$, where $|v|_1 = \sum_{\alpha \in \Delta} v_\alpha$. If ρ belongs to Φ^+ , the coordinates ρ_α are nonnegative, so $|\rho|_1$ is the L_1 -norm of ρ . In particular, by equivalence of the norms, $|\rho_n|_1$ tends to infinity as $\|\rho_n\|$ does. We get

$$q(\widehat{\rho_n}) = q\left(\frac{\rho_n}{|\rho_n|_1}\right) = \frac{q(\rho_n)}{(|\rho_n|_1)^2} = \frac{1}{(|\rho_n|_1)^2} \xrightarrow{n \rightarrow \infty} 0.$$

Suppose now that $\widehat{\rho_n}$ tends to a limit ℓ . Then we obtain $q(\ell) = 0$, i.e., $\ell \in Q$, which concludes the proof of Theorem 2.7. \square

From Theorem 2.7 and its proof, we also get these easy consequences.

Corollary 2.11. *The two following statements hold:*

- (i) *for any $M \geq 0$, the set $\{\rho \in \Phi \mid \|\rho\| \leq M\}$ is finite;*
- (ii) *for any $\varepsilon > 0$, the set $\{\rho \in \Phi \mid \text{dist}(\widehat{\rho}, \widehat{Q}) \geq \varepsilon\}$ is finite.*

Definition 2.12. Let (Φ, Δ) be a based root system in (V, B) , and suppose that Δ is a basis for V . We denote by $E(\Phi)$ (or simply E when there is no possible confusion) the set of accumulation points (or limit points) of $\widehat{\Phi}$, i.e., the set consisting of all the possible limits of injective sequences of normalized roots.

Sometimes, we refer to the points of E as *limit roots* of the root system (Φ, Δ) .

Remark 2.13. As stated in Remark 2.2, we could have studied the roots in the projective space:

$$\mathbb{P}\Phi := \{\mathbb{R}\alpha \mid \alpha \in \Phi\} = \{\mathbb{R}\alpha \mid \alpha \in \Phi^+\} \subseteq \mathbb{P}V$$

with the quotient topology. However, we choose not to for two reasons: the pictures at the base of this article were obtained in an affine hyperplane and we wanted to explain precisely what we see; but also, since we do not use projective geometry technics, it seemed natural to state our result in the simplest way possible. In this context, the accumulation set $E(\Phi)$ of $\widehat{\Phi}$ is the accumulation set of $\mathbb{P}\Phi$, which is (since Φ is discrete) equal to: $\overline{\mathbb{P}\Phi} \setminus \mathbb{P}\Phi = \{\mathbb{R}x \mid x \in E(\Phi)\}$ (where $\overline{\mathbb{P}\Phi}$ denotes the topological closure of $\mathbb{P}\Phi$).

As $\widehat{\Phi}$ is included in the simplex $\text{conv}(\Delta)$ (which is closed), Theorem 2.7 implies

$$E(\Phi) \subseteq Q \cap \text{conv}(\Delta) = \widehat{Q} \cap \text{cone}(\Delta).$$

The reverse inclusion is not always true: we saw some examples of this fact in §2.1, for $\text{rk}(W) \geq 4$, or even for $\text{rk}(W) = 3$ whenever some $B(\alpha, \beta) < -1$. We address a more precise description of $E(\Phi)$ in §3.2.

Since Δ is finite, the convex set $\text{conv}(\Delta)$ is compact. Moreover, if Φ is infinite, then $\widehat{\Phi}$ is infinite as well. Therefore the inclusion $\widehat{\Phi} \subseteq \text{conv}(\Delta)$ implies the following statement.

Proposition 2.14. *Let (Φ, Δ) be a based root system in (V, B) , and suppose that Δ is a basis for V . Then $E(\Phi) \neq \emptyset$ whenever Φ is infinite.*

2.3. Simple examples : rank 2, reducible groups, and affine groups.

When (Φ, Δ) is an infinite based root system of rank 2 (with $\Delta = \{\alpha, \beta\}$), we have already explained that $E(\Phi)$ consists of one or two points (according to whether Φ is affine or not, i.e., whether $B(\alpha, \beta) = -1$ or < -1): see Example 2.1. In any case, we get $E(\Phi) = \widehat{Q}$.

For any based root system (Φ, Δ) , the limit roots coming from a given rank 2 reflection subgroup can be observed inside $E(\Phi)$. Take two distinct positive roots ρ_1 and ρ_2 , denote by W' the dihedral reflection subgroup of W generated by the two reflections s_{ρ_1} and s_{ρ_2} , and consider the associated rank 2 root subsystem (Φ', Δ') as in Proposition 1.5. Denote by $E(\Phi')$ the points of E that are limits of normalized roots of Φ' , and by $L(\widehat{\rho}_1, \widehat{\rho}_2)$ the line passing through $\widehat{\rho}_1$ and $\widehat{\rho}_2$. Then, by construction (and using Proposition 1.5), we have the following properties:

- $E(\Phi') = Q \cap L(\widehat{\rho}_1, \widehat{\rho}_2) = E(\Phi) \cap L(\widehat{\rho}_1, \widehat{\rho}_2)$;
- the cardinality of $E(\Phi')$ is 0, 1 or 2, depending on whether $|B(\rho_1, \rho_2)| < 1$, $|B(\rho_1, \rho_2)| = 1$ or $|B(\rho_1, \rho_2)| > 1$.

Another case where the equality $E(\Phi) = \widehat{Q}$ holds is when Φ is affine. To see that, we first need to discuss the reduction to irreducible root systems. Recall that a Coxeter system (W, S) is *irreducible* if there is no proper partition $S = I \sqcup J$ such that any element of I commutes with any element of J , i.e., the Coxeter graph is connected. Similarly, a based root system (Φ, Δ) is called *irreducible* if its associated Coxeter group is irreducible, i.e., if there is no proper partition $\Delta = \Delta_I \sqcup \Delta_J$ such that $B(\alpha, \beta) = 0$ for all $\alpha \in \Delta_I$ and $\beta \in \Delta_J$.

Now, for any subset Δ_I of Δ , we can associate a (*standard*) *parabolic root subsystem* (Φ_I, Δ_I) by setting:

$$W_I := \langle s_\alpha \mid \alpha \in \Delta_I \rangle \quad \text{and} \quad \Phi_I := W_I(\Delta_I).$$

It is then natural to define the limit roots for this root subsystem as the subset $E(\Phi_I)$ of $E(\Phi)$ whose elements are limits of sequences in $\widehat{\Phi}_I$. We postpone until §5 the discussion on why this definition and its analogue for more general root subsystems (not necessarily standard parabolic) make sense. Now we explain in the following easy proposition why it is possible to limit our study to irreducible root systems.

Proposition 2.15. *Assume that (Φ, Δ) is reducible, and consider proper subsets $\Delta_I, \Delta_J \subset \Delta$ such that $\Delta = \Delta_I \sqcup \Delta_J$, with $B(\alpha, \beta) = 0$ for all $\alpha \in \Delta_I$ and $\beta \in \Delta_J$. Then $E(\Phi) = E(\Phi_I) \sqcup E(\Phi_J)$.*

Proof. We have $\widehat{\Phi} = \widehat{\Phi}_I \cup \widehat{\Phi}_J$. Since $\widehat{\Phi}_I \subseteq \text{conv}(\Delta_I)$ (resp. $\widehat{\Phi}_J \subseteq \text{conv}(\Delta_J)$) and since $\text{conv}(\Delta_I)$ and $\text{conv}(\Delta_J)$ are disjoint compact sets, a converging sequence of elements in $\widehat{\Phi}$ eventually lives either in $\text{conv}(\Delta_I)$ or in $\text{conv}(\Delta_J)$, and so converges to a limit point either in $E(\Phi_I)$ or in $E(\Phi_J)$. \square

Corollary 2.16. *If (Φ, Δ) is an affine based root system, then $E(\Phi)$ is finite. Moreover, if (Φ, Δ) is affine and irreducible, then $E(\Phi)$ is a singleton (and is equal to \widehat{Q}).*

Proof. By definition, if (Φ, Δ) is an irreducible and affine based root system, of rank n , then the signature of B is $(n - 1, 0)$. So \widehat{Q} is a point in this case, and since Φ is infinite, $E \neq \emptyset$ by Proposition 2.14. Since $E \subseteq \widehat{Q}$, we get that $E = \widehat{Q}$ and contains a single element. The proof follows then from Proposition 2.15. \square

3. GEOMETRIC ACTION OF THE COXETER GROUP W ON THE LIMIT ROOTS

In this section, we address some natural questions concerning the pictures we obtained: can W act on the set E of limit roots? Where does the “fractal phenomenon” come from? As in the previous section, we fix an infinite based root system (Φ, Δ) in (V, B) with associated Coxeter system (W, S) , and we assume that Δ is a basis for V . We recall that V_1 denotes the affine hyperplane containing the simple roots (seen as points), with direction V_0 , and $E = E(\Phi)$ the set of limit roots, i.e., the accumulation set of the normalized roots.

3.1. Geometric action.

It is clear that the geometric action of the group W on V does not restrict to an action on V_1 . However, using the normalization map, it induces a natural action on a part of V_1 . We consider the set

$$\begin{aligned} D &:= \bigcap_{w \in W} w(V \setminus V_0) \cap V_1 \\ &= V_1 \setminus \bigcup_{w \in W} w(V_0). \end{aligned}$$

We define an action of W on D as follows: for $x \in D$ and $w \in W$, set

$$w \cdot x := \widehat{w(x)}.$$

Since any element of D stays in the complement of V_0 after action of any $w \in W$, it is straightforward to check that this is a well-defined action of W , and that any $w \in W$ acts continuously on D . The set D is the complement in V_1 of an infinite union of hyperplanes, and is the maximal subset of V_1 on which W acts naturally. The following statement is the main motivation to consider this action.

Proposition 3.1.

- (i) $\widehat{\Phi}$ and E are contained in D ;
- (ii) $\widehat{\Phi}$ and E are stable by the action of W ; moreover $\widehat{\Phi} = W \cdot \Delta$;
- (iii) the topological closure $\widehat{\Phi} \sqcup E$ of $\widehat{\Phi}$ is stable by the action of W .

Proof. First, we show that $\widehat{\Phi}$ is contained in D and that $\widehat{\Phi}$ is stable by W . Recall that Φ lies in $V \setminus V_0$ and is stable under the action of W on V . Let $\widehat{\beta} \in \widehat{\Phi}$ and $w \in W$. Since $\mathbb{R}\beta = \mathbb{R}\widehat{\beta}$, we have $\mathbb{R}w(\widehat{\beta}) = w(\mathbb{R}\widehat{\beta}) = w(\mathbb{R}\beta) = \mathbb{R}w(\beta)$. Therefore, since $w(\beta)$ is a root, the point $w(\widehat{\beta})$ lies in $V \setminus V_0$ and $w \cdot \widehat{\beta} = \widehat{w(\beta)} = \widehat{w(\beta)}$ is a point of $\widehat{\Phi}$. It remains to prove that:

$$\forall w \in W, \forall x \in E, w(x) \in V \setminus V_0 \text{ and } w \cdot x = \widehat{w(x)} \in E.$$

Take $x \in E$, and $w \in W$. Consider an injective sequence $(\rho_n)_{n \in \mathbb{N}}$ in Φ^+ , such that $(\widehat{\rho_n})$ converges to x . Then, for all n , $w(\widehat{\rho_n})$ is in $V \setminus V_0$, as well as its limit $w(x)$. Since w and $\widehat{\cdot}$ are continuous, we have:

$$w \cdot \rho_n = \widehat{w(\rho_n)} = \widehat{w(\widehat{\rho_n})} \xrightarrow{n \rightarrow \infty} \widehat{w(x)} = w \cdot x.$$

So $w \cdot x$ is a limit point of the set $\widehat{\Phi}$, i.e., lies in E . The fact that $\widehat{\Phi} = W \cdot \Delta$ follows directly from the definition of Φ , $\Phi = W(\Delta)$. So (i) and (ii) are proved, and (iii) is a trivial consequence of them. \square

It is possible to restrict the action of W on D to the simplest connected component of D , which is the convex set:

$$D^+ := \bigcap_{w \in W} w(V_0^+) \cap V_1 \subseteq V_0^+ \cap V_1.$$

This convex set is still stable by the action of W and does not contain any element of $\widehat{\Phi}$ (since $s_\alpha(\alpha) = -\alpha$). However D^+ is of interest to study the action of W on E (see for instance the proof of Proposition 3.7):

Proposition 3.2. *The set E is contained in D^+ . In particular, $W(E)$ and $W \cdot E$ are subsets of V_0^+ , that is, $|w(x)|_1 > 0$, for all $x \in E$ and $w \in W$.*

Proof. We need to prove that for any $w \in W$ and $x \in E$, the image $w(x)$ stays in V_0^+ . Using a direct induction on the length of w —that is, the minimal length of the word w in the alphabet S —this is a consequence of the following fact:

$$\forall \alpha \in \Delta, \forall x \in E, s_\alpha(x) \in V_0^+.$$

Indeed we know from Proposition 3.1 that E is stable by the action of W . Let us prove this fact. Take $x \in E$, and $\alpha \in \Delta$. Consider an injective sequence $(\rho_n)_{n \in \mathbb{N}}$ in Φ^+ , such that $(\widehat{\rho_n})$ converges to x . We can suppose that (ρ_n) does not contain α . Then, for all n , $s_\alpha(\rho_n)$ is in Φ^+ , and in particular in V_0^+ , so $s_\alpha \cdot \widehat{\rho_n}$ is also in V_0^+ , as well as its limit $s_\alpha \cdot x$. \square

Remark 3.3. Although we do not know any better description of D^+ than the definition given here, M. Dyer suggested that if we replace V_0^+ with $\text{cone}(\Delta)$ in the definition of D^+ , then the convex set we obtain may be equal to $\text{conv}(E)$. We do not know a proof of this statement or even if D^+ is equal to this new convex set.

Remark 3.4. Obviously, this action is not faithful when Φ is an affine root system (since E is finite, see Corollary 2.16). We do not know whether this action is faithful for infinite irreducible non affine root systems of rank ≥ 3 .

The aim behind studying the set $E \subseteq V_1$ is to be able to represent “limit points of roots” in an affine space. Now we can also study an action of W on these limit points, and it turns out that the action of the reflections of W on E (and more generally on $Q \cap D$) is geometric in essence:

Proposition 3.5 (A geometric description).

- (i) Let $\rho \in \Phi$, and $x \in D \cap Q$. Denote by $L(\widehat{\rho}, x)$ the line containing $\widehat{\rho}$ and x . Then:
 - (a) if $B(\rho, x) = 0$, then $L(\widehat{\rho}, x)$ intersects Q only in x , and $s_\rho \cdot x = x$;
 - (b) if $B(\rho, x) \neq 0$, then $L(\widehat{\rho}, x)$ intersects Q in the two distinct points x and $s_\rho \cdot x$.
- (ii) Let $\rho_1 \neq \rho_2 \in \Phi$, $x \in L(\widehat{\rho}_1, \widehat{\rho}_2) \cap Q$ and $w \in W$. Then $w \cdot x \in L(w \cdot \widehat{\rho}_1, w \cdot \widehat{\rho}_2) \cap Q$.

To visualize this proposition, we refer to Figure 7 in which we draw the images of the two yellow points x and y on the edge $[\beta, \gamma]$ of $\text{conv}(\Delta)$ under the action of s_α and of $s_\beta s_\alpha$.

Proof. (i) Let $\rho \in \Phi$, and $x \in D \cap Q$. First, we know that $s_\rho(x)$ and $s_\rho \cdot x$ belong to Q . Note also that $s_\rho \cdot x \in \text{span}(\rho, x) \cap V_1 = L(\widehat{\rho}, x)$, so x and $s_\rho \cdot x$ are always intersection points of $L(\widehat{\rho}, x)$ with Q . Since s_ρ is a reflection and $x \in V_1$, we have

$$|L(\widehat{\rho}, x) \cap Q| = 1 \Rightarrow s_\rho \cdot x = x \Leftrightarrow s_\rho(x) = x \Leftrightarrow B(\rho, x) = 0.$$

Consider an element $u_\lambda = \lambda\hat{\rho} + (1-\lambda)x$ in $L(\hat{\rho}, x) \cap Q$. Then

$$\begin{aligned} 0 = q(u_\lambda) &= \lambda^2 q(\hat{\rho}) + 2\lambda(1-\lambda)B(\hat{\rho}, x) + (1-\lambda)^2 q(x) \\ &= \lambda^2 q(\hat{\rho}) + 2\lambda(1-\lambda)B(\hat{\rho}, x). \end{aligned}$$

Hence $L(\hat{\rho}, x)$ intersects Q in at most two points. In the case that $B(\rho, x) \neq 0$, the two points $x, s_\rho \cdot x$ are distinct, and thus the third sentence of the present proof yields that $L(\hat{\rho}, x) \cap Q = \{x, s_\rho \cdot x\}$. In the case that $B(\rho, x) = 0$, then $0 = q(u_\lambda) = \lambda^2 q(\hat{\rho})$, so $u_\lambda = x$ and $L(\hat{\rho}, x) \cap Q = \{x\}$.

(ii) Let $\rho_1 \neq \rho_2 \in \Phi$. For any $x \in E \subseteq Q$ and $w \in W$, since $q(x) = B(x, x) = B(w(x), w(x)) = q(w(x))$, it follows that both $w(x)$ and $w \cdot x$ are in Q . Therefore, we only need to check that if $x \in L(\hat{\rho}_1, \hat{\rho}_2)$ then $w \cdot x \in L(w \cdot \hat{\rho}_1, w \cdot \hat{\rho}_2)$.

Let $x \in L(\hat{\rho}_1, \hat{\rho}_2) \cap Q$. Note that $x \in E$ as explained in §2.3. Let $w \in W$ and set $\rho'_1 := w(\rho_1)$ and $\rho'_2 := w(\rho_2)$. Then $w \cdot \rho_1 = \hat{\rho}'_1$ and $w \cdot \rho_2 = \hat{\rho}'_2$. Since $x \in L(\hat{\rho}_1, \hat{\rho}_2) \subseteq \text{span}(\rho_1, \rho_2)$, it follows that $w(x)$ is a point in the plane P spanned by ρ'_1 and ρ'_2 . Since $P \cap V_1 = L(\hat{\rho}'_1, \hat{\rho}'_2)$, and $w \cdot x \in P \cap V_1$, it follows that $w \cdot x \in L(\hat{\rho}'_1, \hat{\rho}'_2) \cap Q$ as required. \square

As we can see on Figure 7, when $L(\hat{\rho}, x)$ intersects Q in two points, one of them is closest to $\hat{\rho}$ than the other; the closest point to $\hat{\rho}$ is then *visible from* $\hat{\rho}$. Let us describe this phenomenon more precisely.

Definition 3.6. We say that $x \in \hat{Q}$ is *visible from* $v \in V_1$ if the segment $[v, x]$ intersects \hat{Q} in only one point, which has to be x ; in other words, x is visible from v if and only if $[v, x] \cap Q = \{x\}$.

The following proposition sums up what we observe in Figure 7.

Proposition 3.7. *Let $x \in E$ and $\rho \in \Phi$. Then,*

- (i) x is visible from $\hat{\rho}$ if and only if $B(\hat{\rho}, x) \geq 0$;
- (ii) if $\rho \in \Phi^+$, x is visible from $\hat{\rho}$ if and only if $B(\rho, x) \geq 0$;
- (iii) if $\rho \in \Phi^+$ and $w \in W$ such that $w(\rho) \in \Phi^+$, then x is visible from $\hat{\rho}$ if and only if $w \cdot x$ is visible from $w \cdot \hat{\rho}$.

Proof. If $B(\rho, x) = 0$, then by Proposition 3.5, $L(\hat{\rho}, x) \cap Q = \{x\}$, so both statements are obviously true. Assume now that $B(\rho, x) \neq 0$, which is equivalent by Proposition 3.5 to $|L(\hat{\rho}, x) \cap Q| = 2$. First suppose that $\rho \in \Phi^+$. So the linear forms $B(\rho, \cdot)$ and $B(\hat{\rho}, \cdot)$ are of the same sign. In this case, the proof follows from the equality $s_\rho(x) = x - 2B(\rho, x)\rho$, which implies that when $B(\rho, x) < 0$, $s_\rho(x)$ is in the interior of cone(x, ρ), and when $B(\rho, x) > 0$, x is in the interior of cone($s_\rho(x), \rho$). Now, if $\rho \in \Phi^-$, apply the above case with $-\rho \in \Phi^+$, which proves (i) and (ii).

For (iii), we use Proposition 3.2: $E \subseteq D^+$, therefore $w(x) \in V_0^+$. So $|w(x)|_1 > 0$, which means that $B(w(x), \cdot)$ and $B(w \cdot x, \cdot)$ have the same sign. Since we have $w(\rho) \in \Phi^+ \subseteq V_0^+$, $B(\cdot, w(\rho))$ and $B(\cdot, w \cdot \hat{\rho})$ have also the same sign. We conclude using (i) and (ii). \square

We end this discussion by completing the example of infinite dihedral groups.

Example 3.8 (The case of infinite dihedral groups). Let (Φ, Δ) be of rank 2 as in Example 2.1 (illustrated in Figure 2 and Figure 3). Recall that $\Delta = \{\alpha, \beta\}$ with $B(\alpha, \beta) \leq -1$ and that V_1 is the line $L(\alpha, \beta)$. Let $x \in \hat{Q}$ such that $B(x, \alpha) \geq 0$, so x is visible from α . Since $s_\beta \cdot \alpha \in V_1 = L(\alpha, \beta)$ and $s_\beta \cdot \beta = \beta \in V_1 = L(\alpha, \beta)$,

it follows from Proposition 3.5 (ii) that $s_\beta \cdot x \in L(\alpha, \beta) \cap Q$, and similarly that $s_\alpha \cdot x \in L(\alpha, \beta) \cap Q$. Now since $L(\alpha, \beta) = L(x, \alpha) = L(x, \beta)$, it follows from Proposition 3.5 (i) that either $x = s_\beta \cdot x$ and $x = s_\alpha \cdot x$ or $L(\alpha, \beta) \cap Q = \{x, s_\beta \cdot x\} = \{x, s_\alpha \cdot x\}$ has cardinality 2. Thus in both case, $s_\beta \cdot x = s_\alpha \cdot x$, and this point is visible from β . So the segment $[\alpha, x]$ intersects \widehat{Q} only in x .

We endow the line $L(\alpha, \beta) = V_1$ with the total order on \mathbb{R} by orienting it from α to β . So from the above discussion we get that

$$\alpha < x \leq s_\alpha \cdot x < \beta.$$

We are particularly interested in the following partition of the segment $[\alpha, \beta]$:

$$[\alpha, \beta] = [\alpha, x] \sqcup [x, s_\alpha \cdot x] \sqcup (s_\alpha \cdot x, \beta].$$

It is straightforward computations⁶ to prove the following nice geometric description of the action of s_α and s_β on $\widehat{\Phi} \sqcup E$:

$$\alpha < s_\alpha \cdot \beta < s_\alpha s_\beta \cdot \alpha < s_\alpha s_\beta s_\alpha \cdot \beta < \cdots < (s_\alpha s_\beta)^n \cdot \alpha < (s_\alpha s_\beta)^n s_\alpha \cdot \beta < \cdots < x$$

and

$$s_\beta \cdot x = s_\alpha \cdot x < \cdots < s_\beta (s_\alpha s_\beta)^n \cdot \alpha < (s_\beta s_\alpha)^n \cdot \beta < \cdots < s_\beta s_\alpha \cdot \beta < s_\beta \cdot \alpha < \beta.$$

Moreover, we note the following two facts:

- the bilinear form B is positive on $[\alpha, x[\times [\alpha, x[$ and $]s_\alpha \cdot x, \beta] \times]s_\alpha \cdot x, \beta]$, and negative on $[\alpha, x[\times]s_\alpha \cdot x, \beta]$;
- the depth (dp) is increasing on $[\alpha, x[\cap \widehat{\Phi}$ and decreasing on $]s_\alpha \cdot x, \beta] \cap \widehat{\Phi}$ (where we set $\text{dp}(\widehat{\rho}) := \text{dp}(\rho)$ for $\rho \in \Phi^+$).

3.2. Fractal description of E .

We know that E is contained in \widehat{Q} (Theorem 2.7), but we could obtain a more precise inclusion, using the action of W . First of all, E is also included in $\text{conv}(\Delta)$. Now consider the example illustrated in Figure 7, and suppose that we can act by W on \widehat{Q} , with the action of §3.1, i.e., that $\widehat{Q} \subseteq D$ (this is not true in general). Thus, as there are no limit points in the red arc which is outside the triangle $\text{conv}(\Delta)$, there are also no limit points on its image by s_α , which is the smaller red arc on the bottom left; and not either in the subsequent image by s_β . So E seems to be contained in a fractal self-similar subset F of \widehat{Q} , obtained by removing from \widehat{Q} all these iterated arcs. The rank 4 pictures are even more convincing of this property; see in particular Figures 9 and 1(b), where the fractal F obtained (an ellipsoid cut out by an infinite number of planes) looks like an Apollonian gasket drawn on \widehat{Q} .

We conjecture that E is actually equal to F . In the particular case where \widehat{Q} is contained in $\text{conv}(\Delta)$ —e.g., Figure 8, and all the cases of a rank 3 Coxeter group with classical representation, as Figures 1(a) and 5—, this means that E fills up \widehat{Q} . We also conjecture the following more precise property:

Conjecture 3.9. *We say that $\Delta_I \subseteq \Delta$ is generating if $\widehat{Q}_I := \widehat{Q} \cap \text{span}(\Delta_I)$ is included in $\text{conv}(\Delta_I)$. Then we have the following properties:*

- (i) *if $\Delta_I \subseteq \Delta$ is generating, then $E(\Phi_I) = \widehat{Q}_I$;*

⁶See for instance [Fu12, p.5 Eq. (1.2)] for details.

(ii) the set E is the topological closure of the fractal self-similar subset F_0 of \widehat{Q} defined by:

$$F_0 := W \cdot \left(\bigcup_{\substack{\Delta_I \subseteq \Delta \\ \Delta_I \text{ generating}}} \widehat{Q}_I \right).$$

Example 3.10. In the example pictured in Figure 9, the set F_0 defined above is the fractal constituted by the infinite union of all the circles. The set F (and, conjecturally, the set E) is the complement, on the red sphere, of the union of the associated open “disks”.

4. CONSTRUCTION OF A DENSE SUBSET OF THE LIMIT ROOTS FROM DIHEDRAL REFLECTION SUBGROUPS

As in the previous sections, we fix an infinite based root system (Φ, Δ) in (V, B) with associated Coxeter system (W, S) , assuming that Δ is a basis for V , and we denote by $E = E(\Phi)$ the accumulation set of its normalized roots. In this section, we construct a nice countable subset of E , easy to describe, and we show its density in E .

4.1. Main Theorem: E is obtained from dihedral reflection subgroups.

Consider two distinct positive roots ρ_1 and ρ_2 , and define the dihedral reflection subgroup $W' := \langle s_{\rho_1}, s_{\rho_2} \rangle$ and

$$\Phi' := \{ \rho \in \Phi \mid s_{\rho} \in W' \}.$$

From Proposition 1.5, we know that Φ' is a based root system of rank 2, associated to the dihedral group W' . Denote by α, β its simple roots, and suppose that $B(\alpha, \beta) \leq -1$, i.e., that W' is infinite and the plane $\text{span}(\rho_1, \rho_2)$ intersects $Q \setminus \{0\}$ (see Proposition 1.5 (ii)(a)). Similarly, the line $L(\widehat{\rho}_1, \widehat{\rho}_2)$ intersects \widehat{Q} inside V_1 . Denote by u, v the elements of the intersection $L(\widehat{\rho}_1, \widehat{\rho}_2) \cap \widehat{Q}$ (where $u = v$ if and only if $B(\alpha, \beta) = -1$). Because of the rank 2 picture (see Examples 2.1 and 3.8 and §2.3), we know that the set $E(\Phi')$ of limit points of normalized roots of W' is equal to the finite set $\{u, v\}$. This leads to the following natural definition.

Definition 4.1. Let E_2 be the subset of E formed by the union of the sets $E(\Phi')$, where (Φ', Δ') is any based root subsystem of rank 2 in (Φ, Δ) . Equivalently,

$$E_2 := \bigcup_{\rho_1, \rho_2 \in \Phi^+} L(\widehat{\rho}_1, \widehat{\rho}_2) \cap \widehat{Q},$$

where $L(\widehat{\rho}_1, \widehat{\rho}_2)$ denotes the line containing $\widehat{\rho}_1$ and $\widehat{\rho}_2$.

Note that E_2 is countable, and geometrically easier to describe than the whole set E . Indeed, it is constructed from Φ with intersections rather than with limits. Moreover, $E_2 = E$ when (Φ, Δ) is affine, or of rank 2. Figure 10 gives an example of construction of some points of E_2 . Surprisingly, E_2 still carries all the information of E , as implied by the following theorem.

Theorem 4.2. *Let Φ be an (infinite) based root system, E the set of limit points of its normalized roots. Then the union E_2 of all limit roots arising from dihedral reflection subgroups is dense in E .*




FIGURE 10. Geometric construction of E_2 , for the based root system with labels 4, 4, 4 (roots of depth ≤ 5 in blue, some elements of E_2 are in yellow diamonds).




FIGURE 11. Construction of a sequence of elements of E_2 (x_n , in yellow diamonds) from a sequence of elements of $\widehat{\Phi}$ (u_n , in blue), both converging to $\ell \in E$.

Remark 4.3. Since E is a limit set, it is closed, so the theorem implies that E is the closure of E_2 . Note also that if we define similarly E_k as the subset of limit points arising from rank k reflection subgroups (with $k \geq 2$), then E_k is dense in E too (since it contains E_2). However, this is not true if we consider only parabolic subgroups. For instance, any proper parabolic subgroup of an irreducible affine Coxeter group is finite and therefore does not provide any limit point (contradicting Corollary 2.16).

It turns out that an even smaller subset than E_2 is enough to recover E . Define E_2° by using the lines passing through a normalized root and a *simple* root:

$$E_2^\circ := \bigcup_{\substack{\alpha \in \Delta \\ \rho \in \Phi^+}} L(\alpha, \widehat{\rho}) \cap \widehat{Q} \subseteq E_2.$$

Proposition 4.4. *The set E_2 is the W -orbit of E_2° for the action defined in §3.1:*

$$E_2 = W \cdot E_2^\circ.$$

Proof. Let $x \in E_2$ and $\rho_1, \rho_2 \in \Phi$ such that $x \in L(\widehat{\rho}_1, \widehat{\rho}_2)$. Since $\Phi = W(\Delta)$, there is w in W such that $\alpha := w(\rho_1) \in \Delta$. By Proposition 3.5 (ii), we have that $w \cdot x \in L(\widehat{\alpha}, \widehat{w(\rho_2)}) \cap \widehat{Q}$, which proves that $w \cdot x \in E_2^\circ$. \square

Theorem 4.2 is in fact a direct consequence of the following stronger property.

Theorem 4.5. *Let Φ be an (infinite) based root system, E its set of limit points of normalized roots, and E_2° the subset of E defined above. Then E_2° is dense in E .*

The rest of this section is devoted to the proof of Theorem 4.5. In regard to Proposition 2.15, we assume that the based root system (Φ, Δ) , and the Coxeter system (W, S) , are irreducible.

Consider an element ℓ of E . By definition, there exists a sequence of normalized roots $(u_n)_{n \in \mathbb{N}} \subseteq \widehat{\Phi}$ converging to ℓ . We want to construct, from (u_n) , a new

sequence of elements of E_2° converging to ℓ as well. The geometric idea is quite simple, as illustrated in Figure 11: we construct x_n as the “right” intersection point of the line $L(\alpha, u_n)$ with Q , for an $\alpha \in \Delta$ well chosen. The core of the proof is to show that it is always possible to find such a simple root that makes the construction work.

We proceed in two steps, detailed in the next subsections.

4.2. First step of proof: when the limit root is outside the radical.

Denote by $V^\perp = \{v \in V \mid B(v, u) = 0, \forall u \in V\}$ the radical of the bilinear form B .

Proposition 4.6. *Let $\ell \in E$. If ℓ is not in the radical V^\perp , then ℓ is in the closure of E_2° .*

This proposition is a consequence of the following construction.

Lemma 4.7. *Let $(\rho_n)_{n \in \mathbb{N}}$ be a sequence in Φ^+ , such that $\widehat{\rho}_n$ converges to a limit ℓ in E . Suppose that there exists α in Δ , such that $B(\alpha, \ell) \neq 0$. Then, there exists a sequence $(x_n)_{n \in \mathbb{N}}$ converging to ℓ , such that, for any n large enough, x_n lies in the intersection $L(\alpha, \widehat{\rho}_n) \cap Q$.*

If $\ell \in E$ and $\ell \notin V^\perp$, then we are in the hypothesis of the lemma since Δ spans V . Moreover, by definition of E_2° , the constructed sequence (x_n) eventually lies in E_2° . Thus, Proposition 4.6 follows directly from the lemma, which we prove now.

Proof. Since $B(\alpha, \ell) \neq 0$, we get from Proposition 3.5 (ii) that $L(\alpha, \ell) \cap Q$ contains two points (see Figure 11). Consider an element $x \in V_1 \setminus \{\alpha\}$, and let us find the points of $L(\alpha, x) \cap Q$. Take an element u_λ in the line $L(\alpha, x)$

$$u_\lambda = \lambda\alpha + (1 - \lambda)x, \text{ with } \lambda \in \mathbb{R}.$$

Then a quick computation gives

$$(4.1) \quad q(u_\lambda) = 0 \Leftrightarrow q(\alpha - x)\lambda^2 + 2B(x, \alpha - x)\lambda + q(x) = 0$$

(where q is the quadratic form of B). For $x = \ell$, we know that Equation (4.1) has two solutions for λ . So by continuity, there exists a neighbourhood Ω_ℓ of ℓ in V_1 , such that, for any $x \in \Omega_\ell$, Equation (4.1) has two solutions as well (and then $q(\alpha - x) \neq 0$). Computing these solutions in function of x , we find, after simplification,

$$\lambda_\pm(x) = \frac{q(x) - B(\alpha, x) \pm \sqrt{B(\alpha, x)^2 - q(x)}}{q(\alpha - x)},$$

and the two intersection points of $L(\alpha, x)$ with Q (for $x \in \Omega_\ell$) are given by $u_+(x)$ and $u_-(x)$, where

$$u_\varepsilon(x) = \lambda_\varepsilon(x)\alpha + (1 - \lambda_\varepsilon(x))x, \text{ for } \varepsilon = + \text{ or } -.$$

For $x = \ell$, the intersection points are obviously ℓ and $s_\alpha \cdot \ell$; so, in regard to Proposition 3.7, either $u_+(\ell) = \ell$ if $B(\alpha, \ell) > 0$ (that is, when ℓ is visible from α), or $u_-(\ell) = \ell$ if $B(\alpha, \ell) < 0$ (that is, when $s_\alpha \cdot \ell$ is visible from α). Define, for $x \in \Omega_\ell$,

$$f_\alpha(x) = u_\varepsilon(x) = \lambda_\varepsilon(x)\alpha + (1 - \lambda_\varepsilon(x))x,$$

where $\varepsilon = +$ if $B(\alpha, \ell) > 0$, and $\varepsilon = -$ if $B(\alpha, \ell) < 0$. Then,

- $f_\alpha(\ell) = \ell$;

- f_α is a continuous map on Ω_ℓ ;
- for all $x \in \Omega_\ell$, $f_\alpha(x) \in L(\alpha, x) \cap Q$.

Now consider a sequence $(\rho_n)_{n \in \mathbb{N}}$ in Φ^+ , such that $\widehat{\rho_n}$ converges to ℓ . For n large enough, $\widehat{\rho_n} \in \Omega_\ell$, and we set $x_n = f_\alpha(\widehat{\rho_n})$. Then $x_n \in L(\alpha, \widehat{\rho_n}) \cap Q$, and by continuity of f_α , x_n converges to $f_\alpha(\ell) = \ell$. \square

4.3. Second step of proof: when the limit root is inside the radical.

To finish the proof, we only need to deal with the case where ℓ is in the radical V^\perp of the form B . Since ℓ is also in $\text{cone}(\Delta)$, the following proposition, which is the same as [Kra09, Lemma 6.1.1], implies that this case can only happen when Φ is an affine based root system.

Proposition 4.8. *Let Φ be an (infinite) irreducible based root system. If Φ is not affine, then*

$$V^\perp \cap \text{cone}(\Delta) = \{0\} .$$

The proof of this statement is an application of the Perron-Frobenius theorem; we refer to [Kra09, Lemma 6.1.1] for details.

So we focus on the last remaining case: Φ is an irreducible affine root system. In this case, by Corollary 2.16, E is a singleton, so it is sufficient to check that E_2° is nonempty. But E_2° is empty if and only if E_2 is empty, if and only if any reflection subgroup of W of rank 2 is finite; and this cannot happen because of the following classical property.

Lemma 4.9. *Let W be a Coxeter group. If every reflection subgroup of rank 2 of W is finite, then W is finite.*

Proof. We use the concept of small roots, described for instance in [BB05, §4.7]; all the references in this proof are to this book. If every reflection subgroup of rank 2 of W is finite, then any covering edge of the root poset on Φ^+ (see Def. 4.6.3) is a short edge (see §4.7), and any root is a small root. But it is known that the set of small roots is finite (Theorem 4.7.3, taken from Brink-Howlett [BH93]), so Φ is finite and W must be finite as well. \square

This concludes the proof of Theorem 4.5 and therefore the proof of Theorem 4.2.

5. HOW TO CONSTRUCT THE LIMIT ROOTS FOR REFLECTION SUBGROUPS

In this section, we explain how to slightly extend the definition of E so that it holds for a reflection subgroup W' of a Coxeter group W . To do so we extend the definition of limit roots such that it applies even when the set of simple roots is not a basis. In the last subsection, we also detail the relations between the set of limit roots associated to a reflection subgroup of W (especially to a parabolic subgroup) and the set E associated to W .

5.1. Root subsystems of a based root system.

First, we define the notion of a root subsystem. Let (Φ, Δ) be a based root system of (V, B) with associated Coxeter system (W, S) , according to Definition 1.2. Consider a finitely generated reflection subgroup W' of W , together with its canonical set S' of Coxeter generators, as explained in §1.2. There is a natural way to construct a root system for W' from the one for W . If

$$\Phi' := \{\rho \in \Phi \mid s_\rho \in W'\} \quad \text{and} \quad \Delta' := \{\rho \in \Phi \mid s_\rho \in S'\},$$

then (Φ', Δ') is a based root system of (V, B) with associated Coxeter system (W', S') (see [BD10, Lemma 3.5] for details). Note that $\Delta' \subseteq \Phi^+$ does not have to intersect Δ . We call (Φ', Δ') a *based root subsystem of (Φ, Δ)* . In §2.3, we already mentioned the specific case of *standard parabolic* root subsystems, which correspond to standard parabolic Coxeter subgroups of W .

The simplest definition of $E(\Phi')$ in this setting is to consider the subset of $E(\Phi)$ consisting of all the points which are limits of sequences in $\widehat{\Phi'}$. A natural question arises: is this definition consistent with the intrinsic one, when we consider (Φ', Δ') as a root system by itself? The first problem here is that, in §2, we only defined E when the set of simple roots is linearly independent, and this is not always the case for (Φ', Δ') , as illustrated in the following example.

Example 5.1. Consider the Coxeter group W of rank 3 with $S = \{s, t, u\}$ and $m_{s,t} = m_{t,u} = m_{s,u} = 4$ (whose Coxeter diagram is on Figure 10). Let (V, B) be the classical geometric representation of W . Consider the reflection subgroup W' generated by $S' = \{s, u, tst, tut\}$. It is easy to check that its simple system Δ' is $\{\alpha_s, \alpha_u, t(\alpha_s), t(\alpha_u)\}$ and satisfies the conditions of Definition 1.2 (Condition (i) is critical here). So (W', S') is a Coxeter group of rank 4 and V is a geometric W' -module of dimension 3 with based root system (Φ', Δ') where Φ' is the set of roots ρ in Φ such that $s_\rho \in W'$. Consider now the matrix $A = (B(\alpha_x, \alpha_y))_{x,y \in S'}$, written accordingly to $(\alpha_s, \alpha_u, t(\alpha_s), t(\alpha_u))$:

$$A = \begin{pmatrix} 1 & -\frac{\sqrt{2}}{2} & 0 & -1 - \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} & 1 & -1 - \frac{\sqrt{2}}{2} & 0 \\ 0 & -1 - \frac{\sqrt{2}}{2} & 1 & -\frac{\sqrt{2}}{2} \\ -1 - \frac{\sqrt{2}}{2} & 0 & -\frac{\sqrt{2}}{2} & 1 \end{pmatrix}.$$

This matrix satisfies Equation (1.2), so we can construct (as in §1.4) the canonical geometric W -module (V_A, B_A) , of dimension 4, associated to W' . Observe that (V_A, B_A) is not the classical geometric representation of W' .

Such examples explain why a satisfying definition of a based root system should not require the simple roots to form a basis, but only to be positively independent (Condition (i) in Definition 1.2). They also raise the question of the possible relations between (V, B) and (V_A, B_A) as W' -modules (see §5.3).

5.2. How to cut the rays of roots in general: hyperplanes transverse to Φ^+ .

In §2 we constructed the set E of limit roots from *normalized* roots, by cutting the rays of positive roots with the hyperplane V_1 . To be able to define properly E in the case where Δ is not necessarily a basis, we need to find analogues for the cutting hyperplane V_1 and the normalization map $\widehat{\cdot}$.

Even when Δ is a basis, it is natural to ask the following question: since $\widehat{\Phi}$ is an affine representative of $\mathbb{P}\Phi$ obtained by “cutting” Φ by a particularly well-chosen affine hyperplane, what happens if we change the affine hyperplane, i.e., the representative of $\mathbb{P}\Phi$? For instance, in the case of an affine dihedral group (see Example 1.4), if we cut the rays of Φ by an affine line directed by $\alpha + \beta$ (thus parallel to the radical of B), then the points representing the roots are not all contained in the convex hull of the points representing the simple roots, and these normalized roots diverge to infinity. This fact suggests the following general definition, that also takes care of the case where Δ is not a basis (but still positively independent).

Definition 5.2. Let (Φ, Δ) be a based root system in (V, B) . We call an affine hyperplane H *transverse to Φ^+* if the intersection of H with any ray $\mathbb{R}^+\alpha$ directed by a simple root $\alpha \in \Delta$ is a nonzero point, i.e., if $\mathbb{R}^{+*}\alpha \cap H$ is a singleton for all $\alpha \in \Delta$.

If Δ is a basis for V , the affine hyperplane V_1 we used to cut the rays of Φ is obviously transverse. In the case where Δ is only positively independent, the definition of V_1 does not work (it could be equal to V). But transverse hyperplanes still exist. Consider a linear hyperplane H_0 , separating Φ^+ and Φ^- : it exists because Δ is positively independent (see Remark 1.3). Then the affine hyperplane $H := \sum_{\alpha \in \Delta} \alpha + H_0$ is transverse.

Let H be a transverse hyperplane directed by a hyperplane H_0 . Let H_0^+ be the open halfspace supported by H_0 and containing H (H is parallel to H_0 and 0 cannot be in H by definition). So H_0^+ contains also Δ and therefore Φ^+ , which is a subset of $\text{cone}(\Delta)$. In particular, we obtain easily the following statement: an affine hyperplane H is transverse if and only if $\mathbb{R}^{+*}\beta \cap H$ is a singleton for all $\beta \in \Phi^+$.

Like for V_1 , a normalization map can be applied to Φ :

$$\pi_H : V \setminus H_0 \rightarrow H$$

where $\pi_H(v)$ is the intersection point of $\mathbb{R}v$ with H . For instance, when Δ is a basis, $\pi_{V_1}(v) = \hat{v}$. Note that since H is transverse, $\pi_H(\Phi)$ is contained in the polytope $\text{conv}(\pi_H(\Delta))$. Denote by $E(\Phi, H)$ the set of accumulation points of $\pi_H(\Phi)$. The following straightforward proposition states that the topology of our object of study, $E(\Phi)$, does not depend on the choice of the transverse hyperplane chosen to cut the rays of Φ , as suggested by the point of view of projective geometry.

Proposition 5.3. *Let (Φ, Δ) be a based root system, and let H, H' be two hyperplanes, transverse to Φ^+ . Then π_H induces a homeomorphism from $\text{conv}(\pi_{H'}(\Delta))$ to $\text{conv}(\pi_H(\Delta))$, whose inverse is the restriction of $\pi_{H'}$. Moreover, π_H maps (bijectively) $\pi_{H'}(\Phi)$ to $\pi_H(\Phi)$ and $E(\Phi, H')$ to $E(\Phi, H)$.*

Proof. For $v \in V$, $\pi_H(v)$ and $\pi_{H'}(v)$ (when they are defined) are colinear to v . As Δ is contained in $V \setminus H_0$, so are $\pi_{H'}(\Delta)$ and $\text{conv}(\pi_{H'}(\Delta))$. So, the mentioned restrictions of π_H and $\pi_{H'}$ are, by construction, continuous, mutual inverses, and swap $\pi_{H'}(\Phi)$ and $\pi_H(\Phi)$. Consequently, they also swap the sets of limit points of these two sets, i.e., $E(\Phi, H')$ and $E(\Phi, H)$. \square

Remark 5.4. The geometric action of W defined in §3.1 does not depend on the choice of the transverse hyperplanes employed to cut Φ : if H and H' are two hyperplanes that are transverse to Φ^+ , then the map $\pi_{H'}$ induces by restriction an isomorphism of W -sets from $\pi_H(\Phi) \sqcup E(\Phi, H)$ to $\pi_{H'}(\Phi) \sqcup E(\Phi, H')$.

Remark 5.5. Changing the transverse hyperplane can still change the geometry overall. For instance, consider a rank 3 based root system with B of signature $(2, 1)$, so that Q is a circular cone. Let H be a transverse affine plane. In general, the intersection $Q \cap H$ of H with the isotropic cone Q can be an ellipse, a parabola or an hyperbola. In the case of an hyperbolic cut, only one branch intersects $\text{conv}(\pi_H(\Delta))$. In a following paper [DHR13], we prove and use the fact that for a hyperbolic group of rank 3, there always exists a hyperplane H , transverse to Φ^+ , such that $Q \cap H$ is a circle (the analogous property is actually valid for hyperbolic groups of higher rank).

For now, we apply this result to explain in the following subsection that all the results of §§2-3-4 transfer easily to the general case where Δ is not necessarily a basis.

5.3. From the linearly independent case to the positively independent case.

In the case where Δ is not a basis, there is another way to study the limit points of roots (without using general transverse hyperplanes), by pulling back all the structure to a space where Δ is a basis, using the canonical geometric W -module defined in §1.4. let us first give the following definition (motivated by Example 5.1).

Definition 5.6. Let (W, S) be a Coxeter group and $A = (a_{s,t})$ be a matrix that satisfies Equation (1.2). A pair (V, B) is a *geometric W -module associated to A* if

- (i) V is a finite dimensional real vector space containing a subset $\Delta = \{\alpha_s \mid s \in S\}$ that is positively independent;
- (ii) B is a bilinear form such that $B(\alpha_s, \alpha_t) = a_{s,t}$ for all $s, t \in S$.

In this case, since Δ satisfies the requirement of Definition 1.2, (V, B) is a geometric W -module (as in Definition 1.2); the associated based root system is called the *based root system associated to (V, B)* .

Note that any geometric W -module (V, B) of Definition 1.2 is a geometric W -module associated to a certain matrix A determined by Conditions (ii) and (iii) of Definition 1.2.

Obviously, for each matrix A satisfying Equation (1.2), the simplest geometric W -module associated to A is the “canonical” geometric W -module (V_A, B_A) (defined in §1.4), where we declare Δ as a basis. However, as Example 5.1 shows, the set of geometric modules (V, B) —up to isomorphism—associated to A may contain more than the class of (V_A, B_A) . Still, the canonical geometric W -module associated to A carries all the combinatorics of the based root system of A , as described in the following proposition.⁷

Proposition 5.7. *Let (W, S) be a Coxeter system and A be an associated matrix satisfying Equation (1.2). Let (V_A, B_A) be the canonical geometric W -module, and (V, B) be a geometric W -module associated to A , with based root system (Φ, Δ) . Then there is a morphism of W -modules $\varphi_A : V_A \rightarrow V$, preserving B , that restricts to a bijection from Φ_A to Φ , such that $\varphi_A(\Delta_A) = \Delta$ and $\varphi_A(\Phi_A^+) = \Phi^+$.*

Proof. Write $\Delta = \{\alpha_s \mid s \in S\}$ and $\Delta_A = \{\gamma_s \mid s \in S\}$. So

$$B(\alpha_s, \alpha_t) = a_{s,t} = B_A(\gamma_s, \gamma_t).$$

Now let

$$\varphi_A : V_A \rightarrow V$$

be the linear map defined by mapping any element γ_s of the basis Δ_A of V_A to α_s . We have $B(\varphi_A(\gamma_s), \varphi_A(\gamma_t)) = B(\alpha_s, \alpha_t) = B_A(\gamma_s, \gamma_t)$; so that it is easy to check that φ_A is a morphism of W -modules, preserving B .

The fact that φ_A restricts to a bijection from Φ_A to Φ is nontrivial, see [Fu12, Proposition 2.1]. Finally, as by construction we have $\varphi_A(\Delta_A) = \Delta$, we get directly that φ_A also maps Φ_A^+ to Φ^+ . \square

⁷This proposition was suggested by M. Dyer (personal communication, September 2011), see also [Fu12, Proposition 2.1].

Proposition 5.7 tends to imply that in the previous sections we did not lose much generality by studying only root systems where Δ is a basis.

Indeed, let (V, B) be a geometric W -module with based root system (Φ, Δ) , where Δ is not necessarily linearly independent. Then we know that (V, B) is a geometric W -module associated to a matrix A ; but we can also construct, from §1.4, the canonical geometric W -module (V_A, B_A) and the based root system (Φ_A, Δ_A) for which Δ_A is a basis. The proposition above tells us that the map φ_A is a morphism of W -modules carrying bijectively the combinatorics of the root system Φ_A to that of the root system Φ . Moreover, since φ_A is a linear map, it is continuous, and the topological properties we established in V_A are carried to V through φ_A .

In V_A , the simple system Δ_A is a basis, so we can apply our usual construction to (Φ_A, Δ_A) , building the hyperplane $(V_A)_1$, the normalized roots $\widehat{\Phi}_A$ and the limit roots $E(\Phi_A)$. Now choose a hyperplane H in V , transverse to Φ^+ . Thanks to Proposition 5.7, we see that

$$\text{conv}(\pi_H(\Delta)) = \pi_H \circ \varphi_A(\text{conv}(\Delta_A)) \quad \text{and} \quad \pi_H(\Phi) = \pi_H \circ \varphi_A(\widehat{\Phi}_A).$$

Since φ_A and π_H are continuous, $\pi_H \circ \varphi_A$ is continuous. Therefore we obtain:

$$E(\Phi, H) = \pi_H \circ \varphi_A(E(\Phi_A)).$$

Thus, the sets of normalized roots and limit roots of a based root system (Φ, Δ) attached to a matrix A are the images of the same objects for the based root system (Φ_A, Δ_A) . Overall, this means that all the properties we proved in the previous sections in the case where Δ is a basis are simply transferred to the general case.

However it is not clear whether the map $\pi_H \circ \varphi_A$ induces a bijection between $E(\Phi_A)$ and $E(\Phi, H)$: we do not know if in general $E(\Phi, H)$ is a faithful image of $E(\Phi_A)$; but $E(\Phi, H)$ is always a projection of $E(\Phi_A)$ by construction.

5.4. On limit roots for reflection subgroups.

We now address how to define the limit roots for a reflection subgroup. Let (Φ, Δ) be a based root system of (V, B) with associated Coxeter system (W, S) . Fix H a hyperplane that is transverse to Φ^+ , according to Definition 5.2 (e.g., take $H = V_1$ if Δ is a basis). Let (Φ', Δ') be a based root subsystem of (Φ, Δ) . Then:

- The hyperplane H is also transverse to Φ'^+ .
- So, we define intrinsically the set $E(\Phi', H)$, constituted with the limit points of $\pi_H(\Phi')$.
- Obviously $E(\Phi', H)$ is a subset of $E(\Phi, H)$.

So, as wanted in §5.1, we get an intrinsic definition of the limit roots for a root subsystem, which is compatible with the inclusion inside the whole set of limit roots. Moreover, thanks to §5.3, if we set

- A the matrix associated to the root subsystem (Φ', Δ') ,
- (V_A, B_A) the canonical geometric W' -module for A , with based root system (Φ_A, Δ_A) ,
- $\varphi_A : V_A \rightarrow V$ the associated map defined in Proposition 5.7,
- $E(\Phi_A)$ the set of limit roots for (Φ_A, Δ_A) ,

then $E(\Phi', H)$ is the renormalization of the image under φ_A of $E(\Phi_A)$

$$E(\Phi', H) = \pi_H(\varphi_A(E(\Phi_A))).$$

In the following, to lighten the notations, the cutting hyperplane H is implicit.

The set $E(\Phi')$ is always included in $\text{conv}(\Delta')$ (and in $E(\Phi)$). One can wonder whether the inclusion $E(\Phi') \subseteq E(\Phi) \cap \text{conv}(\Delta')$ is an equality. Consider the most simple case, where the reflection subgroup W' is in fact a standard parabolic subgroup. Let I be a subset of S , $\Delta_I = \{\alpha_s \mid s \in I\}$ and Φ_I its orbit under $W_I = \langle I \rangle$ (i.e., a parabolic root subsystem). Denote by F_I the convex hull of Δ_I , and $E(\Phi_I)$ the set of accumulation points of $\widehat{\Phi_I}$, which lives inside F_I . In this case, the question is: if a limit root lies in the face F_I of the simplex $\text{conv}(\Delta)$, is this a limit point of normalized roots of W_I ? Surprisingly enough, the answer can be negative, as shown by the following counterexample.

Example 5.8. Take the rank 5 root system Φ with $\Delta = \{\alpha, \beta, \gamma, \delta, \varepsilon\}$ and the labels $m_{\alpha, \beta} = m_{\delta, \varepsilon} = \infty$, $m_{\beta, \gamma} = m_{\gamma, \delta} = 3$, and the others equal to 2. Take $\Delta_I = \Delta \setminus \{\gamma\}$, so that W_I is the direct product of two infinite dihedral groups. Then we have $E(\Phi_I) = \{\frac{\alpha+\beta}{2}, \frac{\delta+\varepsilon}{2}\}$ (see §2.3). But if we consider $\rho_n = (s_\alpha s_\beta s_\varepsilon s_\delta)^n(\gamma)$, it is easy to check that $\widehat{\rho_n}$ tends to $\frac{\alpha+\beta+\delta+\varepsilon}{4}$, that lies in $E(\Phi) \cap F_I$ but not in $E(\Phi_I)$.

In a subsequent paper [DHR13], we show that this property holds nevertheless for the set E_2 studied in §4, i.e., $E_2(\Phi_I) = E_2(\Phi) \cap F_I$. We also define other natural smaller *dense* subsets of E for which this property of parabolic restriction works.

Acknowledgements. We would like to thank Matthew Dyer for having communicated to us his preprint [Dye12], for many fruitful discussions at LaCIM in September 2011, for the idea of the counterexample 5.8 and the suggestion of Proposition 5.7. We are also grateful to the referees of an extended abstract of this article (submitted to the conference FPSAC 2012) for helpful comments. Finally, we gratefully thank the anonymous referee for the careful reading of our manuscript, for the references to [Sau91] and [Kra09, Lemma 6.1.1] and for the valuable comments and suggestions, which contributed to improve the quality of this article.

REFERENCES

- [BB05] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [BD10] Cédric Bonnafé and Matthew Dyer. Semidirect product decomposition of Coxeter groups. *Comm. Algebra*, 38(4):1549–1574, 2010. <http://dx.doi.org/10.1080/00927870902980354>.
- [BH93] Brigitte Brink and Robert B. Howlett. A finiteness property and an automatic structure for Coxeter groups. *Math. Ann.*, 296(1):179–190, 1993.
- [Bou68] Nicolas Bourbaki. *Groupes et algèbres de Lie*. Chapitre 4-6. Paris: Hermann, 1968.
- [Deo89] Vinay V. Deodhar. A note on subgroups generated by reflections in Coxeter groups. *Arch. Math. (Basel)*, 53(6):543–546, 1989. <http://dx.doi.org/10.1007/BF01199813>.
- [Dye90] Matthew Dyer. Reflection subgroups of Coxeter systems. *J. Algebra*, 135(1):57–73, 1990. [http://dx.doi.org/10.1016/0021-8693\(90\)90149-I](http://dx.doi.org/10.1016/0021-8693(90)90149-I).
- [Dye10] Matthew Dyer. On rigidity of abstract root systems of Coxeter systems. Preprint arXiv:1011.2270 <http://arxiv.org/abs/1011.2270>, November 2010.
- [Dye11] Matthew Dyer. On the weak order of Coxeter groups. Preprint arXiv:1108.5557 <http://arxiv.org/abs/1108.5557>, August 2011.
- [Dye12] Matthew Dyer. Imaginary cone and reflection subgroups of Coxeter groups. Preprint arXiv:1210.5206 <http://arxiv.org/abs/1210.5206>, October 2012.
- [DHR13] Matthew Dyer, Christophe Hohlweg, and Vivien Ripoll. Imaginary cones and limit roots of infinite Coxeter groups. Preprint arXiv:1303.6710 <http://arxiv.org/abs/1303.6710>, March 2013.
- [Fu12] Xiang Fu. The dominance hierarchy in root systems of Coxeter groups. *J. Algebra*, 366:187–204, 2012.

- [Fu13] Xiang Fu. Coxeter groups, imaginary cones and dominance. *Pacific J. Math.*, 262(2):339–363, 2013.
- [How96] Robert B. Howlett. Introduction to Coxeter groups. Lectures at A.N.U., available at <http://www.maths.usyd.edu.au/res/Algebra/How/1997-6.html>, February 1996.
- [Hum90] James E. Humphreys. *Reflection groups and Coxeter groups*, volume 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.
- [Kac90] Victor G. Kac. *Infinite-dimensional Lie algebras*. Cambridge University Press, Cambridge, third edition, 1990. <http://dx.doi.org/10.1017/CB09780511626234>.
- [Kan01] Richard Kane. *Reflection groups and invariant theory*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 5. Springer-Verlag, New York, 2001.
- [Kra09] Daan Krammer. The conjugacy problem for Coxeter groups. *Groups Geom. Dyn.*, 3(1):71–171, 2009. <http://dx.doi.org/10.4171/GGD/52>.
- [LN04] Ottmar Loos and Erhard Neher. Locally finite root systems. *Mem. Amer. Math. Soc.*, 171(811):x+214, 2004.
- [MP89] Robert V. Moody and Arturo Pianzola. On infinite root systems. *Trans. Amer. Math. Soc.*, 315(2):661–696, 1989. <http://dx.doi.org/10.2307/2001300>.
- [Sau91] Clive Saunders. The reflection representations of some Chevalley groups. *J. Algebra*, 137(1):145–165, 1991.
- [S+13] William A. Stein et al. *Sage Mathematics Software (Version 5.9)*. The Sage Development Team, 2013. <http://www.sagemath.org>.
- [Vin71] Ernest. B. Vinberg. Discrete linear groups that are generated by reflections. *Izv. Akad. Nauk SSSR Ser. Mat.*, 35:1072–1112, 1971. translation by P. Flor, IOP Science.

(Christophe Hohlweg) UNIVERSITÉ DU QUÉBEC À MONTRÉAL, LACIM ET DÉPARTEMENT DE MATHÉMATIQUES, CP 8888 SUCC. CENTRE-VILLE, MONTRÉAL, QUÉBEC, H3C 3P8, CANADA
E-mail address: hohlweg.christophe@uqam.ca
URL: <http://hohlweg.math.uqam.ca>

(Jean-Philippe Labbé) FREIE UNIVERSITÄT BERLIN, INSTITUT FÜR MATHEMATIK, ARNIMALLEE 2, 14195, BERLIN, DEUTSCHLAND
E-mail address: labbe@math.fu-berlin.de
URL: <http://page.mi.fu-berlin.de/labbe>

(Vivien Ripoll) UNIVERSITÉ DU QUÉBEC À MONTRÉAL, LACIM ET DÉPARTEMENT DE MATHÉMATIQUES, CP 8888 SUCC. CENTRE-VILLE, MONTRÉAL, QUÉBEC, H3C 3P8, CANADA
E-mail address: vivien.ripoll@lacim.ca
URL: <http://www.math.ens.fr/~vripoll>