

CHARACTERISTIC SUBMANIFOLD THEORY AND TOROIDAL DEHN FILLING

STEVEN BOYER, CAMERON MCA. GORDON, AND XINGRU ZHANG

March 27, 2012

ABSTRACT. The exceptional Dehn filling conjecture of the second author concerning the relationship between exceptional slopes α, β on the boundary of a hyperbolic knot manifold M has been verified in all cases other than small Seifert filling slopes. In this paper we verify it when α is a small Seifert filling slope and β is a toroidal filling slope in the generic case where M admits no punctured-torus fibre or semi-fibre, and there is no incompressible torus in $M(\beta)$ which intersects ∂M in one or two components. Under these hypotheses we show that $\Delta(\alpha, \beta) \leq 5$. Our proof is based on an analysis of the relationship between the topology of M , the combinatorics of the intersection graph of an immersed disk or torus in $M(\alpha)$, and the two sequences of characteristic subsurfaces associated to an essential punctured torus properly embedded in M .

1. INTRODUCTION

This is the first of four papers concerned with the relationship between Seifert filling slopes and toroidal filling slopes on the boundary of a hyperbolic knot manifold M . Such results are part of the exceptional surgery problem, which we describe now.

A *hyperbolic knot manifold* M is a compact, connected, orientable, hyperbolic 3-manifold whose boundary is a torus. A *slope* on ∂M is a ∂M -isotopy class of essential simple closed curves. Slopes can be visualized by identifying them with \pm -classes of primitive elements of $H_1(\partial M)$ in the *surgery plane* $H_1(\partial M; \mathbb{R})$. The *distance* $\Delta(\alpha_1, \alpha_2)$ between slopes α_1, α_2 is the absolute value of the algebraic intersection number of their associated classes in $H_1(\partial M)$. Given a set of slopes \mathcal{S} , set $\Delta(\mathcal{S}) = \sup\{\Delta(\alpha, \beta) : \alpha, \beta \in \mathcal{S}\}$.

To each slope α on ∂M we associate the α -*Dehn filling* $M(\alpha) = (S^1 \times D^2) \cup_f M$ of M where $f : \partial(S^1 \times D^2) \rightarrow \partial M$ is any homeomorphism such that $f(\{*\} \times \partial D^2)$ represents α .

Set $\mathcal{E}(M) = \{\alpha \mid M(\alpha) \text{ is not hyperbolic}\}$ and call the elements of $\mathcal{E}(M)$ *exceptional slopes*. It follows from Thurston's hyperbolic Dehn surgery theorem that $\mathcal{E}(M)$ is finite, while Perelman's solution of the geometrisation conjecture implies that

$$\mathcal{E}(M) = \{\alpha \mid M(\alpha) \text{ is either reducible, toroidal, or small Seifert}\}$$

Steven Boyer was partially supported by NSERC grant RGPIN 9446-2008.

Cameron Gordon was partially supported by NSF grant DMS-0906276.

2000 MATHEMATICS SUBJECT CLASSIFICATION. PRIMARY 57M25, 57M50, 57M99

Here, a manifold is *small Seifert* if it admits a Seifert structure with base orbifold of the form $S^2(a, b, c)$, where $a, b, c \geq 1$.

Much work has been devoted to understanding the structure of $\mathcal{E}(M)$ and describing the topology of M when $|\mathcal{E}(M)| \geq 2$. For instance, sharp upper bounds are known for the distance between two reducible filling slopes [GL], between two toroidal filling slopes [Go], [GW], and between a reducible filling slope and a toroidal filling slope [Oh], [Wu1]. More recently, strong upper bounds were obtained for the distance between a reducible filling slope and a small Seifert filling slope [BCSZ2], [BGZ2]. In this paper, and its sequel, we examine the distance between toroidal filling slopes and small Seifert filling slopes.

Let W be the exterior of the right-handed Whitehead link and T one of its boundary components. Consider the following hyperbolic knot exteriors obtained by the indicated Dehn filling of W along T : $M_1 = W(T; -1)$, $M_2 = W(T; -2)$, $M_3 = W(T; 5)$, $M_4 = W(T; \frac{5}{2})$. One of the key conjectures concerning $\mathcal{E}(M)$ is the following:

Conjecture 1.1 (C.McA. Gordon). *For any hyperbolic knot manifold M , we have $\#\mathcal{E}(M) \leq 10$ and $\Delta(\mathcal{E}(M)) \leq 8$. Moreover, if $M \neq M_1, M_2, M_3, M_4$, then $\#\mathcal{E}(M) \leq 7$ and $\Delta(\mathcal{E}(M)) \leq 5$.*

It is shown in [BGZ1] that the conjecture holds if the first Betti number of M is at least 2. (By duality, it is at least 1.) Lackenby and Meyerhoff have proven that the first statement of the conjecture holds in general [LM]. See §2 of their paper for a historical survey of results concerning upper bounds for $\#\mathcal{E}(M)$ and $\Delta(\mathcal{E}(M))$. Agol has shown that there are only finitely many hyperbolic knot manifolds M with $\Delta(\mathcal{E}(M)) > 5$ [Ag], though no practical fashion to determine this finite set is known.

It follows from [GL], [Oh], [Wu1], [Go] and [GW] that Conjecture 1.1 holds if $\mathcal{E}(M)$ is replaced by $\mathcal{E}(M) \setminus \{\alpha \mid M(\alpha) \text{ is small Seifert}\}$. It remains, therefore, to consider pairs α, β such that $M(\alpha)$ is small Seifert and $M(\beta)$ is either reducible, toroidal or small Seifert. The first case is treated in [BCSZ2], where it is shown that, generically, $\Delta(\alpha, \beta) \leq 4$. (See below for a more precise statement.) In the present paper we are interested in the second case. (We remark that if $M(\alpha)$ is toroidal Seifert fibred and $M(\beta)$ is toroidal then $\Delta(\alpha, \beta) \leq 4$ [BGZ3].) Here, Conjecture 1.1 implies

Conjecture 1.2. *Suppose that M is a hyperbolic knot manifold M and α, β are slopes on ∂M such that $M(\alpha)$ is small Seifert and $M(\beta)$ toroidal. If $\Delta(\alpha, \beta) > 5$, then M is the figure eight knot exterior.*

Understanding the relationship between Dehn fillings which yield small Seifert manifolds and other slopes in $\mathcal{E}(M)$ has proven difficult. The techniques used to obtain sharp distance bounds in other cases either provide relatively weak bounds here or do not apply at all. For instance, the graph intersection method (see e.g. [CGLS], [GL]) cannot be used as typically, small Seifert manifolds do not admit closed essential surfaces. On the other hand, they usually do admit essential immersions of tori, a fact which can be exploited. Suppose that α and β are slopes on ∂M such that $M(\alpha)$ is small Seifert and M contains an essential surface F of slope β . It

was shown in [BCSZ1] how to construct an immersion $h : Y \rightarrow M(\alpha)$ where Y is a disk or torus, a labeled “intersection” graph $\Gamma_F = h^{-1}(F) \subset Y$, and, for each sign $\epsilon = \pm$, a sequence of characteristic subsurfaces

$$F = \dot{\Phi}_0^\epsilon \supset \dot{\Phi}_2^\epsilon \supset \dot{\Phi}_3^\epsilon \supset \dots \supset \dot{\Phi}_n^\epsilon \supset \dots$$

The relationship between the combinatorics of Γ_F , the two sequences of characteristic subsurfaces, and the topology of M was exploited in [BCSZ2] to show that if $M(\alpha)$ is small Seifert, $M(\beta)$ is reducible, and the (planar) surface F is neither a fibre nor semi-fibre in M , then $\Delta(\alpha, \beta) \leq 4$. (See also [CL], [Li] where a related method is used to study the existence of immersed essential surfaces in Dehn fillings of knot manifolds.) The main contributions of this paper are the further refinement of this technique and its application in the investigation of Conjecture 1.2.

When M is the figure eight knot exterior there are (up to orientation-reversing homeomorphism of M) two pairs (α, β) with $\Delta(\alpha, \beta) > 5$ such that $M(\alpha)$ is small Seifert and $M(\beta)$ is toroidal, namely $(-3, 4)$ and $(-2, 4)$. The toroidal manifold $M(4)$ contains a separating incompressible torus which intersects ∂M in two components. Moreover the corresponding punctured torus is not a fibre or semi-fibre in M . We show that if a hyperbolic knot manifold M has a small Seifert filling $M(\alpha)$ and a toroidal filling $M(\beta)$ then $\Delta(\alpha, \beta) \leq 5$ in the generic case when M admits no punctured-torus fibre or semi-fibre, and there is no incompressible torus in $M(\beta)$ which intersects ∂M in one or two components. Our precise result is stated in §2 where we also detail our underlying assumptions and provide an outline of the paper. We will examine the non-generic cases of Conjecture 1.2 in the forthcoming manuscripts [BGZ3], [BGZ4].

We are indebted to Marc Culler and Peter Shalen for their role in the development of the ideas in this paper.

2. BASIC ASSUMPTIONS AND STATEMENT OF MAIN RESULT

Throughout the paper we work under the following assumptions.

Assumption 2.1. *$M(\alpha)$ is a small Seifert manifold with base orbifold $S^2(a, b, c)$ where $a, b, c \geq 1$ and $M(\beta)$ is toroidal.*

Assumption 2.2. *Among all embedded essential tori in $M(\beta)$, \widehat{F} is one whose intersection with ∂M has the least number of components.*

Then $F = \widehat{F} \cap M$ is a properly embedded essential punctured torus in M with boundary slope β . Set $m = |\partial F| \geq 1$.

Assumption 2.3. *If there is an essential separating torus in $M(\beta)$ satisfying Assumption 2.2, \widehat{F} has been chosen to be separating.*

Assumption 2.4. *If there is an essential torus in $M(\beta)$ satisfying Assumption 2.2 which bounds a twisted I -bundle over the Klein bottle in $M(\beta)$, \widehat{F} has been chosen to bound such an I -bundle.*

Note that it is possible that there are essential tori $\widehat{F}_1, \widehat{F}_2$ in $M(\beta)$ which bound twisted I -bundles over the Klein bottle in $M(\beta)$, such that $\widehat{F}_1 \cap M$ is the frontier of a twisted I -bundle in M but $\widehat{F}_2 \cap M$ is not.

Assumption 2.5. *If there is an essential torus \widehat{F} in $M(\beta)$ satisfying Assumption 2.2 such that there is a twisted I -bundle in M with frontier $F = \widehat{F} \cap M$, \widehat{F} has been chosen so that F is the frontier of a twisted I -bundle.*

Let S be the surface in M which is F when F is separating and is a union of two parallel copies F_1, F_2 of F when F is non-separating. Then S splits M into two components X^+ and X^- .

Let \widehat{S} be a closed surface in $M(\beta)$ obtained by attaching disjoint meridian disks of the β -filling solid torus to S . Then \widehat{S} splits $M(\beta)$ into two compact submanifolds \widehat{X}^+ containing X^+ and \widehat{X}^- containing X^- , each having incompressible boundary \widehat{S} .

We call F a *fibre* in M if it is a fibre of a surface bundle map $M \rightarrow S^1$. Equivalently, the exterior of F in M is homeomorphic to $F \times I$. We call F a *semi-fibre* in M if it separates and splits M into two twisted I -bundles.

Assumption 2.6. *Assume that F , chosen as above, is neither a fibre nor a semi-fibre in M . In particular, assume that X^+ is not an I -bundle over a surface.*

Here is our main theorem.

Theorem 2.7. *Suppose that M is a hyperbolic knot manifold and α, β slopes on ∂M such that $M(\alpha)$ is a small Seifert manifold and $M(\beta)$ is toroidal. Let F be an essential genus 1 surface of slope β which is properly embedded in M and which satisfies the assumptions listed above. If $m \geq 3$, then $\Delta(\alpha, \beta) \leq 5$.*

When the first Betti number of M is at least 2 or one of $M(\alpha)$ and $M(\beta)$ is reducible, Theorem 2.7 holds by [Ga], [BGZ1], [BCSZ2], [Oh], [Wu1]. Thus we make the following assumption.

Assumption 2.8. *The first Betti number of M is 1 and both $M(\alpha)$ and $M(\beta)$ are irreducible.*

The paper is organised as follows. Section 3 contains background information on characteristic submanifolds associated to the pair (X^ϵ, S) ($\epsilon = \pm$). Sections 4 and 5 are devoted to exploring the restrictions forced on essential annuli in (X^ϵ, S) by our assumptions on F . These results will be applied in §7 to the study the structure of the characteristic submanifolds of (X^ϵ, S) and the topology of \widehat{X}^ϵ . An analysis of the existence and numbers of certain characteristic subsurfaces of S is made in §6, §8, and §9. The relation between the number of such surfaces and the length of essential homotopies in (M, S) is determined in §10. Section 11 introduces the intersection graphs associated with certain immersions in $M(\alpha)$ and relates their structure to lengths of essential homotopies, leading to bounds on $\Delta(\alpha, \beta)$. Conditions which guarantee the existence of faces of the graph with few edges are investigated in §12, while the relations in the fundamental groups of X^+ and X^- associated to these faces are considered in §15. Theorem 2.7 is proved when F is non-separating in §13 and in the presence of “tight” characteristic

subsurfaces in §14 and §16. The implications of certain combinatorial configurations in the intersection graph are examined in §17. The proof of Theorem 2.7 in the absence of tight components is achieved in the last two sections.

CONTENTS

1. Introduction	1
2. Basic assumptions and statement of main result	3
3. Characteristic submanifolds of (X^ϵ, S)	6
3.1. General subsurfaces of S	6
3.2. Characteristic subsurfaces of S	6
4. Essential embedded annuli in (X^ϵ, S)	8
5. Pairs of embedded essential annuli in (M, S)	13
6. The dependence of the number of tight components of $\check{\Phi}_j^\epsilon$ on j	15
7. The structure of $\dot{\Phi}_1^\epsilon$ and the topology of X^ϵ	17
7.1. S is connected	18
7.2. S is not connected	21
8. \widehat{S} -essential annuli in $\dot{\Phi}_j^\epsilon$	23
9. The existence of tight components in $\check{\Phi}_j^\epsilon$ for small values of j	25
10. Lengths of essential homotopies	28
11. The intersection graph of an immersed disk or torus	29
12. Counting faces in $\overline{\Gamma}_S$	34
13. Proof of Theorem 2.7 when F is non-separating	36
14. Proof of Theorem 2.7 when F is separating and $t_1^+ + t_1^- \geq 4$	37
15. The relation associated to a face of Γ_S	37
16. Proof of Theorem 2.7 when F is separating and $t_1^+ + t_1^- = 2$	39
17. Extended S -cycles in Γ_S	42
18. Proof of Theorem 2.7 when X^- is not an I -bundle and $t_1^+ = t_1^- = 0$	44
18.1. Background results	44
18.2. Proof	46
19. Proof of Theorem 2.7 when X^- is a twisted I -bundle and $t_1^+ = 0$	47
19.1. Background results	47
19.2. Proof when $\dot{\Phi}_3^+$ is a union of tight components	49

19.3. Proof when $\dot{\Phi}_3^+$ is not a union of tight components	68
References	74

3. CHARACTERISTIC SUBMANIFOLDS OF (X^ϵ, S)

3.1. General subsurfaces of S . A surface is called *large* if each of its components has negative Euler characteristic.

A connected subsurface S_0 of S is called *neat* if it is either a collar on a boundary component of S or it is large and each boundary component of S that can be isotoped into S_0 is contained in S_0 . A subsurface of S is *neat* if each of its components has this property.

For each boundary component b of S , let \hat{b} denote a meridian disk which it bounds in the β -filling solid torus of $M(\beta)$. The *completion* of a neat subsurface S_0 of S is the surface $\hat{S}_0 \subseteq M(\beta)$ obtained by attaching the disks \hat{b} to S_0 for each boundary component b of S contained in S_0 .

A simple closed curve $c \subseteq S$ is called *outer* if it is parallel in S to a component of ∂S . Otherwise it is called *inner*.

A boundary component of a subsurface is called an *inner boundary component* if it is an inner curve, and an *outer boundary component* otherwise.

A neat subsurface S_0 of S is called *tight* if \hat{S}_0 is a disk. Equivalently, S_0 is a connected, planar, neat subsurface of S with at most one inner boundary component.

A simple closed curve $c \subseteq S$ which is essential in \hat{S} will be called *\hat{S} -essential*.

We call a subsurface S_0 of S an *\hat{S} -essential annulus* if \hat{S}_0 is an essential annulus in \hat{S} .

Two essential annuli in \hat{S} are called *parallel* if their core circles are parallel in \hat{S} .

3.2. Characteristic subsurfaces of S . For the rest of the paper we use ϵ to denote either of the signs $\{\pm\}$.

A map f of a path-connected space Y to S is called *large* if $f_{\#}(\pi_1(Y))$ contains a non-abelian free group.

A map of pairs $f : (Y, Z) \rightarrow (X^\epsilon, S)$ is called *essential* if it is not homotopic, as a map of pairs, to a map $f' : (Y, Z) \rightarrow (X^\epsilon, S)$ where $f'(Y) \subseteq S$.

An *essential annulus* in (X^ϵ, S) is the image of an essential proper embedding $(S^1 \times I, S^1 \times \partial I) \rightarrow (X^\epsilon, S)$.

An *essential homotopy of length n* in (M, S) of $f : Y \rightarrow S$ which starts on the ϵ -side of S is a homotopy

$$H : (Y \times [a, a+n], Y \times \{a, a+1, \dots, a+n\}) \rightarrow (M, S)$$

such that

- (1) $H(y, a) = f(y)$ for each $y \in Y$;
- (2) $H^{-1}(S) = Y \times \{a, a + 1, \dots, a + n\}$;
- (3) for each $i \in \{1, 2, \dots, n\}$, $H|_{Y \times [a + i - 1, a + i]}$ is an essential map in $(X^{(-1)^{i-1}\epsilon}, S)$.

Let $(\Sigma^\epsilon, \Phi^\epsilon) \subseteq (X^\epsilon, S)$ be the characteristic I -bundle pair of (X^ϵ, S) [JS], [Jo]. We shall use τ_ϵ to denote the free involution on Φ^ϵ which interchanges the endpoints of the I -fibres of Σ^ϵ .

The union of the components P of Σ^ϵ for which $P \cap S$ is large is denoted by Σ_1^ϵ . Set

$$\Phi_0^\epsilon = S$$

and

$$\Phi_1^\epsilon = \Sigma_1^\epsilon \cap S$$

More generally, for $j \geq 0$ we define $\Phi_j^\epsilon \subseteq S$ to be the j -th characteristic subsurface with respect to the pair (M, S) as defined in §5 of [BCSZ1]. We shall assume throughout the paper that Φ_j^ϵ is neatly embedded in S . It is characterised up to ambient isotopy by the following property:

- (*) $\left\{ \begin{array}{l} \text{a large function } f_0 : Y \rightarrow S \text{ admits an essential homotopy of length } j \text{ which starts} \\ \text{on the } \epsilon\text{-side of } S \text{ if and only if it is homotopic in } S \text{ to a map with image in } \Phi_j^\epsilon \end{array} \right.$

See [BCSZ1, Proposition 5.2.8].

A compact connected 3-dimensional submanifold P of X^ϵ is called *neat* if

- (1) $\partial P \cap S$ is a neat subsurface of S ;
- (2) each component of $\overline{\partial P \setminus S}$ is an essential annulus in (X^ϵ, S) ;
- (3) some component of $\partial M \cap X^\epsilon$ is isotopic in (X^ϵ, S) into P , then it is contained in P .

A compact 3-dimensional submanifold P of X^ϵ is called *neat* if each of its components has this property.

Given a neat submanifold P of X^ϵ , we use \widehat{P} to denote the submanifold of \widehat{X}^ϵ obtained by attaching to P those components H of $\widehat{X}^\epsilon \setminus \text{int}(M)$ for which $H \cap \partial M \subseteq P$.

For convenience we describe some properties of the characteristic I -bundle pair $(\Sigma^\epsilon, \Phi^\epsilon)$ which hold under the assumptions on F listed in §2, even though their justification will only be addressed in §4.

It follows from Proposition 4.9 that if c and $\tau_\epsilon(c)$ are two outer boundary components of Φ_1^ϵ , then Φ_1^ϵ can be isotoped so that the annulus component in $\partial M \cap X^\epsilon$ bounded by c and $\tau_\epsilon(c)$ is contained in Σ_1^ϵ . We will therefore assume from §5 on that Σ_1^ϵ is neatly embedded in S .

Let $\dot{\Phi}_j^\epsilon$ denote the union of the components of Φ_j^ϵ which contain some outer boundary components. We will see in Proposition 4.4 that τ_ϵ preserves the set of outer, respectively inner, essential simple closed curves in Φ_1^ϵ . Hence, it restricts to a free involution on $\dot{\Phi}_1^\epsilon$, which we continue to denote τ_ϵ . Let $\dot{\Sigma}_1^\epsilon$ denote the corresponding I -bundle.

Let $\check{\Phi}_j^\epsilon$ be the neat subsurface in S obtained from the union of $\dot{\Phi}_j^\epsilon$ and a closed collar neighbourhood of $\partial S \setminus \partial \dot{\Phi}_j^\epsilon$ in $S \setminus \dot{\Phi}_j^\epsilon$. It follows from the previous paragraph that there is an I -bundle pair $(\check{\Sigma}_1^\epsilon, \check{\Phi}_1^\epsilon)$ properly embedded in (X^ϵ, S) where $\check{\Sigma}_1^\epsilon$ is the union of $\dot{\Sigma}_1^\epsilon$ and closed collar neighbourhoods of the annular components of $\partial M \cap X^\epsilon$ cobounded by components of $\partial S \setminus \partial \dot{\Phi}_1^\epsilon$. Thus $\tau_\epsilon : \dot{\Phi}_1^\epsilon \rightarrow \dot{\Phi}_1^\epsilon$ extends to an involution $\tau_\epsilon : \check{\Phi}_1^\epsilon \rightarrow \check{\Phi}_1^\epsilon$.

A properly embedded annulus in (X^ϵ, S) is called *vertical* if it is a union of I -fibres of $\check{\Sigma}_1^\epsilon$. A subsurface of $\check{\Phi}_1^\epsilon$ is called *horizontal*.

It follows from the defining property $(*)$ of the surfaces $\dot{\Phi}_j^\epsilon$ that if (X^-, S) is an I -bundle pair, then for each $j \geq 0$,

$$\begin{aligned} (\check{\Phi}_{2j}^-, \dot{\Phi}_{2j}^-) &= (\check{\Phi}_{2j+1}^-, \dot{\Phi}_{2j+1}^-) \\ (\check{\Phi}_{2j+1}^+, \dot{\Phi}_{2j+1}^+) &= (\check{\Phi}_{2j+2}^+, \dot{\Phi}_{2j+2}^+) \\ (\check{\Phi}_{2j+2}^-, \dot{\Phi}_{2j+2}^-) &= (\tau_-(\check{\Phi}_{2j+1}^+), \tau_-(\dot{\Phi}_{2j+1}^+)) \end{aligned}$$

Recall from [BCSZ1, Proposition 5.3.1] that for each ϵ and $j \geq 0$ there is a homeomorphism

$$h_j^\epsilon : (\check{\Phi}_j^\epsilon, \dot{\Phi}_j^\epsilon) \rightarrow (\check{\Phi}_j^{(-1)^{j+1}\epsilon}, \dot{\Phi}_j^{(-1)^{j+1}\epsilon})$$

obtained by concatenating alternately restrictions of τ_+ and τ_- . These homeomorphisms satisfy some useful properties:

$$\begin{aligned} h_1^\epsilon &= \tau_\epsilon \\ h_{2j}^\epsilon : (\check{\Phi}_{2j}^\epsilon, \dot{\Phi}_{2j}^\epsilon) &\xrightarrow{\cong} (\check{\Phi}_{2j}^{-\epsilon}, \dot{\Phi}_{2j}^{-\epsilon}) \\ h_{2j+1}^\epsilon : (\check{\Phi}_{2j+1}^\epsilon, \dot{\Phi}_{2j+1}^\epsilon) &\xrightarrow{\cong} (\check{\Phi}_{2j+1}^\epsilon, \dot{\Phi}_{2j+1}^\epsilon) \text{ is a free involution} \end{aligned}$$

Finally, consider two large subsurfaces S_0, S_1 of S . Their *large essential intersection* is a large, possibly empty, subsurface $S_0 \wedge S_1$ of S characterised up to isotopy in S by the property:

$$(**) \left\{ \begin{array}{l} \text{a large function } f : Y \rightarrow S \text{ is homotopic into both} \\ S_0 \text{ and } S_1 \text{ if and only if it is homotopic into } S_0 \wedge S_1 \end{array} \right.$$

See [BCSZ1, Proposition 4.2]. It follows from the defining property $(*)$ of the surfaces $\dot{\Phi}_j^\epsilon$ that

$$(3.2.1) \quad h_j^\epsilon(\dot{\Phi}_{j+k}^\epsilon) = \dot{\Phi}_j^{(-1)^{j+1}\epsilon} \wedge \dot{\Phi}_k^{(-1)^j\epsilon}$$

4. ESSENTIAL EMBEDDED ANNULI IN (X^ϵ, S)

The next two sections are devoted to exploring the restrictions forced on essential annuli in (X^ϵ, S) by our assumptions on F . These results will be applied in §7 to the study of the structure of $\dot{\Phi}_1^+$ and $\dot{\Phi}_1^-$.

Lemma 4.1. *Let U be a submanifold of $M(\beta)$ which is homeomorphic to a Seifert fibred space over the disk with two cone points. If U contains a closed curve which is non-null homotopic in $M(\beta)$, then either*

(i) ∂U is an incompressible torus in $M(\beta)$, or

(ii) $\overline{M(\beta) \setminus U}$ is a solid torus and $M(\beta)$ is a torus bundle over the circle which admits a Seifert structure with base orbifold of the form $S^2(a, b, c)$ where $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 1$.

Proof. Suppose that ∂U is compressible in $M(\beta)$. Then it is compressible in $\overline{M(\beta) \setminus U}$. The surgery of the torus ∂U using a compressing disk produces a separating 2-sphere. Since $M(\beta)$ is irreducible, this 2-sphere bounds a 3-ball B in $M(\beta)$. By hypothesis, U is not contained in B . Thus $\overline{M(\beta) \setminus U}$ is a solid torus. As $M(\beta)$ is irreducible (Assumption 2.8) and a Dehn filling of U , it is a Seifert fibred manifold over the 2-sphere with at most three cone points. But if such manifold contains an incompressible torus, it is a torus bundle over the circle and admits a Seifert structure of the type described in (ii). \diamond

Lemma 4.2. *Suppose that $(A, \partial A) \subseteq (X^\epsilon, S)$ is an embedded essential annulus. Let c_1, c_2 be the two boundary components of A .*

- (1) c_1 is essential in \widehat{S} if and only if c_2 is essential in \widehat{S} .
- (2) If c_1 and c_2 cobound an annulus E in \widehat{S} , then A is not parallel in \widehat{X}^ϵ to E . Furthermore E is essential in \widehat{S} .
- (3) If one of c_1 and c_2 is not essential in \widehat{S} , then c_1 and c_2 bound disjoint disks D_1 and D_2 in \widehat{S} such that $|D_1 \cap \partial M| = |D_2 \cap \partial M|$.

Proof. (1) This follows from the incompressibility of \widehat{S} in \widehat{X}^ϵ .

(2) Suppose otherwise that A is parallel to E in \widehat{X}^ϵ . Since A is essential in X^ϵ , $E \cap \partial M$ is not empty. But then we may consider E as an annulus in \widehat{F} , and if we replace E in \widehat{F} by A , we get a torus in $M(\beta)$ which is incompressible (since it is isotopic to \widehat{F}) but has fewer than m components of intersection with ∂M . This contradicts Assumption 2.2.

Now we show that E is essential in \widehat{S} . Suppose otherwise. Then one of c_1 and c_2 , say c_1 , bounds a disk D in \widehat{S} with interior disjoint from E . If A is non-separating in \widehat{X}^ϵ then $A \cup E$ is a non-separating Klein bottle or torus with compressing disk D with non-separating boundary. Compression of $A \cup E$ along D yields a non-separating 2-sphere in \widehat{X}^ϵ , which is impossible since \widehat{X}^ϵ is irreducible (Assumption 2.8). Thus A is separating in \widehat{X}^ϵ and therefore $T = E \cup A$ is a torus. Denote by W_1 and W_2 the two components of \widehat{X}^ϵ cut open along A and assume that W_1 is the component whose boundary is T . Assumption 2.8 shows that a regular neighborhood Y of $W_1 \cup D$ in \widehat{X}^ϵ is a 3-ball. Hence the disk $E \cup D$ is isotopic in Y to the disk $A \cup D$. So by Assumption 2.2 we have $|E \cap \partial M| = 0$. Therefore W_1 is contained in X^ϵ .

Since F is not contained in a regular neighborhood of ∂M , $T = \partial W_1$ is not parallel to ∂M . The hyperbolicity of M then implies that T is compressible in M . Since F is essential, we may assume that a compressing disk D_* for T in M is contained in X^ϵ . If the interior of D_* is disjoint from W_1 , then W_1 is contained in a 3-ball in M contrary to the fact that c_1 is essential in S . Thus $D_* \subset W_1$. The 2-sphere obtained by compressing T along D_* bounds a 3-ball contained in W_1 . (Otherwise ∂M would be contained in $W_1 \subseteq X^\epsilon$.) Hence W_1 is a solid torus.

But A is not parallel to E in W_1 . Therefore Y is once-punctured lens space with non-trivial fundamental group and not a 3-ball. This contradiction completes the proof of (2).

(3) By part (1) of this lemma, c_i bounds a disk D_i in \widehat{S} for each of $i = 1, 2$. If D_1 and D_2 are not disjoint, then one is contained in the other, say $D_1 \subseteq D_2$. Thus c_1 and c_2 bound an annulus E in D_2 . This contradicts part (2) of this theorem.

So D_1 and D_2 are disjoint. Let $d_i = |D_i \cap \partial M|$, $i = 1, 2$. Suppose otherwise that $d_1 \neq d_2$, say $d_1 < d_2$. Since \widehat{X}^ϵ is irreducible, $A \cup D_1 \cup D_2$ bounds a 3-ball B in \widehat{X}^ϵ with the interior of B disjoint from \widehat{S} . Then it is not hard to see that the disk D_2 can be isotoped rel its boundary in B to have at most d_1 intersection components with ∂M . This implies that the incompressible torus \widehat{F} can be isotoped in $M(\beta)$ to have less than m intersection components with ∂M , which again contradicts our minimality assumption on $m = |\partial F|$. \diamond

A *root torus* in (X^ϵ, S) is a solid torus $\Theta \subseteq X^\epsilon$ such that $\Theta \cap S$ is an incompressible annulus in $\partial\Theta$ whose winding number in Θ is at least 2 in absolute value. For instance, a regular neighbourhood of an embedded Möbius band $(B, \partial B) \subseteq (X^\epsilon, S)$ is a root torus. Note that for such a Θ , $\overline{\partial\Theta \setminus (\Theta \cap F)}$ is an essential annulus in (X^ϵ, F) .

Lemma 4.2(2) yields the following lemma.

Proposition 4.3. *If Θ is a root torus in (X^ϵ, S) , then $\Theta \cap S$ is an essential annulus in \widehat{S} . In particular, the boundary of a Möbius band properly embedded in X^ϵ is essential in \widehat{S} . \diamond*

Proposition 4.4. *A simple closed curve $c \subseteq \check{\check{\Phi}}_1^\epsilon$ is inner, respectively outer, if and only if $\tau_\epsilon(c)$ is inner, respectively outer. In particular, the image by τ_ϵ of a tight subsurface of $\check{\check{\Phi}}_1^\epsilon$ is a tight subsurface of $\check{\check{\Phi}}_1^\epsilon$.*

Proof. Suppose that c is inner. It suffices to see that $\tau_\epsilon(c)$ is inner as well. If c is \widehat{S} -essential, so is $\tau_\epsilon(c)$ since they cobound a singular annulus. Thus $\tau_\epsilon(c)$ is inner. Otherwise, c bounds a disk D in \widehat{S} containing at least two components of ∂S . If c cobounds an annulus with $\tau_\epsilon(c)$ in X^ϵ , for instance if c is contained in a product bundle component of $\check{\check{\Sigma}}_1^\epsilon$, then Lemma 4.2(3) shows that $\tau_\epsilon(c)$ is inner. In general, let ϕ be the component of $\check{\check{\Phi}}_1^\epsilon$ which contains c and Σ the component of $\check{\check{\Sigma}}_1^\epsilon$ which contains ϕ . Each boundary component of ϕ cobounds a vertical annulus in Σ with its image under τ_ϵ , so if c is boundary-parallel in ϕ , we are done. On the other hand, if it is not boundary-parallel in ϕ , $\tau_\epsilon(c)$ is not boundary-parallel in $\tau_\epsilon(\phi)$ and therefore cannot be boundary-parallel in S . Thus $\tau_\epsilon(c)$ is inner. \diamond

Proposition 4.5. *If ϕ is a tight component of $\check{\check{\Phi}}_{2j+1}^\epsilon$, then $h_{2j+1}^\epsilon(\phi) \cap \phi = \emptyset$.*

Proof. If $h_{2j+1}^\epsilon(\phi) \cap \phi \neq \emptyset$, then $h_{2j+1}^\epsilon(\phi) = \phi$. Hence as $h_{2j+1}^\epsilon = g_j \circ \tau_\epsilon \circ g_j^{-1}$ where $g_j = \tau_{(-1)^{j\epsilon}} \circ \tau_{(-1)^{j-1\epsilon}} \circ \dots \circ \tau_{-\epsilon}$, we have $\tau_\epsilon(\phi') = \phi'$ where ϕ' is the tight subsurface $g_j(\phi)$ of $\check{\check{\Phi}}_1^\epsilon$. It follows from Proposition 4.4 that if c is the inner boundary component of ϕ' , then $\tau_\epsilon(c) = c$. Thus c bounds a Möbius band properly embedded in X^ϵ . But then Proposition 4.3 implies that c is \widehat{S} -essential, contrary to the tightness of ϕ' . Thus $h_{2j+1}^\epsilon(\phi) \cap \phi = \emptyset$. \diamond

Proposition 4.6. *Let P be a component of Σ_1^ϵ and suppose that c is an inner boundary component of $P \cap S$ which is inessential in \widehat{S} . Let $D \subset \widehat{S}$ be the disk with boundary c and suppose that the component of $\overline{P \cap S}$ containing c is disjoint from $\text{int}(D)$. Then $P \cap D = c$ and if H is the component of $\widehat{X}^\epsilon \setminus P$ which contains D and A is the annulus $P \cap H$, then $(H, A) \cong (D^2 \times I, (\partial D^2) \times I)$. In particular, the I -bundle structure on P extends over $P \cup H$.*

Proof. Lemma 4.2(3) implies that there is a disk $D' \subset \widehat{S}$ disjoint from D such that $\partial D' = \tau_\epsilon(c)$. Thus $P \cap D = c$. Note that $\partial A = c \cup \tau_\epsilon(c)$. Then $D \cup A \cup D'$ is a 2-sphere which bounds a 3-ball $B \subseteq \widehat{X}^\epsilon$ such that $B \cap P = A$. The desired conclusions follow from this. \diamond

Lemma 4.7. *Suppose that $(A, \partial A) \subseteq (X^\epsilon, S)$ is a non-separating essential annulus with boundary components c_1, c_2 . Then c_1 and c_2 are essential in \widehat{S} . Further, either*

(i) $S = F$, X^- is a twisted I -bundle, and ∂A splits \widehat{F} into two annuli E_1, E_2 such that $|E_j \cap \partial M| = m/2$ and $A \cup E_j$ is a Klein bottle for $j = 1, 2$; or

(ii) $S = F_1 \cup F_2$ has two components where c_j is contained in F_j .

Proof. The components of ∂A are either both inessential in \widehat{S} or both essential (Lemma 4.2(1)). In the former case, Lemma 4.2(3) implies that there are disjoint disks D_1, D_2 in \widehat{S} such that $c_j = \partial D_j$. Then $D_1 \cup A \cup D_2$ is a 2-sphere in the irreducible manifold \widehat{X}^ϵ , which therefore bounds a 3-ball. This is impossible since A is non-separating. Thus the components of ∂A are essential in \widehat{S} .

If conclusion (ii) does not hold, there is a component S_0 of S such that ∂A splits \widehat{S}_0 into two annuli: $\widehat{S}_0 = E_1 \cup_{\partial A} E_2$. We assume, without loss of generality, that $|E_1 \cap \partial M| \leq m/2$. Since $E_j \cup A$ is non-separating and intersects ∂M in fewer than m components, $E_1 \cup A$ is a Klein bottle. A regular neighbourhood U of $E_1 \cup A$ is a twisted I -bundle over $E_1 \cup A$ and contains a loop which is not null-homotopic in M . Since \widehat{S} is isotopic into $\overline{M(\beta)} \setminus \overline{U}$, the latter cannot be a solid torus. Thus Lemma 4.1 implies that ∂U is an incompressible torus in $M(\beta)$. Hence $m \leq |\partial U \cap \partial M| = 2|(E_1 \cup A) \cap \partial M| = 2|E_1 \cap \partial M| \leq m$. It follows that $|E_1 \cap \partial M| = |E_2 \cap \partial M| = m/2$ and $|\partial U \cap \partial M| = m$. In particular, $(E_1 \cup A) \cap M$ is an $\frac{m}{2}$ -punctured Klein bottle properly embedded in M with twisted I -bundle neighbourhood $U \cap M$. Assumptions 2.3 and 2.5 then imply that $S = F$ and X^- is a twisted I -bundle. Hence situation (i) holds. \diamond

Lemma 4.8. *Suppose that $(A_j, \partial A_j)$ ($j = 1, 2$) are disjoint essential annuli contained in (X^ϵ, S) . If a boundary component c_1 of A_1 cobounds an annulus $E \subseteq S$ with a boundary component c_2 of A_2 and c_1 is \widehat{S} -inessential, then A_1 is isotopic to A_2 in X^ϵ .*

Proof. Let $\partial A_j = c_j \cup c'_j$ ($j = 1, 2$). We can suppose that c_j bounds a disk D_j in \widehat{S} ($j = 1, 2$) such that $D_2 = D_1 \cup E$. According to Lemma 4.2(3), c'_j bounds a disk D'_j in \widehat{S} such that $D_j \cap D'_j = \emptyset$ ($j = 1, 2$). Since $M(\beta)$ is irreducible, the 2-sphere $\Pi_2 = D_2 \cup A_2 \cup D'_2$ bounds a 3-ball $B_2 \subseteq \widehat{X}^\epsilon$.

Since $c_1 \subseteq \text{int}(D_2)$ and $\text{int}(A_1)$ is disjoint from Π_2 , A_1 is contained in B_2 . If $D'_1 \cap D_2 \neq \emptyset$, then $D'_1 \subseteq \text{int}(E) \subseteq S$. But this is impossible as c'_1 is essential in S . Thus $D'_1 \subseteq \text{int}(D'_2)$ and therefore c'_1 and c'_2 cobound an annulus $E' \subseteq D'_2 \subseteq \widehat{S}$. It then follows from Lemma 4.2(3) that $E' \subset S$.

The torus $T = E \cup A_1 \cup E' \cup A_2 \subset X^\epsilon$ is not boundary-parallel in the hyperbolic manifold M , so must compress in X^ϵ . It cannot be contained in a 3-ball in M since ∂A_1 is essential in S . Hence it bounds a solid torus Θ in X^ϵ . Proposition 4.3 shows that Θ is not a root torus, so A_1 must be parallel to A_2 in X^ϵ . \diamond

Proposition 4.9. *Let $(A, \partial A)$ be an essential annulus in (X^ϵ, S) such that a component c of ∂A cobounds an annulus $E \subseteq S$ with a component c' of ∂S . Then $(A, \partial A)$ is isotopic in (X^ϵ, S) to a component of $\partial M \cap X^\epsilon$.*

Proof. Let A' be the component of $\partial M \cap X^\epsilon$ which contains c' . Then A' is a properly embedded essential annulus in (X^ϵ, S) . Since c is inessential in \widehat{S} , Lemma 4.8 implies that A is isotopic to A' in X^ϵ . \diamond

As mentioned in §3.2, this corollary allows us to assume that Σ_1^ϵ is neatly embedded in X^ϵ .

Lemma 4.10. *Let ϕ_1 and ϕ_2 be components of Φ_1^ϵ , possibly equal, and suppose that there are a component c_1 of $\partial\phi_1$, a component c_2 of $\partial\phi_2$, and an annulus $E \subseteq \overline{S \setminus \Phi_1^\epsilon}$ such that $\partial E = c_1 \cup c_2$. Then E is essential in \widehat{S} .*

Proof. There are I -bundles $\Sigma_j \subseteq X^\epsilon$ such that $\phi_j \subseteq \Sigma_j \cap S$ is a component of the associated S^0 -bundle ($j = 1, 2$). Let $(A_j, \partial A_j) \subseteq (X^\epsilon, S)$ be the essential annulus in the frontier of Σ_j in X^ϵ which contains c_j ($j = 1, 2$).

If $A_1 = A_2$, then $A_1 \cup E$ is a torus in $X^\epsilon \subseteq M$ and so is either contained in a 3-ball in X^ϵ or bounds a solid torus $\Theta \subseteq X^\epsilon$. Since c_1 is essential in S , the latter must occur, and since A_1 is an essential annulus in (X^ϵ, S) , the winding number of E in Θ is at least 2. Thus Θ is a root torus of the type described. Proposition 4.3 now implies that E is essential in \widehat{S} .

Next suppose that $A_1 \neq A_2$, so these two annuli are disjoint. Note that they cannot be parallel as otherwise the I -bundle structures on Σ_1 and Σ_2 can be extended across an embedded $(E \times I, E \times \partial I) \subseteq (X^\epsilon, S)$, which contradicts the defining properties of Φ_1^ϵ . Hence Lemma 4.8 implies that E is essential in \widehat{S} . \diamond

Proposition 4.11. *Let ϕ_1 and ϕ_2 be components of $\check{\Phi}_j^\epsilon$, possibly equal, and suppose that there are a component c_1 of $\partial\phi_1$, a component c_2 of $\partial\phi_2$, and an annulus $E \subseteq \overline{S \setminus \check{\Phi}_j^\epsilon}$ such that $\partial E = c_1 \cup c_2$. Then E is essential in \widehat{S} .*

Proof. As $\check{\Phi}_0^\epsilon = S$, there is an integer k such that $1 \leq k \leq j$ and E is contained in $\check{\Phi}_{k-1}^\epsilon$ but not in $\check{\Phi}_k^\epsilon$. Then $\phi_1 \cup \phi_2 \subseteq \check{\Phi}_j^\epsilon \subseteq \check{\Phi}_k^\epsilon$. Further, as E is not contained in $\check{\Phi}_k^\epsilon$, there must be inner boundary component of $\check{\Phi}_k^\epsilon$, call it c_0 , contained in E . If there is an arc a in $E \cap \check{\Phi}_k^\epsilon$ connecting c_1 and c_2 , then $c_0 \subset (E \setminus a)$ and therefore c_0 is contained in a disk in $E \subset S$, which contradicts

the essentiality of the inner components of $\partial\check{\Phi}_k^\epsilon$ in S . Hence $(E, \partial E) \subseteq \overline{(\check{\Phi}_{k-1}^\epsilon \setminus \check{\Phi}_k^\epsilon, \partial\check{\Phi}_k^\epsilon)}$. When $k = 1$ set $E_0 = E$ and when $k > 1$ set $E_0 = (\Pi_{i=1}^{k-1} \tau_{(-1)^{k-i-1}\epsilon})(E)$ so that $(E_0, \partial E_0) \subseteq \overline{(S \setminus \Phi_1^{(-1)^{k-1}\epsilon}, \partial\Phi_1^{(-1)^{k-1}\epsilon})}$. By Lemma 4.10, E_0 is \widehat{S} -essential, and therefore E is as well. \diamond

5. PAIRS OF EMBEDDED ESSENTIAL ANNULI IN (M, S)

In this section we consider pairs of essential annuli lying on either side of S in M .

Lemma 5.1. *Suppose that $S = F$ and that there are embedded, separating, essential annuli $(A^+, \partial A^+) \subseteq (X^+, F)$ and $(A^-, \partial A^-) \subseteq (X^-, F)$ such that ∂A^+ and ∂A^- are four parallel essential mutually disjoint curves in \widehat{F} . Then*

(1) ∂A^ϵ does not separate $\partial A^{-\epsilon}$ in \widehat{F} .

(2) Let E be an annulus in \widehat{F} bounded by a component of ∂A^+ and a component ∂A^- , with the interior of E disjoint from $A^+ \cup A^-$. Then $|E \cap \partial M| = m/2$.

Proof. For each ϵ , the boundary ∂A^ϵ of A^ϵ separates \widehat{F} into two parallel essential annuli, E_1^ϵ and E_2^ϵ , in \widehat{F} .

In order to prove the first assertion of the lemma, assume that ∂A^ϵ separates $\partial A^{-\epsilon}$ in \widehat{F} , that is, ∂A^- is not contained in E_1^+ or E_2^+ . Then $|E_j^+ \cap \partial A^-| = 1$ for $j = 1, 2$. Hence ∂A^- splits E_1^+ and E_2^+ into four annuli, which we denote by A_1, A_2, A_3, A_4 with $A_1 = E_1^+ \cap E_1^-$, $A_2 = E_1^+ \cap E_2^-$, $A_3 = E_2^+ \cap E_2^-$, $A_4 = E_2^+ \cap E_1^-$. Note that A_1, \dots, A_4 are four parallel essential annuli in \widehat{F} with disjoint interiors and with $A_1 \cup \dots \cup A_4 = \widehat{F}$.

Suppose that $|A_1 \cap \partial M| > 0$. Then the torus $A^+ \cup E_2^+$ bounds a solid torus V^+ in \widehat{X}^+ (since it intersects ∂M in fewer than m components) such that A^+ is not parallel to E_2^+ in V^+ (Lemma 4.2). Similarly the torus $A^- \cup E_2^-$ bounds a solid torus V^- in \widehat{X}^- such that A^- is not parallel to E_2^- in V^- . Hence $U = V^+ \cup_{A_3} V^-$ is a submanifold of $M(\beta)$ which is a Seifert fibred space over the disk with two cone points. Also a core circle of A_3 is non-null homotopic in $M(\beta)$. If ∂U compresses in $M(\beta)$, then Lemma 4.1 implies that $V = \overline{M(\beta) \setminus U}$ is a solid torus. Hence A_1 is ∂ -parallel in V and therefore is isotopic to either $A^+ \cup A_2$ or to $A^- \cup A_4$ in V , contrary to construction. Thus ∂U is an incompressible torus in $M(\beta)$. But $\partial U = A_2 \cup A^+ \cup A_4 \cup A^-$ intersects ∂M in fewer than m components, which contradicts Assumption 2.2. Thus $|A_1 \cap \partial M| = 0$, and similarly $|A_j \cap \partial M| = 0$ for $j = 2, 3, 4$, which is impossible. This proves (1).

Next we prove the lemma's second assertion. By (1), we can suppose that ∂A^- is contained in E_1^+ or E_2^+ , say, E_1^+ . Then we may assume that E_1^- is contained in E_1^+ and that E_2^+ is contained in E_2^- . Let E, E_* be the two annulus components of $E_2^- \cap E_1^+$. We need to show that $|E \cap \partial M| = |E_* \cap \partial M| = m/2$.

First we show that $|E_1^- \cap \partial M| = |E_2^+ \cap \partial M| = 0$. Suppose otherwise that $|E_1^- \cap \partial M| \neq 0$, say. Then the torus $A^+ \cup E_2^+$ bounds a solid torus V^+ in \widehat{X}^+ such that A^+ is not parallel to E_2^+ in V^+ , and the torus $A^- \cup E_2^-$ bounds a solid torus V^- in \widehat{X}^- such that A^- is not parallel to E_2^-

in V^- . Hence $U = V^+ \cup_{E_2^+} V^-$ is a submanifold of $M(\beta)$ which is a Seifert fibred space over the disk with two cone points. Also the center circle of E_2 is non-null homotopic in $M(\beta)$. As in the proof of assertion (1), we can use Lemma 4.1 to see that ∂U is an incompressible torus in $M(\beta)$. But $\partial U = A^- \cup E \cup A^+ \cup E_*$ intersects ∂M in fewer than m components, contradicting Assumption 2.2. Thus $|E_1^- \cap \partial M| = 0$ and a similar argument yields $|E_2^+ \cap \partial M| = 0$. Hence $\partial F \subseteq E \cup E_*$.

Next we prove $|E \cap \partial M| = |E_* \cap \partial M| = m/2$. Suppose otherwise, say $|E \cap \partial M| < m/2$ and $|E_* \cap \partial M| > m/2$. By the previous paragraph, the torus $A^+ \cup E_2^+$ bounds a solid torus V^+ in \widehat{X}^+ such that A^+ is not parallel to E_2^+ in V^+ , and the torus $A^- \cup E_1^-$ bounds a solid torus V^- in \widehat{X}^- such that A^- is not parallel to E_1^- in V^- . Hence a regular neighborhood U of $V^+ \cup E \cup V^-$ in $M(\beta)$ is a submanifold of $M(\beta)$ which is a Seifert fibred space over the disk with two cone points, and the core circle of E , which is contained in U , is non-null homotopic in $M(\beta)$. As above, Lemma 4.1 implies that ∂U is incompressible in $M(\beta)$. But by construction, $|\partial U \cap \partial M| < m$, contradicting Assumption 2.2. Thus $|E \cap \partial M| = |E_* \cap \partial M| = m/2$, which completes the proof of the lemma. \diamond

Proposition 5.2. *Suppose that $S = F$ and that there are embedded, separating, essential annuli $(A^+, \partial A^+) \subseteq (X^+, F)$ and $(A^-, \partial A^-) \subseteq (X^-, F)$ such that ∂A^+ and ∂A^- are four parallel essential mutually disjoint curves in \widehat{F} . Then no component of ∂A^+ is isotopic in F to a component of ∂A^- .*

Proof. Otherwise we may isotope A^ϵ in X^ϵ , so that ∂A^+ and ∂A^- remain disjoint but ∂A^+ separates ∂A^- in \widehat{F} . This is impossible by Lemma 5.1. \diamond

Proposition 5.3. *Suppose that $S = F$ and that there is an embedded Möbius band $(B, \partial B) \subseteq (X^\epsilon, F)$. Then ∂B cannot be isotopic in F to a boundary component of an embedded, separating, essential annulus $(A, \partial A) \subseteq (X^{-\epsilon}, F)$.*

Proof. Suppose otherwise. By Proposition 4.3, ∂B is essential in \widehat{F} . Let P be a regular neighborhood of B in X^ϵ . Then the frontier A_* of P in X^ϵ is an essential annulus in X^ϵ . Also ∂A and ∂A_* are essential curves in \widehat{F} which can be assumed to be mutually disjoint since ∂B is isotopic in F to a component of ∂A and each component of ∂A_* is isotopic to ∂B in F . But such a situation is impossible by Proposition 5.2. \diamond

Lemma 5.4. *Suppose that $S = F$ and that there are disjoint embedded, essential annuli $(A^+, \partial A^+) \subseteq (X^+, F)$ and $(A^-, \partial A^-) \subseteq (X^-, F)$ such that A^+ is separating, A^- is non-separating, and ∂A^+ and ∂A^- are four parallel essential mutually disjoint curves in \widehat{F} which split it into four annuli E_1, E_2, E_3, E_4 where $E_i \cap E_{i+1} \neq \emptyset$ for all $i \pmod{4}$.*

(1) *Suppose that ∂A^+ does not separate ∂A^- in \widehat{F} and that the annuli E_i are numbered so that $\partial A^+ = \partial E_1$ and $\partial A^- = \partial E_3$. Then $|E_1 \cap \partial F| = 0$.*

(2) *Suppose that ∂A^+ separates ∂A^- in \widehat{F} and that the annuli E_i are numbered so that the components of ∂A^+ are $E_1 \cap E_2$ and $E_3 \cap E_4$. Then $|E_1 \cap \partial F| = |E_4 \cap \partial F|$, $|E_2 \cap \partial F| = |E_3 \cap \partial F|$, and $|E_1 \cap \partial F| + |E_2 \cap \partial F| = m/2$.*

Proof. The proof is based on Lemma 4.7. Since we have assumed that $S = F$, conclusion (i) of this lemma holds.

Assume first that ∂A^+ does not separate ∂A^- in \widehat{F} . Then Lemma 4.7 implies that $|E_3 \cap \partial F| = |E_1 \cap \partial F| + |E_2 \cap \partial F| + |E_4 \cap \partial F| = m/2$. Since $A^+ \cup E_2 \cup A^- \cup E_4$ is a Klein bottle, $|E_2 \cap \partial F| + |E_4 \cap \partial F| \geq m/2$ and therefore $|E_1 \cap \partial F| = 0$.

Next assume that ∂A^+ separates ∂A^- in \widehat{F} . A tubular neighbourhood U of the Klein bottle $A^+ \cup E_1 \cup A^- \cup E_3$ is a twisted I -bundle over the Klein bottle, and as no Dehn filling of U is toroidal, the torus ∂U must be incompressible in $M(\beta)$ (cf. Assumption 2.8). Hence $m \leq |\partial U \cap \partial M| \leq 2(|E_1 \cap \partial F| + |E_3 \cap \partial F|)$ and therefore $|E_1 \cap \partial F| + |E_3 \cap \partial F| \geq m/2$. Similarly, consideration of the Klein bottle $A^+ \cup E_2 \cup A^- \cup E_4$ shows that $|E_2 \cap \partial F| + |E_4 \cap \partial F| \geq m/2$. On the other hand, Lemma 4.7 implies that $|E_1 \cap \partial F| + |E_2 \cap \partial F| = |E_3 \cap \partial F| + |E_4 \cap \partial F| = m/2$, from which we deduce the desired conclusion. \diamond

6. THE DEPENDENCE OF THE NUMBER OF TIGHT COMPONENTS OF $\check{\Phi}_j^\epsilon$ ON j

Let \mathcal{T}_j^ϵ be the union of the tight components of $\check{\Phi}_j^\epsilon$ and set

$$t_j^\epsilon = |\mathcal{T}_j^\epsilon|$$

If j is odd, the free involution $h_j : \check{\Phi}_j^\epsilon \rightarrow \check{\Phi}_j^\epsilon$ preserves \mathcal{T}_j^ϵ but none of its components (cf. Proposition 4.5). Thus t_j^ϵ is even for j odd. Further, as $\check{\Phi}_j^\epsilon \cong \check{\Phi}_j^{-\epsilon}$ for j even, $t_{2k}^+ = t_{2k}^-$ for all k .

Lemma 6.1. *Suppose that $C \subseteq (S \setminus \check{\Phi}_j^\epsilon)$ is an essential simple closed curve which bounds a disk $D \subseteq \widehat{S}$. Then D contains a tight component of $\check{\Phi}_j^\epsilon$. Further, if C is not isotopic in S into the boundary of a tight component of $\check{\Phi}_j^\epsilon$ (i.e. $D \cap S$ is not isotopic in S to a tight component of $\check{\Phi}_j^\epsilon$), then D contains at least two tight components of $\check{\Phi}_j^\epsilon$.*

Proof. Since C is essential, D contains at least one boundary component of S and hence at least one component of $\check{\Phi}_j^\epsilon$. Amongst all the inner boundary components of $\check{\Phi}_j^\epsilon$ which are contained in D , choose one, C_1 say, which is innermost in D . It is easy to see that this circle is the inner boundary component of a tight component ϕ_1 of $\check{\Phi}_j^\epsilon$. This proves the first assertion of the lemma.

Next suppose that C is not isotopic in S into the boundary of a tight component of $\check{\Phi}_j^\epsilon$. Then C and C_1 do not cobound an annulus in $D \cap S$, so there is a component of ∂S contained in $\overline{D \setminus \phi_1}$. Hence if $\phi_1, \phi_2, \dots, \phi_n$ are the components of $\check{\Phi}_j^\epsilon$ contained in $D \cap S$, then $n \geq 2$. If every inner boundary component of $\phi_2 \cup \phi_3 \cup \dots \cup \phi_n$ is essential in the annulus $\widehat{D \setminus \phi_1}$, some such boundary component cobounds an annulus $E \subseteq S$ with C_1 . Without loss of generality we may suppose $\partial E = C_1 \cup C_2$ where $C_2 \subseteq \partial \phi_2$. But this is impossible as Proposition 4.11 would then imply that E is essential in \widehat{S} . Hence some inner boundary component of $\phi_2 \cup \phi_3 \cup \dots \cup \phi_n$ bounds a subdisk D' of D which is disjoint from ϕ_1 , the argument of the first paragraph of this proof shows that D contains another tight component of $\check{\Phi}_j^\epsilon$, so we are done. \diamond

An immediate consequence of the lemma is the following corollary.

Corollary 6.2.

(1) If $\check{\Phi}_j^\epsilon$ has a component ϕ which is contained in a disk $D \subseteq \widehat{S}$, then either ϕ is tight or D contains at least two tight components of $\check{\Phi}_j^\epsilon$.

(2)(a) If ϕ_0 is a tight component of $\check{\Phi}_j^\epsilon$, there is a tight component ϕ_1 of $\check{\Phi}_{j+1}^\epsilon$ contained in ϕ_0 .

(b) If ϕ_1 is not isotopic to ϕ_0 in S , there are at least two tight components of $\check{\Phi}_{j+1}^\epsilon$ contained in ϕ_0 . \diamond

Proposition 6.3.

(1)(a) $t_j^\epsilon \leq t_{j+1}^\epsilon$ with equality if and only if \mathcal{T}_j^ϵ is isotopic to $\mathcal{T}_{j+1}^\epsilon$ in S .

(b) $t_j^\epsilon \leq t_{j+1}^{-\epsilon}$

(2) If $0 < t_j^\epsilon = t_{j+2}^\epsilon$, then $t_j^\epsilon = |\partial S|$, so \mathcal{T}_j^ϵ is a regular neighbourhood of ∂S .

Proof. Part (1)(a) follows immediately from Corollary 6.2. For part (1)(b), note that if j is odd then $t_j^\epsilon \leq t_{j+1}^\epsilon = t_{j+1}^{-\epsilon}$, while if j is even, $t_j^\epsilon = t_j^{-\epsilon} \leq t_{j+1}^{-\epsilon}$.

Next we prove part (2). Suppose that $0 < t_j^\epsilon = t_{j+2}^\epsilon$. Then Lemma 6.1 implies that up to isotopy, $\mathcal{T}_j^\epsilon = \mathcal{T}_{j+2}^\epsilon$. We claim that $(\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_{j+2}^\epsilon) = \mathcal{T}_j^\epsilon$, at least up to isotopy fixed on ∂S . To see this, first note that $(\tau_{-\epsilon}\tau_\epsilon)(\check{\Phi}_{j+2}^\epsilon) \subseteq \check{\Phi}_j^\epsilon$. Fix a tight component ϕ_0 of $\check{\Phi}_j^\epsilon$ and let $\phi_1, \phi_2, \dots, \phi_n$ be the components of $\check{\Phi}_{j+2}^\epsilon$ such that for each $i = 1, 2, \dots, n$, $\phi'_i = (\tau_{-\epsilon}\tau_\epsilon)(\phi_i) \subseteq \phi_0$. Since each component of $\partial S \cap \phi_0$ is contained in some ϕ'_i , the argument of the first paragraph of the proof of Lemma 6.1 shows that at least one of the ϕ'_i , or equivalently ϕ_i , is tight. Since ϕ_0 is an arbitrary tight component of $\check{\Phi}_j^\epsilon$ and $t_j^\epsilon = t_{j+2}^\epsilon$, it follows that $(\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_{j+2}^\epsilon) \subseteq \mathcal{T}_j^\epsilon$ and each component of \mathcal{T}_j^ϵ contains a unique component of $(\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_{j+2}^\epsilon)$. Note as well that as $\mathcal{T}_j^\epsilon = \mathcal{T}_{j+2}^\epsilon$, we have $|\partial S \cap \mathcal{T}_j^\epsilon| = |\partial S \cap \mathcal{T}_{j+2}^\epsilon| = |\partial S \cap (\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_{j+2}^\epsilon)|$, so if ϕ_1 is a tight component of $\mathcal{T}_{j+2}^\epsilon$ and ϕ_0 the tight component of \mathcal{T}_j^ϵ containing $\phi'_1 = (\tau_{-\epsilon}\tau_\epsilon)(\phi_1)$, then $|\partial S \cap \phi'_1| = |\partial S \cap \phi_0|$. But then as ϕ_0 and ϕ'_1 are tight, ϕ'_1 is isotopic to ϕ_0 by an isotopy fixed on ∂S . Hence we can assume that $(\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_{j+2}^\epsilon) = \mathcal{T}_j^\epsilon$, or in other words, $(\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_j^\epsilon) = \mathcal{T}_j^\epsilon$. It follows that $(\tau_{-\epsilon}\tau_\epsilon)^k(\mathcal{T}_j^\epsilon) = \mathcal{T}_j^\epsilon \subseteq \check{\Phi}_j^\epsilon$, and so $\mathcal{T}_j^\epsilon \subseteq \check{\Phi}_{j+2k}^\epsilon$ for all k . But since F is neither a fibre nor a semi-fibre, $\check{\Phi}_{j+2k}^\epsilon$ is a regular neighbourhood of ∂S for large k (cf. [BCSZ1, proof of Theorem 5.4.1]). Thus \mathcal{T}_j^ϵ is a union of annuli.

The boundary components of S can be numbered $b_1, b_2, \dots, b_{|\partial S|}$ so that they arise successively around ∂M and $(\tau_{-\epsilon}\tau_\epsilon)(b_i) = b_{i+(-1)^i 2}$, where the indices are considered (mod $|\partial S|$). Hence as $(\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_j^\epsilon) = (\tau_{-\epsilon}\tau_\epsilon)(\mathcal{T}_{j+2}^\epsilon) = \mathcal{T}_j^\epsilon$, $\partial \mathcal{T}_j^\epsilon \cap \partial S$ is the union of either all even-indexed b_i , or all odd-indexed b_i , or all the b_i (recall that we have assumed $t_j^\epsilon > 0$). In particular, for either all even i or all odd i , the component of $\check{\Phi}_j^\epsilon$ containing b_i is an annulus. If j is odd, we have a free involution $\check{h}_j^\epsilon : \check{\Phi}_j^\epsilon \rightarrow \check{\Phi}_j^\epsilon$ which the reader will verify preserves \mathcal{T}_j^ϵ and exchanges the even-indexed b_i with the odd-indexed b_i . Hence for any i , the component of $\check{\Phi}_j^\epsilon$ containing b_i is an annulus. Thus $t_j^\epsilon = m$, so \mathcal{T}_j^ϵ is a regular neighbourhood of ∂S .

Next suppose that j is even. After possibly adding 1 (mod $|\partial S|$) to the indices of the labels of the components of ∂S , we can assume that $\partial \mathcal{T}_j^\epsilon \cap \partial S$ contains the union of all even-indexed b_i . Then as $\check{\Phi}_{j+1}^{-\epsilon} = \tau_{-\epsilon}(\check{\Phi}_j^\epsilon \wedge \check{\Phi}_1^{-\epsilon}) \supseteq \tau_{-\epsilon}(\mathcal{T}_j^\epsilon \wedge \check{\Phi}_1^{-\epsilon})$, the component of $\mathcal{T}_{j+1}^{-\epsilon}$ containing an odd-indexed b_i is an annulus. Consideration of the free involution $\check{h}_{j+1}^{-\epsilon} : \check{\Phi}_{j+1}^{-\epsilon} \rightarrow \check{\Phi}_{j+1}^{-\epsilon}$ shows that the same is true for the even-indexed b_i . Thus $m = t_{j+1}^{-\epsilon} \leq t_{j+2}^\epsilon = t_j^\epsilon$. It follows that \mathcal{T}_j^ϵ is a regular neighbourhood of ∂S . \diamond

Corollary 6.4.

(1) *If some non-tight component of $\check{\Phi}_1^\epsilon$ has an \widehat{S} -inessential inner boundary component, then $t_1^\epsilon \geq 4$.*

(2) *If $\text{genus}(\check{\Phi}_1^\epsilon) = 1$ but $\check{\Phi}_1^\epsilon \neq S$, then $t_1^\epsilon \geq 4$.*

Proof. Let ϕ be a non-tight component of $\check{\Phi}_1^\epsilon$ and c an \widehat{S} -inessential inner boundary component of ϕ . Let $D \subseteq \widehat{S}$ be the disk with boundary c . Lemma 6.1 implies that D contains a tight component ϕ_0 . By Lemma 4.10, c is not isotopic in S into the boundary of ϕ_0 , so D contains at least two tight components (Lemma 6.1). It follows from Lemma 4.2(3) that $\tau_\epsilon(c)$ bounds a disk $D' \subseteq \widehat{S}$ disjoint from D and as above, D' contains at least two tight components of $\check{\Phi}_1^\epsilon$. Thus $t_1^\epsilon \geq 4$, which proves (1).

Next suppose that $\text{genus}(\check{\Phi}_1^\epsilon) = 1$ but $\check{\Phi}_1^\epsilon \neq S$. Then there is a component $\phi \neq S$ of $\check{\Phi}_1^\epsilon$ of genus 1. In particular, ϕ is not tight. Since $\phi \neq S$, it has inner boundary components. Since $\text{genus}(\phi) = 1$, each such inner boundary component is \widehat{S} -inessential. Hence part (2) of the corollary follows from part (1). \diamond

7. THE STRUCTURE OF $\check{\Phi}_1^\epsilon$ AND THE TOPOLOGY OF X^ϵ

In this section we study how the existence of a component ϕ of $\check{\Phi}_1^\epsilon$ such that $\widehat{\phi}$ contains an \widehat{S} -essential annulus constrains $\check{\Phi}_1^\epsilon$ and the topology of \widehat{X}^ϵ . The following construction will be useful to our analysis.

Let P be a component of $\check{\Sigma}_1^\epsilon$. For each \widehat{S} -inessential inner component c of $P \cap S$ let $D_c \subset \widehat{S}$ be the disk with boundary c and suppose that the component of $P \cap S$ containing c is disjoint from $\text{int}(D_c)$. The component H_c of $\widehat{X}^\epsilon \setminus P$ containing c satisfies $(H_c, H_c \cap P) \cong (D^2 \times I, (\partial D^2) \times I)$ (cf. Proposition 4.6). Let

$$(7.0.1) \quad Q_P = \widehat{P} \cup (\cup_c H_c)$$

where c ranges over all \widehat{S} -inessential inner components of $P \cap S$ such that $P \cap \text{int}(D_c) = \emptyset$. The I -fibre structure on P extends over Q_P .

We prove the following results.

Proposition 7.1. *Suppose that F is separating in M , so $S = F$ is connected.*

(1) *There is at most one component P of $\check{\Sigma}_1^\epsilon$ such that $\widehat{P \cap F}$ contains an \widehat{F} -essential annulus.*

- (2) *There is exactly one such component if $\dot{\Sigma}_1^\epsilon$ contains a twisted I -bundle.*
- (3) *Suppose that P is a component of $\dot{\Sigma}_1^\epsilon$ such that $\widehat{P \cap F}$ contains an \widehat{F} -essential annulus. Then \widehat{X}^ϵ admits a Seifert structure and if*
- (a) *genus($\widehat{P \cap F}$) = 1, then \widehat{X}^ϵ is a twisted I -bundle over the Klein bottle.*
 - (b) *genus($\widehat{P \cap F}$) = 0, then an \widehat{F} -essential annulus in $\widehat{P \cap F}$ is vertical in the Seifert structure and Q_P splits \widehat{X}^ϵ into a union of solid tori. Moreover, if*
 - (i) *P is a twisted I -bundle, then \widehat{X}^ϵ has base orbifold a disk with two cone points, at least one of which has order 2.*
 - (ii) *P is a product I -bundle and Q_P separates \widehat{X}^ϵ , then \widehat{X}^ϵ has base orbifold a disk with two cone points.*
 - (iii) *P is a product I -bundle and Q_P does not separate \widehat{X}^ϵ , then X^- is a twisted I -bundle and \widehat{X}^ϵ has base orbifold a Möbius band with at most one cone point.*

Proposition 7.2. *Suppose that F is non-separating in M , so $S = F_1 \cup F_2$ is not connected.*

- (1) *$\dot{\Sigma}_1^+$ is a (possibly empty) product bundle and for each $j = 1, 2$ and component P of $\dot{\Sigma}_1^+$, genus($P \cap F_j$) = 0.*
- (2) *If P is a component of $\dot{\Sigma}_1^+$ such that $P \cap S \subseteq P \cap F_j$ for some j , then $\widehat{P \cap F_j}$ contains no \widehat{S} -essential annulus.*
- (3) *If $t_1^+ = 0$, then $\dot{\Sigma}_1^+$ has exactly one component P and for each $j = 1, 2$, $\widehat{P \cap F_j}$ is an annulus which is essential in $\widehat{F_j}$. Further, \widehat{X}^+ admits a Seifert structure with base orbifold an annulus with exactly one cone point.*

We consider the cases S connected and S disconnected separately.

7.1. S is connected. In this subsection we prove Proposition 7.1.

Lemma 7.3. *Suppose that F is separating in M and $(A, \partial A) \subseteq (X^\epsilon, F)$ is an essential separating annulus whose boundary separates \widehat{F} into two annuli E_1 and E_2 . If $|E_1 \cap \partial M| < m$ then $A \cup E_1$ bounds a solid torus in \widehat{X}^ϵ in which A has winding number at least 2. Hence if either*

- (a) *$|E_1 \cap \partial M| < m$ and $|E_2 \cap \partial M| < m$; or*
- (b) *$|E_1 \cap \partial M| = m$ and $E_1 \cup A$ bounds a solid torus in \widehat{X}^ϵ ,*

then A splits \widehat{X}^ϵ into two solid tori in each of which A has winding number at least 2. In particular, \widehat{X}^ϵ admits a Seifert structure with base orbifold a disk with two cone points in which A is vertical.

Proof. If $|E_1 \cap \partial M| < m$, then $A \cup E_1$ is a torus which compresses in $M(\beta)$ but is not contained in a 3-ball. Hence it bounds a solid torus V which is necessarily contained in \widehat{X}^ϵ . Lemma 4.2(2) shows that A has winding number at least 2 in V . It follows that if condition (a) holds, \widehat{X}^ϵ

admits a Seifert structure with base orbifold a 2-disk with two cone points. Note that A is vertical in this structure and splits \widehat{X}^ϵ into two solid tori. A similar argument yields the same conclusion under condition (b). \diamond

Proof of part (3) of Proposition 7.1. First suppose that $\text{genus}(\widehat{P \cap F}) = 1$. Then P is necessarily a twisted I -bundle and $\phi = \widehat{P \cap F}$ is connected. Further, each inner boundary component of $P \cap F$ is inessential in \widehat{F} . Thus $\widehat{X}^\epsilon = Q_P$ is a twisted I -bundle over the Klein bottle. Hence part (3)(a) of Proposition 7.1 holds.

Next suppose that $\text{genus}(\widehat{P \cap F}) = 0$ and ϕ is a component of $P \cap F$. Then ϕ has two inner boundary components, c_1, c_2 say, which are \widehat{F} -essential. Any other inner boundary component c of ϕ is inessential in \widehat{F} so Q_P is either a twisted I -bundle over a Möbius band or product I -bundle over an annulus.

Let A_1, A_2 be the vertical annuli in the frontier of P , possibly equal, that contain c_1, c_2 respectively. There are three cases to consider.

Case 1. P is a twisted I -bundle.

In this case, $A_1 = A_2$ and Q_P is a twisted I -bundle over a Möbius band. In particular Q_P is a solid torus in which a core of $\widehat{\phi}$ has winding number 2. Lemma 7.3 shows that Proposition 7.1(3)(b)(i) holds.

Case 2. P is a product I -bundle and Q_P separates X^ϵ .

Then $A_1 \neq A_2$ where A_1 is separating in X^ϵ and Q_P is a product I -bundle over an annulus. Let V and W be the components of the exterior of Q_P in \widehat{X}^ϵ and define E_1, E_2 to be the \widehat{F} -essential annuli $V \cap \widehat{F}, W \cap \widehat{F}$. Since $|P \cap \partial F| > 0$, we have $|E_1 \cap \partial F| < m$ and $|E_2 \cap \partial F| < m$. Then Lemma 7.3 implies that both V and W are solid tori and therefore that Proposition 7.1(3)(b)(ii) holds.

Case 3. P is a product I -bundle and Q_P does not separate X^ϵ .

Here $A_1 \neq A_2$ where A_1 is non-separating and X^- is a twisted I -bundle by Lemma 4.7. Also the boundary of the complement of the interior of Q_P in X^ϵ is a torus which intersects ∂M in fewer than m components but is not contained in any 3-ball in $M(\beta)$. Thus it bounds a solid torus V in \widehat{X}^ϵ , from which we can see that Proposition 7.1(3)(b)(iii) holds. \diamond

Proof of parts (1) and (2) of Proposition 7.1. If $\dot{\Sigma}_1^\epsilon$ contains a twisted I -bundle P , then P contains a subbundle homeomorphic to a Möbius band. Proposition 4.3 then shows that $P \cap F$ contains an \widehat{F} -essential annulus. Thus part (1) implies part (2). We prove part (1) by contradiction.

Suppose that $\dot{\Sigma}_1^\epsilon$ has at least two components P_1, P_2 such that $\widehat{P_i \cap F}$ contains an \widehat{F} -essential annulus for $i = 1, 2$. Let $\phi_i = P_i \cap F$ be the horizontal boundary of P_i ($i = 1, 2$). Clearly, both ϕ_1 and ϕ_2 have genus 0. Since each properly embedded incompressible annulus in a solid torus is separating, Proposition 7.1(3), which we proved above, implies that both Q_{P_1} and Q_{P_2} are separating in \widehat{X}^ϵ and split it into a union of solid tori. We have three cases to consider.

Case 1. ϕ_i is connected for $i = 1, 2$.

Then Q_{P_i} is a twisted I -bundle over a Möbius band whose frontier in \widehat{X}^ϵ is an essential annulus A_i in X^ϵ which is not parallel to the annulus $\psi_i = Q_{P_i} \cap \widehat{F}$ in Q_{P_i} . Let E_1, E_2 be the components of the closure of the complement of $\psi_1 \cup \psi_2$ in \widehat{F} . Then the torus $E_1 \cup A_1 \cup E_2 \cup A_2$ bounds a solid torus V in \widehat{X}^ϵ in which A_1 is not parallel to A_2 . Therefore $U = V \cup \widehat{P}_1$ satisfies the hypotheses of Lemma 4.1. Since \widehat{F} is isotopic into $\overline{M(\beta)} \setminus \overline{U}$, the latter cannot be a solid torus. Thus $\partial U = \psi_1 \cup E_1 \cup A_2 \cup E_2$ is incompressible in $M(\beta)$. But this torus intersects ∂M in fewer than m components, which contradicts Assumption 2.2.

Case 2. ϕ_1 is connected but ϕ_2 is not.

Then Q_{P_1} is a twisted I -bundle over a Möbius band and the frontier of Q_{P_1} in \widehat{X}^ϵ is an essential annulus $A_1 \subseteq X^\epsilon$ which is not parallel to the annulus $\psi_1 = Q_{P_1} \cap \widehat{F}$ in Q_{P_1} . Further, ϕ_2 has two components, ϕ_{21}, ϕ_{22} say, and P_2 is a product I -bundle over ϕ_{21} . The frontier of Q_{P_2} is a pair of essential annuli $A_{21}, A_{22} \subseteq X^\epsilon$. We noted above that Q_{P_2} is separating in X^ϵ , and so the same is true for A_{21} and A_{22} .

We may suppose that A_{21} is adjacent to A_1 . That is, $\partial A_1 \cup \partial A_{21}$ cobounds the union of two disjoint annuli $E_1, E_2 \subseteq \widehat{F}$ whose interiors are disjoint from ϕ_1, ϕ_2 . Then the torus $A_1 \cup E_1 \cup A_{21} \cup E_2$ bounds a solid torus V in \widehat{X}^ϵ such that A_1 is not parallel to A_{21} in V . Therefore $U = V \cup \widehat{P}_1$ is a submanifold of $M(\beta)$ satisfying the hypotheses of Lemma 4.1. As in case 1, this lemma implies that $\partial U = \psi_1 \cup E_1 \cup A_{21} \cup E_2$ is incompressible in $M(\beta)$. But this torus intersects ∂M in fewer than m components, contrary to Assumption 2.2.

Case 3. Neither ϕ_1 nor ϕ_2 is connected.

The frontier of Q_{P_i} in \widehat{X}^ϵ is a pair of annuli A_{i1}, A_{i2} contained in X^ϵ . We may assume that ∂A_{12} and ∂A_{21} cobound two annuli E_1, E_2 in \widehat{F} whose interiors are disjoint from $\phi_1 \cup \phi_2$. The torus $A_{12} \cup E_1 \cup A_{21} \cup E_2$ bounds a solid torus V in \widehat{X}^ϵ in which A_{12} is not parallel to A_{21} . Let E_* be the annulus in \widehat{F} with $\partial E_* = \partial A_{11}$ and whose interior is disjoint from $\phi_1 \cup \phi_2$. The torus $A_{11} \cup E_*$ bounds a solid torus V_* in \widehat{X}^ϵ in which A_{11} is not parallel to E_* . Therefore $U = V_* \cup Q_{P_1} \cup V$ is a submanifold of $M(\beta)$ satisfying the hypotheses of Lemma 4.1 and as above, this lemma implies that $\partial U = E_* \cup (Q_{P_1} \cap \widehat{F}) \cup E_1 \cup E_2 \cup A_{21}$ is incompressible in $M(\beta)$. But this is impossible as $|\partial U \cap \partial M| < m$. \diamond

We can refine Proposition 7.1 somewhat in the absence of tight components of $\check{\Phi}_1^\epsilon$.

Lemma 7.4. *If $\dot{\Sigma}_1^\epsilon = \emptyset$, then $t_1^\epsilon = |\partial S|$.*

Proof. If $\dot{\Sigma}_1^\epsilon$ is empty, then so is $\check{\Phi}_1^\epsilon$ and therefore $\check{\Phi}_1^\epsilon$ is a collar on ∂S , so $t_1^\epsilon = |\partial S|$. \diamond

Proposition 7.5. *When F is separating and $t_1^\epsilon = 0$, then $\dot{\Sigma}_1^\epsilon$ has a unique component P and either $P = X^\epsilon$ or each component of $\check{\Phi}_1^\epsilon = P \cap F$ completes to an essential annulus in \widehat{F} . Further, the base orbifold of the Seifert structure on \widehat{X}^ϵ described in Proposition 7.1 has*

(1) *no cone points of order 2 if P is a product I -bundle,*

- (2) one cone point of order 2 if P is a twisted I -bundle and $\dot{\Phi}_1^\epsilon \neq F$,
- (3) two cone points of order 2 if $P = X^\epsilon$, i.e. X^ϵ is a twisted I -bundle so $\epsilon = -$.

Proof. Since $t_1^\epsilon = 0$, $\dot{\Sigma}_1^\epsilon$ has at least one component (Lemma 7.4) and each inner boundary component of $\dot{\Phi}_1^\epsilon$ is \widehat{F} -essential (Corollary 6.4). Proposition 7.1 then shows that $\dot{\Sigma}_1^\epsilon$ has exactly one component. Call it P . Proposition 7.1 also shows that either $P = X^\epsilon$ or $X^\epsilon \setminus P$ is a union of solid tori. Since the I -bundle structure on P does not extend over these solid tori, the result follows. \diamond

Corollary 7.6. *If F is separating and $t_1^+ = 0$, the base orbifold of the Seifert structure on \widehat{X}^+ described in Proposition 7.1 is $D^2(a, b)$ where $(a, b) \neq (2, 2)$. Further, $M(\beta)$ is not a union of two twisted I -bundles over the Klein bottle.*

Proof. The first assertion follows from part (3) of the previous proposition. Suppose that $M(\beta)$ is a union of two twisted I -bundles over the Klein bottle along their common boundary T . Then T is not isotopic to \widehat{F} by the first assertion. Hence as T splits $M(\beta)$ into two atoroidal Seifert manifolds, $M(\beta)$ must be Seifert. If \widehat{F} is horizontal, it splits $M(\beta)$ into two twisted I -bundles, necessarily over the Klein bottle, which contradicts Assumption 2.6. Thus it is vertical and T is horizontal. It follows that the base orbifold \mathcal{B} of $M(\beta)$ is Euclidean. Further, \mathcal{B} is non-orientable as T separates. Thus \mathcal{B} is either a Klein bottle or $P^2(2, 2)$. In either case \widehat{F} splits $M(\beta)$ into the union of two twisted I -bundles over the Klein bottle, contrary to the first assertion of the corollary. This completes the proof. \diamond

7.2. S is not connected. In this subsection we prove Proposition 7.2. It will follow from the four lemmas below.

Lemma 7.7. *When F is non-separating, $\dot{\Sigma}_1^+$ is a (possibly empty) product I -bundle.*

Proof. Suppose that $\dot{\Phi}_1^+$ has a τ_+ -invariant component, ϕ say. Then there is a Möbius band $(B, \partial B) \subseteq (X^+, \phi)$. According to Proposition 4.3, ∂B is essential in \widehat{S} . Our hypotheses imply that ϕ/τ_+ contains a once-punctured Möbius band. Its inverse image in \widehat{S} is a τ_+ -invariant twice-punctured annulus $\phi_0 \subseteq \phi$ such that $\widehat{\phi}_0$ is essential in \widehat{S} . Without loss of generality we can suppose that $\widehat{\phi}_0 \subseteq \widehat{F}_1$.

Now ϕ_0 has at least two outer boundary components and two inner ones. We denote the latter by c_1, c_2 . By construction $c_2 = \tau_+(c_1)$ and c_1 and c_2 cobound an essential annulus A in (X^+, F_1) . Note that $E = \widehat{F}_1 \setminus \widehat{\phi}_0$ is an annulus and $A \cup E$ a non-separating torus in $M(\beta)$ which intersects ∂M in fewer than m components. Hence it is compressible. But then $M(\beta)$ contains a non-separating 2-sphere, which is impossible by Assumption 2.8. Thus there is no τ_+ -invariant component of $\dot{\Phi}_1^+$. \diamond

Lemma 7.8. *Suppose F is non-separating. Let P be a component of $\dot{\Sigma}_1^+$ and let $\phi_1, \phi_2 \subseteq S$ be the two horizontal boundary components of P . If ϕ_1 contains an \widehat{S} -essential annulus, then ϕ_1 and ϕ_2 are contained in different components of S .*

Proof. Suppose that both ϕ_1 and ϕ_2 are contained in F_1 , say. Choose a neat subsurface $\phi_{1,0}$ of ϕ_1 such that $\widehat{\phi}_{1,0}$ is an \widehat{S} -essential annulus and $|\phi_{1,0} \cap \partial M| > 0$. Set $\phi_{2,0} = \tau_+(\phi_{1,0})$. Then $\widehat{\phi}_{1,0}$ and $\widehat{\phi}_{2,0}$ are disjoint essential annuli in \widehat{F}_1 . The frontier of P in X^+ is a set of two essential annuli in (X^+, F_1) , which we denote by A_1 and A_2 . According to Lemma 4.7, each A_i is separating in X^+ . For $i = 1, 2$, ∂A_i bounds an annulus E_i in \widehat{F}_1 whose interior is disjoint from $\phi_{1,0} \cup \phi_{2,0}$.

The annulus A_1 splits \widehat{X}^+ into two components, which we denote by W_1 and W_2 . We may suppose that the torus $A_1 \cup E_1$ is a boundary component of W_1 . Now $\widehat{F}_2 \subseteq \partial W_i$ for some i , and in this case $A_i \cup E_i$ is a non-separating torus in $M(\beta)$ whose intersection with ∂M has fewer than m components, contrary to Assumption 2.2. Thus the conclusion of the lemma holds. \diamond

Lemma 7.9. *If there is a component P of $\dot{\Sigma}_1^+$ and $j \in \{1, 2\}$ such that $|P \cap F_j| = 2$, then $t_1^+ \geq 4$.*

Proof. Without loss of generality, we can suppose that $j = 1$. Lemma 7.8 implies that no component ϕ of $P \cap F_1$ contains an \widehat{S} -essential annulus. Thus there is a disk in \widehat{F}_1 containing ϕ and this disk must contain a tight component of $\check{\Phi}_1^\natural$. The same is true for the other component of $P \cap F_1$, so the number of tight components of $\check{\Phi}_1^+$ contained in F_1 is at least 2. To see that the same is true for F_2 , it suffices to show that there is a component P' of $\dot{\Sigma}_1^+$ such that $|P' \cap F_2| = 2$. But it is clear that such a component exists since Lemma 7.7 implies that the number of boundary components of F_1 contained in a component of $\dot{\Sigma}_1^+$ which intersects both F_1 and F_2 equals the number of such boundary components of F_2 . \diamond

Lemma 7.10. *Suppose F is non-separating and $t_1^+ = 0$. Then there is a unique component P of $\dot{\Sigma}_1^+$ such that $\widehat{P \cap F_1}$ contains an \widehat{F}_1 -essential annulus. Further, \widehat{X}^+ admits a Seifert structure in which $\widehat{\Phi}_1^+$ is vertical and whose base orbifold is an annulus with exactly one cone point.*

Proof. First observe that $\dot{\Sigma}_1^+$ has at least one component, P say, since $t_1^+ = 0$. By Lemma 7.9, $|P \cap F_j| = 1$ for each j . Set $\phi_j = P \cap F_j$. Corollary 6.4 implies that each inner boundary component of ϕ_j is \widehat{F}_j -essential. There must be such boundary components since X^+ is not a product. Thus $\widehat{\phi}_j$ is an \widehat{F}_j -essential annulus.

Let P_1, \dots, P_k be the components of $\dot{\Sigma}_1^+$ and set $\phi_{1i} = P_i \cap F_1$ and $\phi_{2i} = P_i \cap F_2$. Then each $\widehat{\phi}_{j,i}$ is an \widehat{F}_j -essential annulus. The closure of the complement of $\cup_i \widehat{\phi}_{j,i}$ in \widehat{F}_j is a set of annuli which we denote by $E_{ji}, i = 1, \dots, k$. We may assume that $\widehat{\phi}_{j1}, E_{j1}, \widehat{\phi}_{j2}, E_{j2}, \dots, \widehat{\phi}_{jk}, E_{jk}$ appear consecutively in \widehat{F}_j .

Let $d_i = |\phi_{1i} \cap \partial M| = |\phi_{2i} \cap \partial M|$. Since $\check{\Phi}_1^+$ has no tight components, $d_1 + \dots + d_k = m$. We will assume that $k > 1$ in order to derive a contradiction. Then without loss of generality, $2d_1 \leq m$.

For each $i = 1, \dots, k$, let A_i, A'_i be the two components of the frontier of P_i in X^+ . Then each of A_i and A'_i is an essential annulus in (X^+, S) . We may assume that $\partial A'_i \cup \partial A_{i+1} = \partial E_{1i} \cup \partial E_{2i}$, so $A_1, A'_1, A_2, A'_2, \dots, A_k, A'_k$ appear consecutively in X^+ .

Now $A'_i \cup E_{1i} \cup A_{i+1} \cup E_{2i}$ is a torus in X^+ which contains a curve which is null-homotopic in M . (Here the indices are defined (mod k .) It therefore bounds a solid torus V_i in X^+ . Note that A'_i is not parallel in V to A_{i+1} as otherwise P_i and P_{i+1} would be contained in a component of $\dot{\Sigma}_1^+$. Then $U_i = V_{i-1} \cup \widehat{P}_i \cup V_i$ is a submanifold of $M(\beta)$ which is a Seifert fibred space over the disk with two cone points. Since \widehat{S} can be isotoped into $\overline{M(\beta) \setminus U_i}$, Lemma 4.1 implies that ∂U_i is an incompressible torus in $M(\beta)$. By construction, ∂U_1 contains $2d_1 \leq m$ components of ∂M . Assumption 2.2 then implies that $2d_1 = m$. But this is impossible by Assumption 2.3. Thus $k = 1$.

Finally note that the closure of the complement of $\dot{\Sigma}_1^+$ in X^+ is a solid torus V such that $\dot{\Sigma}_1^+ \cap V = A_1 \cup A'_1$, in X^+ . Hence \widehat{X}^+ is homeomorphic to the manifold obtained from V by identifying A_1 with A'_1 . It is therefore a Seifert fibred space over the annulus with at most one cone point. If there is no cone point, then F is a fibre in M , contrary to Assumption 2.6. This completes the proof. \diamond

Corollary 7.11. *If F is non-separating and $t_1^+ = 0$, then $M(\beta)$ does not fibre over the circle with torus fibre.*

Proof. Suppose otherwise and let T be the fibre. Isotope \widehat{F} so that it intersects T transversally and in a minimal number of components. Since T is a fibre, the previous lemma shows that $T \cap \widehat{F} \neq \emptyset$ and so T cuts \widehat{F} into a finite collection of incompressible annuli which run from one side of T to the other. It follows that $M(\beta)$ admits a Seifert structure in which T is horizontal. If \widehat{F} is horizontal it is a fibre in $M(\beta)$, which contradicts Assumption 2.6. Thus it is vertical. It follows that the base orbifold \mathcal{B} of $M(\beta)$ is Euclidean. Further, the projection image of \widehat{F} in \mathcal{B} is a non-separating two-sided curve. Thus \mathcal{B} is either a torus or Klein bottle. In either case \widehat{F} splits $M(\beta)$ into the product of a torus and an interval, which is impossible by Lemma 7.10. This completes the proof. \diamond

8. \widehat{S} -ESSENTIAL ANNULI IN $\dot{\Phi}_j^\epsilon$

Proposition 8.1. *Suppose that F is separating. If $\dot{\Phi}_2^+$ or $\dot{\Phi}_2^-$ contains an \widehat{F} -essential annulus, then \widehat{X}^ϵ admits a Seifert structure with base orbifold of the form $D^2(a, b)$ for some $a, b \geq 2$ for both ϵ . Further, one of the following situations arises:*

(i) $t_1^+ + t_1^- \geq 4$.

(ii) X^- is a twisted I -bundle.

(iii) $M(\beta)$ admits a Seifert structure with base orbifold $S^2(a, b, c, d)$. Further, if $t_1^\epsilon = 0$ for some ϵ , then $(a, b, c, d) \neq (2, 2, 2, 2)$.

Proof. As $h_2^+ : \dot{\Phi}_2^+ \xrightarrow{\cong} \dot{\Phi}_2^-$, we can suppose that $\dot{\Phi}_2^-$ contains an \widehat{F} -essential annulus. Since $\dot{\Phi}_2^- = \tau_-(\dot{\Phi}_1^- \wedge \dot{\Phi}_1^+)$, if $\dot{\Phi}_2^-$ contains an \widehat{F} -essential annulus, so do $\dot{\Phi}_1^+$ and $\dot{\Phi}_1^-$. Hence Proposition 7.1 implies that \widehat{X}^ϵ admits a Seifert structure with base orbifold of the form $D^2(a, b)$ for some $a, b \geq 2$ for both ϵ .

If $\text{genus}(\dot{\Phi}_1^\epsilon) = 1$ for some ϵ , then either X^- is a twisted I -bundle or $t_1^\epsilon \geq 4$ (Corollary 6.4). Thus (i) or (ii) holds. Assume that $\text{genus}(\dot{\Phi}_1^\epsilon) = 0$ for both ϵ , so X^- is not a twisted I -bundle, and let φ_ϵ be the slope on \widehat{F} of an \widehat{F} -essential annulus contained in $\dot{\Phi}_1^\epsilon$. Then φ_ϵ is the fibre slope of the Seifert structure on \widehat{X}^ϵ given by Proposition 7.1. As $\dot{\Phi}_2^- = \tau_-(\dot{\Phi}_1^- \wedge \dot{\Phi}_1^+)$, we see that $\dot{\Phi}_1^-$ contains curves of slope φ_+ and φ_- . Hence if these slopes are distinct, $\text{genus}(\dot{\Phi}_1^-) = 1$, contrary to our assumptions. Thus $\varphi_+ = \varphi_-$ so $M(\beta)$ admits a Seifert structure with base orbifold of the form $S^2(a, b, c, d)$. Finally if $t_1^\epsilon = 0$ for some ϵ , Proposition 7.5 shows that $(a, b, c, d) \neq (2, 2, 2, 2)$. \diamond

Proposition 8.2. *Suppose that F is separating. If $\dot{\Phi}_3^+$ contains an \widehat{F} -essential annulus then either*

(i) $t_1^+ \geq 4$, or

(ii) X^- is a twisted I -bundle and $M(\beta)$ is Seifert with base orbifold $P^2(2, n)$ for some $n > 2$. Further, $t_1^+ = 0$, $\dot{\Phi}_1^+$ is an \widehat{F} -essential annulus, $\dot{\Phi}_3^+$ is the union of two \widehat{F} -essential annuli, and there are disjoint, non-separating annuli A_1^-, A_2^- properly embedded in X^- such that $\partial A_1^- \cup \partial A_2^- \subseteq \dot{\Phi}_1^+$ and for each j , $\partial \dot{\Phi}_1^+ \cap \partial A_j^-$ is a boundary component of $\dot{\Phi}_1^+$.

Proof. Assume that $t_1^+ \leq 2$. We will show that (ii) holds.

Suppose that some component ϕ_0 of $\dot{\Phi}_3^+$ contains an \widehat{F} -essential annulus and let ψ_0 be the component of $\dot{\Phi}_1^+$ containing ϕ_0 . By Assumption 2.6, $\psi_0 \neq F$. Corollary 6.4 then shows that $\text{genus}(\psi_0) = 0$ and $\widehat{\psi}_0$ completes to an \widehat{F} -essential annulus.

We can suppose that $\phi_0 \subseteq \text{int}(\psi_0)$. Set $\phi_1 = \tau_+(\phi_0) \subseteq \dot{\Phi}_1^+ \wedge \dot{\Phi}_2^-$. Now $h_3^+ = \tau_+ \circ \tau_- \circ \tau_+ | \dot{\Phi}_3^+$ is a free involution of $\dot{\Phi}_3^+$. In particular, either $h_3^+(\phi_0) = \phi_0$ or $h_3^+(\phi_0) \cap \phi_0 = \emptyset$. Equivalently, either $\tau_-(\phi_1) = \phi_1$ or $\tau_-(\phi_1) \cap \phi_1 = \emptyset$. In the first case there are an essential annulus A^- properly embedded in (X^-, ϕ_1) such that $\partial A^- = \partial \widehat{\phi}_1$ and a Möbius band B properly embedded in $(X^-, \text{int}(\phi_1))$. Proposition 5.3 then implies that ψ_0 is τ_+ -invariant. Hence there is an annulus A^+ properly embedded in (X^+, ψ_0) with $\partial A^+ = \partial \widehat{\psi}_0$. Lemma 5.1 implies that ϕ_1 , and therefore ϕ_0 , has no outer boundary components, which is impossible.

Next suppose that $\tau_-(\phi_1) \cap \phi_1 = \emptyset$. Then there is an embedding $(\phi_1 \times I, \phi_1 \times \{0\}, \phi_1 \times \{1\}) \rightarrow (X^-, \phi_1, \tau_-(\phi_1))$. First suppose that the components A_1^-, A_2^- of the image of $\partial \widehat{\phi}_1 \times I$ are separating annuli in X^- . Let A^+ be a properly embedded annulus in (X^+, ψ_0) such that at least one boundary component of A^+ is contained in $\partial \widehat{\psi}_0$. According to Lemma 5.1(1), ∂A^+ does not separate ∂A_j^- for $j = 1, 2$. Lemma 5.1(2) then implies that ϕ_1 has no outer boundary components, which is impossible. Thus A_1^- and A_2^- are non-separating in X^- . In particular, X^- is a twisted I -bundle (Lemma 4.7).

Let P be the unique component of $\dot{\Sigma}_1^+$ whose intersection with F contains an \widehat{F} -essential annulus (Proposition 7.1). Then ψ_0 is a component of $P \cap F$ and $\phi_0 \cup \phi_1 \subseteq P \cap F$. If P is a product I -bundle, let A_1^+, A_2^+ be the annuli in its frontier in X^+ , and consider the torus T obtained from the union of $A_1^+, A_1^-, A_2^+, A_2^-$ and four annuli in \widehat{F} disjoint from $\text{int}(\phi_1) \cup \text{int}(\tau_-(\phi_1))$. The reader will verify that T bounds a twisted I -bundle over the Klein bottle in $M(\beta)$ and so is

essential in $M(\beta)$ by Lemma 4.1. Hence it intersects ∂M in at least m components. But this implies $|\phi_1 \cap \partial F| = 0$, which is impossible. Thus P is a twisted I -bundle. It follows from Proposition 7.1 that \widehat{X}^+ is Seifert with base orbifold $D^2(2, n)$ with ∂A_1^- vertical. Thus $M(\beta)$ is Seifert with base orbifold $P^2(2, n)$. Let A^+ be the frontier of P in X^+ . By construction, ∂A^+ does not separate ∂A_1^- or ∂A_2^- in \widehat{F} . Lemma 5.4 then shows that $|(\widehat{F} \setminus \psi_0) \cap \partial F| = 0$. Hence, $t_1^+ = 0$. This implies that $n > 2$ (Corollary 7.6) and τ_- is defined on $\widehat{F} \setminus \psi_0$ and sends it into the interior of $\dot{\Phi}_1^+$. Thus, there are disjoint, non-separating annuli E_1^-, E_2^- properly embedded in X^- such that $\partial E_1^- \cup \partial E_2^- \subseteq \dot{\Phi}_1^+$ and for each j , $\partial \dot{\Phi}_1^+ \cap \partial E_j^-$ is a boundary component of $\dot{\Phi}_1^+$. Write $\partial E_j^- = c_j \cup c'_j$ where $\partial \dot{\Phi}_1^+ = c_1 \cup c_2$. Since $\widehat{F} \setminus \psi_0$ is an annulus, it follows from our constructions that the disjoint subsurfaces of $\dot{\Phi}_1^+$ with inner boundaries $c_1 \cup c'_2$ and $c_2 \cup c'_1$ lie in $\dot{\Phi}_3^+$ and contain ∂F . Thus their union is $\dot{\Phi}_3^+$. This proves the proposition. \diamond

Definition 8.3. ([BGZ1, page 266]) Given a closed, essential surface G in M , we let $\mathcal{C}(G)$ denote the set of slopes δ on ∂M such that S compresses in $M(\delta)$. A slope η on ∂M is called a *singular slope* for G if $\eta \in \mathcal{C}(G)$ and $\Delta(\delta, \eta) \leq 1$ for each $\delta \in \mathcal{C}(G)$.

A fundamental result of Wu [Wu1] states that if $\mathcal{C}(G) \neq \emptyset$, then there is at least one singular slope for G .

Proposition 8.4. *Let η and δ be slopes on the boundary of a hyperbolic knot manifold M .*

- (1) ([BGZ1, Theorem 1.5]) *If η is a singular slope for some closed essential surface in M and $M(\delta)$ is not hyperbolic, then $\Delta(\delta, \eta) \leq 3$.*
- (2) ([BGZ1, Theorem 1.7]) *If $M(\eta)$ is a Seifert fibred manifold whose base orbifold is hyperbolic but a 2-sphere with three cone points, then η is a singular slope for some closed essential surface in M .* \diamond

Proposition 8.5. *Suppose that F is non-separating. If $\dot{\Phi}_3^+$ contains an \widehat{S} -essential annulus and $t_1^+ = 0$, then $M(\beta)$ is Seifert fibred with base orbifold a torus or a Klein bottle with exactly one cone point. In particular, β is a singular slope for a closed essential surface in M and thus, $\Delta(\alpha, \beta) \leq 3$.*

Proof. Since $t_1^+ = 0$, Proposition 7.2 implies that $\dot{\Phi}_1^+ = \phi_1 \cup \phi_2$ where ϕ_1, ϕ_2 lie in different components of S and complete to \widehat{S} -essential annuli. This proposition also implies that \widehat{X}^+ admits a Seifert fibred structure with base orbifold an annulus with one cone point. Further, $\widehat{\phi}_j$ is vertical in this structure for both j . To see that $M(\beta)$ is Seifert with base orbifold as claimed, it suffices to show that the slope of $\widehat{\tau}_-(\phi_j)$ coincides with that of $\widehat{\phi}_{3-j}$. But this is an immediate consequence of the fact that $\tau_+(\dot{\Phi}_3^+) = \dot{\Phi}_1^+ \wedge \dot{\Phi}_2^- = \dot{\Phi}_1^+ \wedge \tau_-(\dot{\Phi}_1^+)$ contains an \widehat{S} -essential annulus. \diamond

9. THE EXISTENCE OF TIGHT COMPONENTS IN $\check{\Phi}_j^\epsilon$ FOR SMALL VALUES OF j

In this section we examine the existence of tight components in $\check{\Phi}_j^\epsilon$ for small values of j . Note that if $t_1^\epsilon \neq 0$ for some ϵ , then Proposition 6.3 implies that $t_2^{-\epsilon} = t_2^\epsilon \geq t_1^\epsilon > 0$. Thus we examine

the case $t_1^+ = t_1^- = 0$. Recall that under this hypothesis, $\dot{\Phi}_1^+$ and $\dot{\Phi}_1^-$ are non-empty (Lemma 7.4).

Lemma 9.1. *Suppose that $t_1^+ = t_1^- = 0$. If $\Delta(\alpha, \beta) > 3$ and $M(\beta)$ is Seifert fibred, then its base orbifold is of the form $P^2(a, b)$ for some $(a, b) \neq (2, 2)$ and X^- is a twisted I -bundle.*

Proof. Since $\Delta(\alpha, \beta) > 3$, β is not a singular slope of a closed essential surface in M ([BGZ1, Theorem 1.5]). Hence, as $M(\beta)$ is toroidal, Seifert but not the union of two twisted I -bundles over the Klein bottle (Corollary 7.6), its base orbifold is of the form $P^2(a, b)$ ([BGZ1, Theorem 1.7]) where $(a, b) \neq (2, 2)$. Each essential torus in $M(\beta)$ splits it into the union of a twisted I -bundle over the Klein bottle and a Seifert manifold with base orbifold $D^2(a, b)$. Since $t_1^+ = t_1^- = 0$, Proposition 7.5(3) implies that X^- is a twisted I -bundle. \diamond

Lemma 9.2. *Let S_1, S_2 be large, neat, connected surfaces contained in the same component of S . Suppose, for each j , that either S_j is tight or \widehat{S}_j is an \widehat{S} -essential annulus.*

(1) *Each component of $S_1 \wedge S_2$ is either tight or an \widehat{S} -essential annulus.*

(2) *If we further assume that when both \widehat{S}_1 and \widehat{S}_2 are \widehat{S} -essential annuli, their slopes are distinct, then each component of $S_1 \wedge S_2$ is tight.*

Proof. Let S_0 be a component of $S_1 \wedge S_2$.

First suppose that S_0 is contained in a disk D in \widehat{S} . If S_0 is not tight, it has at least two inner boundary components. Let C be an inner boundary component of S_0 which is innermost in D amongst all the other inner boundary components of S_0 . Let $D_0 \subseteq D \subseteq \widehat{S}$ be the disk with boundary C . By construction, $D_0 \cap S_0 = C$. Further, the neatness of S_1 and S_2 implies that $D_0 \cap S$ is large. Since \widehat{S}_j is either a disk or an \widehat{S} -essential annulus, $D_0 \cap S \subseteq S_j$ for each j . Hence it is contained in $S_1 \wedge S_2$ and therefore S_0 , contrary to our construction. Thus S_0 must be tight. In particular, this proves (2).

Next suppose that S_0 contains an \widehat{S} -essential annulus but S_0 is not itself an \widehat{S} -essential annulus. Then S_0 has at least three inner boundary components and all but exactly two of them are inessential in \widehat{S} . Fix an inessential inner boundary component C of S_0 . The argument of the previous paragraph is easily adapted to this case and leads to a contradiction. Thus \widehat{S}_0 must be an \widehat{S} -essential annulus. \diamond

Proposition 9.3. *If $t_1^+ = t_1^- = 0$, then one of the following three scenarios arises.*

(i) $\check{\Phi}_3^+$ is a union of tight components.

(ii) $t_3^+ = 0$, X^- is a twisted I -bundle, and $M(\beta)$ is Seifert with base orbifold $P^2(2, n)$ for some $n > 2$. Further, $\dot{\Phi}_1^+$ completes to an \widehat{F} -essential annulus, $\dot{\Phi}_3^+$ completes to the union of two \widehat{F} -essential annuli, and there are disjoint, non-separating annuli A_1^-, A_2^- properly embedded in (X^-, F) such that $\partial A_1^- \cup \partial A_2^- \subseteq \dot{\Phi}_1^+$ and for each j , $\partial \dot{\Phi}_1^+ \cap \partial A_j^-$ is a boundary component of $\dot{\Phi}_1^+$.

(iii) X^- is a product I -bundle and $M(\beta)$ is Seifert fibred with base orbifold a torus or a Klein bottle with exactly one cone point. In particular, β is a singular slope for a closed essential surface in M and thus, $\Delta(\alpha, \beta) \leq 3$.

Proof. Propositions 7.1 and 7.2 imply that $\check{\Phi}_1^\epsilon = \dot{\Phi}_1^\epsilon$ is either S or a union of subsurfaces whose completions are \widehat{S} -essential annuli. If no component of $\check{\Phi}_3^+$ contains an \widehat{S} -essential annulus, Lemma 9.2 shows that (i) holds. If some component of $\check{\Phi}_3^+$ does contain an \widehat{S} -essential annulus, Propositions 8.2 and 8.5 show that (ii) and (iii) hold. \diamond

Proposition 9.4. *Suppose that $t_1^+ = t_1^- = 0$ and $\Delta(\alpha, \beta) > 3$.*

- (1) *If X^- is not an I -bundle, each component of $\check{\Phi}_j^\epsilon$ is tight for all $j \geq 2$ and both ϵ .*
- (2) *If X^- is a product I -bundle, or X^- is a twisted I -bundle and $\check{\Phi}_3^+$ does not contain an \widehat{S} -essential annulus, each component of $\check{\Phi}_j^+$ is tight for all $j \geq 3$.*
- (3) *If X^- is a twisted I -bundle and $\check{\Phi}_3^+$ contains an \widehat{S} -essential annulus, then $t_3^+ = 0$, $M(\beta)$ is Seifert with base orbifold $P^2(2, n)$ for some $n > 2$, $\dot{\Phi}_1^+$ and $\dot{\Phi}_3^+$ are as described in Proposition 9.3(ii), and each component of $\check{\Phi}_j^+$ is tight for all $j \geq 5$.*

Proof. Propositions 7.1 and 7.2 show that for each ϵ , $\check{\Phi}_1^\epsilon = \dot{\Phi}_1^\epsilon$ is either S or a union of subsurfaces whose completions are \widehat{S} -essential annuli.

First note that in order to prove assertion (1), it suffices to show that each component of $\dot{\Phi}_2^+$ is tight. For if this holds, the same is true of $\dot{\Phi}_2^- = h_2^+(\dot{\Phi}_2^+)$. Suppose inductively that each component of $\dot{\Phi}_j^\epsilon$ is tight for some $j \geq 2$ and both ϵ . Lemma 9.2 combines with the identity $\dot{\Phi}_{j+1}^\epsilon = \tau_\epsilon(\dot{\Phi}_1^\epsilon \wedge \dot{\Phi}_j^{-\epsilon})$ to show that each component of $\dot{\Phi}_{j+1}^\epsilon$ is tight.

Consider $\dot{\Phi}_2^+$ then. Since $t_1^+ = t_1^- = 0$ and X^- is not an I -bundle, $S = F$ is separating. Proposition 7.1 implies that for each ϵ and component ϕ of $\dot{\Phi}_1^\epsilon$, $\widehat{\phi}$ is an \widehat{F} -essential annulus. Lemma 9.1 implies that $M(\beta)$ is not Seifert, so the slopes of an \widehat{F} -essential annulus in $\widehat{\Phi}_1^+$ and an \widehat{F} -essential annulus in $\widehat{\Phi}_1^-$ are distinct. Hence Lemma 9.2 implies that each component of $\dot{\Phi}_2^+ = \tau_+(\dot{\Phi}_1^+ \wedge \dot{\Phi}_1^-)$ is tight.

Next consider the hypotheses of assertion (2). Proposition 9.3 implies that each component of $\dot{\Phi}_3^+$ is tight. Since X^- is an I -bundle, $\dot{\Phi}_3^+ = \dot{\Phi}_4^+$, so the lemma holds when $j = 3, 4$. But when $j \geq 5$, we have $\dot{\Phi}_{j+1}^+ = \tau_\epsilon(\dot{\Phi}_1^\epsilon \wedge \tau_-(\dot{\Phi}_{j-1}^+))$, so Lemma 9.2 combines with an inductive argument to show that (2) holds for all $j \geq 3$.

Finally consider assertion (3). Now $\tau_-\tau_+(\check{\Phi}_5^+) = \check{\Phi}_3^+ \wedge \check{\Phi}_2^- = \check{\Phi}_3^+ \wedge \tau_-(\check{\Phi}_1^+) = \check{\Phi}_3^+ \wedge \tau_+(\check{\Phi}_3^+)$ where the latter identity follows from Proposition 9.3(ii). Now $\tau_+(\check{\Phi}_3^+) = \phi_1 \cup \phi_2$ where $\widehat{\phi}_j$ is an \widehat{F} -essential annulus and $\phi_2 = \tau_-(\phi_1)$. Hence

$$\check{\Phi}_3^+ \wedge \tau_+(\check{\Phi}_3^+) = \phi_1 \wedge \tau_+(\phi_1) \sqcup \phi_1 \wedge \tau_+(\phi_2) \sqcup \phi_2 \wedge \tau_+(\phi_1) \sqcup \phi_2 \wedge \tau_+(\phi_2)$$

By construction, $\phi_j \cap \tau_+(\phi_{3-j})$ contains an inner boundary component of $\dot{\Phi}_1^+$ for both j . If $\phi_j \wedge \tau_+(\phi_{3-j}) = \phi_j$ for some j , then $\phi_j \wedge \tau_+(\phi_{3-j}) = \phi_j$ and $\phi_j \wedge \tau_+(\phi_j) = \emptyset$ for both j .

Hence $\tau_- \tau_+(\check{\Phi}_5^+) = \check{\Phi}_3^+$ from which it follows that $\check{\Phi}_3^+ = \check{\Phi}_5^+$, which is impossible. Hence $\phi_j \wedge \tau_+(\phi_{3-j}) \neq \phi_j$ for both j . It follows that $\phi_j \wedge \tau_+(\phi_j) \neq \emptyset$ is a non-empty union of tight components for each j .

Next consider $\phi_j \wedge \tau_+(\phi_{3-j})$. By Lemma 9.2, each of its components is either tight or completes to an \widehat{S} -essential annulus. Since $\tau_+(\phi_{3-j})$ contains the inner component c_j of $\widehat{\Phi}_1^+$ contained in ϕ_j , there is exactly one component of $\phi_j \wedge \tau_+(\phi_{3-j})$, E_j say, which completes to an \widehat{S} -essential annulus. To complete the proof we need only show that $E_j \cap \partial S = \emptyset$.

By construction, $\tau_+(E_1) = E_2$ and $\tau_+(E_2) = E_1$. Let $E_0 = \overline{\widehat{\Phi}_1^+ \setminus (E_1 \cup E_2)}$. Then E_0 completes to an \widehat{F} -essential annulus \widehat{E}_0 which is invariant under τ_+ . Hence the associated I -bundle over \widehat{E}_0 is a solid torus V_1 whose frontier in \widehat{X}^+ is an essential annulus in X^+ which has winding number 2 in V_1 .

Next consider the solid torus $V_2 = \overline{X^+ \setminus \widehat{\Sigma}_1^+}$. The frontier of V_2 in X^+ is an essential annulus in X^+ cobounded by c_1 and c_2 . Let A be the other annulus in ∂V_2 cobounded by c_1 and c_2 . Then $\tau_-(A) = \overline{\widehat{\Phi}_1^+ \setminus (\phi_1 \cup \phi_2)}$ is a core annulus in \widehat{E}_0 . The I -bundle in X^- over A is a solid torus in which A has winding number 1. It follows that $W = V_1 \cup V_2 \cup V_3 \subset M(\beta)$ is a Seifert fibered space over a disk with two cone points. In particular, ∂W is incompressible in W . The exterior of W in $M(\beta)$ is a Seifert fibered space over a Möbius band so ∂W is also incompressible in $M(\beta)$. If $E_j \cap \partial S \neq \emptyset$ for some j then ∂W intersects ∂M in fewer than m components contrary to Assumption 2.2. Thus we must have $E_j \cap \partial S = \emptyset$ for both j , which completes that proof of (3). \diamond

10. LENGTHS OF ESSENTIAL HOMOTOPIES

It is clear that $\chi(\check{\Phi}_j^\epsilon) = 0$ if and only if $\chi(\check{\Phi}_j^\epsilon)$ is a regular neighbourhood of ∂S . Thus if we set

$$l_\epsilon = \max\{j : \chi(\check{\Phi}_j^\epsilon) \neq 0\}$$

then l_ϵ is the maximal length of an essential homotopy in (M, S) of a large function which begins on the ϵ -side of S . Hence

$$l_S = \max\{l_+, l_-\}$$

is the maximal length of an essential homotopy in (M, S) of a large function. It is evident that $|l_+ - l_-| \leq 1$ and therefore $|l_\epsilon - l_S| \leq 1$ for each ϵ .

Proposition 10.1. *Suppose that $\Delta(\alpha, \beta) > 3$ if F is non-separating.*

(1) $l_+ \leq |\partial S| - t_1^+$ if $t_3^+ > 0$ and $l_+ \leq |\partial S| - t_1^+ + 2$ otherwise. Hence,

$$l_S \leq \begin{cases} |\partial S| - t_1^+ + 1 & \text{if } t_3^+ > 0 \\ |\partial S| - t_1^+ + 3 & \text{otherwise} \end{cases}$$

(2) If X^- is not an I -bundle, then $l_- \leq |\partial S| - t_1^-$. Hence,

$$l_S \leq |\partial S| - \begin{cases} \max\{t_1^+, t_1^-\} & \text{if } t_1^+ = t_1^- \\ \max\{t_1^+, t_1^-\} - 1 & \text{if } t_1^+ \neq t_1^- \end{cases}$$

Proof. For each ϵ we know that $\mathcal{T}_{l_{\epsilon+1}}^\epsilon$ is a regular neighbourhood of ∂S while $\mathcal{T}_{l_\epsilon}^\epsilon$ isn't, so $t_{l_\epsilon} < t_{l_{\epsilon+1}} = |\partial S|$. If $t_3^+ > 0$, Proposition 6.3 implies that if $2k+1 \leq l_+$, then $t_1^+ < t_3^+ < \dots < t_{2k+1}^+ < |\partial S|$. As each of these numbers is even, $|\partial S| > t_{2k+1}^+ \geq 2k+t_1^+$. Hence $2k+2 \leq |\partial S| - t_1^+$. It follows that $l_+ \leq |\partial S| - t_1^+$ and therefore $l_S \leq |\partial S| - t_1^+ + 1$. In general, Proposition 9.4 implies that $t_5^+ > 0$, which yields $l_+ \leq |\partial S| - t_1^+ + 2$. Thus assertion (1) holds.

If X^- is not an I -bundle, then Proposition 9.4 implies that $t_3^- > 0$, so the argument of the previous paragraph shows that $l_- \leq |\partial S| - t_1^-$ and therefore

- $l_S \leq |\partial S| - t_1^- + 1$; and
- $l_S = \max\{l_+, l_-\} \leq \max\{|\partial S| - t_1^+, |\partial S| - t_1^-\} = |\partial S| - \min\{t_1^+, t_1^-\}$.

Part (1) of the proposition combines with the first inequality to show that $l_S \leq \min\{|\partial S| - t_1^+ + 1, |\partial S| - t_1^- + 1\} = |\partial S| - \max\{t_1^+, t_1^-\} + 1$. The latter combines with the second inequality to yield the upper bound for l_S described in (2). \diamond

Proposition 9.4 and Proposition 10.1 imply the following corollary.

Corollary 10.2. *Suppose that $\Delta(\alpha, \beta) > 3$ if F is non-separating. Then*

$$l_S \leq \begin{cases} |\partial S| & \text{if } X^- \text{ is not an } I\text{-bundle} \\ |\partial S| + 1 & \text{if } X^- \text{ is an } I\text{-bundle and } \hat{\Phi}_3^+ \text{ contains no } \hat{S}\text{-essential annulus when it is twisted} \\ |\partial S| + 3 & \text{if } X^- \text{ is a twisted } I\text{-bundle and } \hat{\Phi}_3^+ \text{ contains an } \hat{S}\text{-essential annulus} \end{cases} \diamond$$

11. THE INTERSECTION GRAPH OF AN IMMERSSED DISK OR TORUS

We recall some of the set up from [BCSZ2, §12].

A 3-manifold is *very small* if its fundamental group does not contain a non-abelian free group.

By Assumption 2.1, $M(\alpha)$ is a small Seifert manifold with base orbifold $S^2(a, b, c)$ where $a, b, c \geq 1$. It is well-known that $M(\alpha)$ is very small if and only if $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \geq 1$. In this case, the non-abelian free group $\pi_1(F)$ cannot inject into $\pi_1(M(\alpha))$. Hence for either ϵ we can find maps $h : D^2 \rightarrow M(\alpha)$ such that the loop $h(\partial D^2)$ is contained in $X^\epsilon \setminus F$ and represents a non-trivial element of $\pi_1(X^\epsilon)$. This won't necessarily be possible when $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} < 1$ since $M(\alpha)$ is not very small. Nevertheless, the inverse image in $M(\alpha)$ of an essential immersed loop contained in the exterior of the cone points of $S^2(a, b, c)$ will be an essential immersed torus in $M(\alpha)$. Hence we can find π_1 -injective immersions $h : T \rightarrow M(\alpha)$ where T is a torus.

Let V_α be the filling solid torus used in forming $M(\alpha)$. It is shown in [BCSZ2, §12] that we can choose an immersion $h : Y \rightarrow M(\alpha)$, where Y is a disk D if $M(\alpha)$ is very small or a torus T if $M(\alpha)$ otherwise, such that

- (1) When Y is a disk D , $h(\partial D) \subseteq M \setminus F \subseteq M \subseteq M(\alpha)$;
- (2) $h^{-1}(V_\alpha)$ is a non-empty set of embedded disks in the interior of Y and h is an embedding when restricted on $h^{-1}(V_\alpha)$;
- (3) $h^{-1}(F)$ is a set of arcs or circles properly embedded in the punctured surface $Y_0 = Y \setminus \text{int}(h^{-1}(V_\alpha))$;
- (4) If e is an arc component of $h^{-1}(F)$, then $h| : e \rightarrow F$ is an essential (immersed) arc;
- (5) If c is a circle component of $h^{-1}(F)$, then $h| : c \rightarrow F$ is an essential (immersed) 1-sphere.

For any subset s of Y , we use s^* to denote its image under the map h . Denote the components of $\partial(h^{-1}(V_\alpha))$ by a_1, \dots, a_n so that a_1^*, \dots, a_n^* appear consecutively on ∂M . Note again that $h| : a_i \rightarrow a_i^* \subseteq \partial M$ is an embedding and that a_i^* has slope α in ∂M , for each $i = 1, \dots, n$. We fix an orientation on Y_0 and let each component a_i of ∂Y_0 have the induced orientation. Two components a_i and a_j are said to have the same orientation if a_i^* and a_j^* are homologous in ∂M . Otherwise, they are said to have different orientations.

Denote the components of ∂F by b_1, \dots, b_m so that they appear consecutively in ∂M . Similar definitions apply to the components of ∂F . Since Y_0 , F and M are all orientable, one has the following

Parity rule: An arc component e of $h^{-1}(F)$ in Y_0 connects components of ∂Y_0 with the same orientation (respectively opposite orientations) if and only if the corresponding e^* in F connects components of ∂F with opposite orientations (respectively the same orientation).

We define an *intersection graph* Γ_F on the surface Y by taking $h^{-1}(V_\alpha)$ as (fat) vertices and taking arc components of $h^{-1}(F)$ as edges. Note that Γ_F has no trivial loops, i.e. no 1-edge disk faces. Also note that we can assume that each a_i^* intersects each component b_j in ∂M in exactly $\Delta(\alpha, \beta)$ points. If e is an edge in Γ_F with an endpoint at the vertex a_i , then the corresponding endpoint of e^* is in $a_i^* \cap b_j$ for some b_j , and the endpoint of e is thus given the label j . So when we travel around a_i in some direction, we see the labels of the endpoints of edges appearing in the order $1, \dots, m, \dots, 1, \dots, m$ (repeated $\Delta(\alpha, \beta)$ times). It also follows that each vertex of Γ_F has valency $m\Delta(\alpha, \beta)$.

Define the *double* of Γ_F to be the graph $D(\Gamma_F)$ in Y as follows: the vertices of $D(\Gamma_F)$ are the vertices of Γ_F ; the edges of $D(\Gamma_F)$ are obtained by doubling the edges of Γ_F (i.e. each edge e is replaced by two parallel copies of e). Finally we set

$$\Gamma_S = \begin{cases} \Gamma_F & \text{if } F \text{ separates} \\ D(\Gamma_F) & \text{if } F \text{ does not separate} \end{cases}$$

It is clear that

- (1) Γ_S is a graph in Y determined by the intersection of an immersed disk or torus with S .
- (2) each vertex of Γ_S has valency $|\partial S|\Delta(\alpha, \beta)$.
- (3) if two faces of Γ_S share a common edge, then they lie on different sides of S .

- (4) if F does not separate, then a face of Γ_S which is sent by h into X^- is a bigon bounded by parallel edges.

Suppose that e and e' are two adjacent parallel edges of Γ_S . Let R be the bigon face between them, realizing the parallelism. Then $(R, e \cup e')$ is mapped into (X^ϵ, S) by the map h for some ϵ . Moreover $h|_R$ provides a basic essential homotopy between the essential paths $h|_e$ and $h|_{e'}$. We may and shall assume that $R^* = h(R)$ is contained in the characteristic I -bundle pair $(\dot{\Sigma}_1^\epsilon, \dot{\Phi}_1^\epsilon)$ of (X^ϵ, S) . We may consider R as $e \times I$ and assume that the map $h : R \rightarrow \dot{\Sigma}_1^\epsilon$ is I -fibre preserving.

Let $\bar{\Gamma}_S$ be the *reduced graph* of Γ_S obtained from Γ_S by amalgamating each maximal family of parallel edges into a single edge. It is evident that $\bar{\Gamma}_S$ coincides with the similarly defined graph $\bar{\Gamma}_F$. The following lemma is a simple consequence of the construction of Γ_S .

Lemma 11.1.

- (1) *There is at most one 1-edge face of $\bar{\Gamma}_S$ and if one, it is a collar on ∂Y when Y is a disk and a once-punctured torus when Y is a torus.*
- (2) *A 2-edge face of $\bar{\Gamma}_S$ is either*
- (a) *a collar on ∂Y bounded by a circuit of two edges and two vertices when Y is a disk;*
 - (b) *a once-punctured torus bounded by a circuit of two edges and two vertices;*
 - (c) *an annulus cobounded by two circuits, each with one edge and one vertex;*
 - (d) *a twice-punctured torus bounded by a circuit of one edge and one vertex. \diamond*

The *weight* of an edge \bar{e} of $\bar{\Gamma}_S$ is the number of parallel edges in Γ_S that \bar{e} represents.

Call the vertex of Γ_S (or $\bar{\Gamma}_S$) with boundary a_i *positive* if a_i and a_1 are like-oriented on ∂M . Otherwise call it *negative*.

Call an edge e , respectively \bar{e} , of Γ_S , respectively $\bar{\Gamma}_S$, *positive* if it connects two positive vertices or two negative vertices. Otherwise it is said to be *negative*.

Proposition 11.2. *If Y is a torus, the number of positive vertices of Γ_S equals the number of negative vertices.*

Proof. Up to taking absolute value, the difference between the number of positive vertices and the number of negative vertices is the intersection number between a class in $H_1(M(\alpha))$ carried by the core of the α -filling torus and $h_*([Y]) \in H_2(M(\alpha))$. Thus the lemma holds as long as $H_2(M(\alpha)) = 0$. Suppose then that $H_2(M(\alpha)) \neq 0$. Since $M(\alpha)$ is small Seifert, we have $H_2(M(\alpha)) \cong \mathbb{Z}$ and is generated by an embedded horizontal surface, G say, which is a fibre in a locally trivial surface bundle $M(\alpha) \rightarrow S^1$. Thus $h_*([Y])$ is a non-zero multiple of $[G]$. In particular, the Thurston norm of $[G]$ is zero. Hence G is a torus (cf. Assumption 2.8). Thus $M(\alpha)$ is toroidal small Seifert. But then $M(\alpha)$ is very small contrary to our assumption that Y is a torus. This completes the proof. \diamond

To each orientation of an edge \bar{e} of $\bar{\Gamma}_S$ of weight $|\partial S|$ or more we can associate a permutation σ of the labels as follows: if e is an edge of Γ_S in the \bar{e} -family and j is the label of its tail, then $\sigma(j)$ is the label of its head. The parity rules implies that there is an integer k such that

$$\sigma(j) \equiv \begin{cases} j + 2k \pmod{m} & \text{if } e \text{ is negative} \\ -j + 2k + 1 \pmod{m} & \text{if } e \text{ is positive} \end{cases}$$

We say that a face of Γ_S or $\bar{\Gamma}_S$ *lies on the ϵ -side of F* if it is mapped to X^ϵ by h .

The discussion in [BCSZ1, §3.4] implies the conclusion of the following proposition.

Proposition 11.3. *If $\bar{\Gamma}_S$ has an edge of weight k , then there is an essential homotopy in (M, S) of length $k - 1$ of a large map with image in S . \diamond*

This result combines with Corollary 10.2 to yield the following corollary.

Corollary 11.4. *Suppose that $\Delta(\alpha, \beta) > 3$.*

- (1) *If X^- is not an I -bundle, the weight of an edge of $\bar{\Gamma}_S$ is at most $|\partial S| + 1$.*
- (2) *If X^- is a product I -bundle, or a twisted I -bundle and Φ_3^+ does not contain an \widehat{F} -essential annulus, the weight of an edge of $\bar{\Gamma}_S$ is at most $|\partial S| + 2$.*
- (3) *If X^- is a twisted I -bundle and Φ_3^+ contains an \widehat{F} -essential annulus, the weight of an edge of $\bar{\Gamma}_S$ is at most $|\partial S| + 4$. \diamond*

We call a graph *hexagonal* if it is contained in a torus, each vertex has valency 6, and each face is a triangle. We call it *rectangular* if it is contained in a torus, each vertex has valency 4, and each face is a rectangle. Such graphs are connected.

The following proposition follows from simple Euler characteristic calculations.

Proposition 11.5.

- (1) *If each vertex of $\bar{\Gamma}_S$ has valency 6 or more, then it is hexagonal, so Y is a torus. Moreover there is a vertex of $\bar{\Gamma}_S$ incident to at least two positive edges.*
- (2) *If $\bar{\Gamma}_S$ has no triangle faces, it has a vertex of valency at most 4. If it has no vertices of valency less than 4, then it is rectangular, so Y is a torus. \diamond*

Proof. We have

$$0 \leq \chi(Y) = \sum_{\text{faces } f \text{ of } \bar{\Gamma}_S} \left\{ \chi(f) - \sum_{v \in \partial f} \left(\frac{1}{2} - \frac{1}{\text{valency}_{\bar{\Gamma}_S}(v)} \right) \right\}$$

Set $\chi_f = \chi(f) - \sum_{v \in \partial f} \left(\frac{1}{2} - \frac{1}{\text{valency}_{\bar{\Gamma}_S}(v)} \right)$. Lemma 11.1 implies that $\chi_f \leq 0$ for each monogon and bigon f in $\bar{\Gamma}_S$. The hypotheses of assertions (1) and (2) of the lemma imply that $\chi_f \leq 0$ for faces with three or more sides. Thus under either set of hypotheses, $\chi(Y) \leq 0$, so Y is a torus. Then $\chi(Y) = 0$ so $\chi_f = 0$ for all faces f . It follows that $\bar{\Gamma}_S$ is hexagonal under the

conditions of (1) and rectangular under those of (2). Finally, it is easy to check that when $\bar{\Gamma}_S$ is hexagonal, it has a vertex incident to at least two positive edges. \diamond

Lemma 11.6. *Suppose that F is non-separating and each component of $\check{\Sigma}_1^+$ intersects both F_1 and F_2 . Then every edge of $\bar{\Gamma}_S$ is negative. Hence every face of $\bar{\Gamma}_S$ has an even number of edges. In particular this is true if $t_1^+ \leq 2$.*

Proof. If each component of $\check{\Sigma}_1^+$ intersects both F_1 and F_2 , all the boundary components b_1, \dots, b_m of F have the same orientation. Hence by the parity rule, every edge of $\bar{\Gamma}_S$ is negative. The second assertion follows from the first, while the third is a consequence of Lemma 7.9 and the others. \diamond

A disk face of k -edges in the graph Γ_S is called a *Scharlemann k -gon with label pair $\{j, j+1\}$* if each edge of the face is positive with the fixed label pair $\{j, j+1\}$ at its two endpoints. The set of edges of a Scharlemann k -gon is called a *Scharlemann k -cycle*. A Scharlemann 2-cycle is also called an *S -cycle*. An S -cycle $\{e_1, e_2\}$ is called an *extended S -cycle* if $m \geq 4$ and the two edges e_1 and e_2 are the middle edges in a family of four adjacent parallel edges of Γ_S . An S -cycle $\{e_1, e_2\}$ is called a *doubly-extended S -cycle* if $m \geq 6$ and the two edges e_1 and e_2 are the middle edges in a family of six adjacent parallel edges of Γ_S .

The method of proof of [BCSZ2, Lemma 12.3] yields the following proposition.

Proposition 11.7. *Suppose that two vertices v and v' of Γ_S have the same orientation and are connected by a family of n parallel consecutive edges e_1, \dots, e_n .*

- (1) *If $n > m/2$, then there is an S -cycle in this family of edges.*
- (2)(a) *If $m \geq 4$ and $n > \frac{m}{2} + 1$, then either there is an extended S -cycle in this family of edges or both $\{e_1, e_2\}$ and $\{e_{n-1}, e_n\}$ are S -cycles.*
 - (b) *If $m \geq 4$ and $n > \frac{m}{2} + 2$, then there is an extended S -cycle in this family of edges.*
- (3)(a) *If $m \geq 6$ and $n > \frac{m}{2} + 3$, then either there is a doubly-extended S -cycle in this family of edges or both $\{e_2, e_3\}$ and $\{e_{n-2}, e_{n-1}\}$ are extended S -cycles.*
 - (b) *If $m \geq 6$ and $n > \frac{m}{2} + 4$, there is a doubly-extended S -cycle in this family of edges. \diamond*

Lemma 11.8. *Suppose that $\{e_1, e_2\}$ is an S -cycle in Γ_S and R the associated bigon face of Γ_S . If R lies on the ϵ -side of F , then $\dot{\Phi}_1^\epsilon$ contains a τ_ϵ -invariant component, so F is separating. Further, this component contains an \widehat{S} -essential annulus and \widehat{X}^ϵ admits a Seifert structure with base orbifold a disk with two cone points, at least one of which has order 2.*

Proof. Suppose that the S -cycle has label pair $\{j, j+1\}$. Then $\tau_\epsilon(b_j \cup e_1^* \cup b_{j+1}) = b_{j+1} \cup e_2^* \cup b_j$. Hence $b_j \cup e_1^* \cup b_{j+1}$ and $b_{j+1} \cup e_2^* \cup b_j$ are contained in the same component ϕ of $\dot{\Phi}^\epsilon$ and this component is τ_ϵ -invariant. Proposition 7.7 implies that S is connected and ϕ contains an \widehat{S} -essential annulus. Proposition 7.1 shows that ϕ is the unique component of $\dot{\Phi}^\epsilon$ to contain such an annulus. Finally, Proposition 7.1(3) implies that \widehat{X}^ϵ is of the form described in (4). \diamond

12. COUNTING FACES IN $\overline{\Gamma}_S$

In this section we examine the existence of triangle faces of $\overline{\Gamma}_S$ incident to vertices of small valency.

For each vertex v of Γ_S let $\varphi_j(v)$ be the number of corners of j -gons incident to v . Then

$$\text{valency}_{\overline{\Gamma}_S}(v) = |\partial S| \Delta(\alpha, \beta) - \varphi_2(v)$$

Set

$$\begin{aligned} \psi_3(v) &= \text{valency}_{\overline{\Gamma}_S}(v) - \varphi_3(v) \geq 0, \\ \mu(v) &= \varphi_2(v) + \frac{\varphi_3(v)}{3} \in \left\{ \frac{k}{3} : k \in \mathbb{Z} \right\} \end{aligned}$$

Lemma 12.1. *Suppose that v is a vertex of Γ_S and set $\mu(v) = |\partial S| \Delta(\alpha, \beta) - 4 + x$. Then*

$$\text{valency}_{\overline{\Gamma}_S}(v) = 6 - \frac{1}{2}(3x + \psi_3(v))$$

and

$$\varphi_3(v) = 3(\text{valency}_{\overline{\Gamma}_S}(v) - 4 + x)$$

Proof. We noted above that $\text{valency}_{\overline{\Gamma}_S}(v) = |\partial S| \Delta(\alpha, \beta) - \varphi_2(v)$. Thus

$$\text{valency}_{\overline{\Gamma}_S}(v) = |\partial S| \Delta(\alpha, \beta) - \mu(v) + \frac{\varphi_3(v)}{3} = 4 - x + \frac{\text{valency}_{\overline{\Gamma}_S}(v)}{3} - \frac{\psi_3(v)}{3},$$

and therefore

$$\text{valency}_{\overline{\Gamma}_S}(v) = \frac{3}{2} \left(4 - x - \frac{\psi_3(v)}{3} \right) = 6 - \frac{1}{2}(3x + \psi_3(v)).$$

On the other hand,

$$\varphi_3(v) = 3(\mu(v) - \varphi_2(v)) = 3(|\partial S| \Delta(\alpha, \beta) - 4 + x - \varphi_2(v)) = 3(\text{valency}_{\overline{\Gamma}_S}(v) - 4 + x).$$

Thus the lemma holds. \diamond

Proposition 12.2. *Suppose that v is a vertex of Γ_S .*

(1) *If $\mu(v) > |\partial S| \Delta(\alpha, \beta) - 4$, then $\text{valency}_{\overline{\Gamma}_S}(v) \leq 5$. Further,*

(a) *if $\text{valency}_{\overline{\Gamma}_S}(v) = 3$, then $\varphi_3(v) \geq 0$.*

(b) *if $\text{valency}_{\overline{\Gamma}_S}(v) = 4$, then $\varphi_3(v) \geq 1$.*

(c) *if $\text{valency}_{\overline{\Gamma}_S}(v) = 5$, then $\varphi_3(v) \geq 4$.*

(2) *If $\mu(v) = |\partial S| \Delta(\alpha, \beta) - 4$, then $4 \leq \text{valency}_{\overline{\Gamma}_S}(v) \leq 6$. Further,*

(a) *if $\text{valency}_{\overline{\Gamma}_S}(v) = 4$ then $\varphi_3(v) = 0$.*

(b) *if $\text{valency}_{\overline{\Gamma}_S}(v) = 5$ then $\varphi_3(v) = 3$.*

(c) *if $\text{valency}_{\overline{\Gamma}_S}(v) = 6$ then $\varphi_3(v) = 6$.*

Proof. Write $\mu(v) = |\partial S|\Delta(\alpha, \beta) - 4 + x$ where $x \geq 0$ is an element of $\{\frac{k}{3} : k \in \mathbb{Z}\}$. By Lemma 12.1 we have $\text{valency}_{\overline{\Gamma}_S}(v) \leq 6 - \frac{3x}{2}$. Thus $\text{valency}_{\overline{\Gamma}_S}(v) \leq \begin{cases} 6 & \text{if } x = 0 \\ 5 & \text{if } x > 0 \end{cases}$. Further, if $x = 0$, the same lemma implies that $\text{valency}_{\overline{\Gamma}_S}(v) = 6 - \frac{\psi_3(v)}{2}$. Since $\psi_3(v) = \text{valency}_{\overline{\Gamma}_S}(v) - \varphi_3(v)$, this is equivalent to $\text{valency}_{\overline{\Gamma}_S}(v) = 4 + \frac{\varphi_3(v)}{3}$. Thus $\text{valency}_{\overline{\Gamma}_S}(v) \geq 4$. The remaining conclusions follow from the identity $\varphi_3(v) = 3(\text{valency}_{\overline{\Gamma}_S}(v) - 4 + x)$ of Lemma 12.1. \diamond

Let V, E, F be the number of vertices, edges, and faces of $\overline{\Gamma}_S$.

Proposition 12.3.

- (1) *If the immersion surface is a disk, then $\sum_v \mu(v) \geq (|\partial S|\Delta(\alpha, \beta) - 4)V + 4$.*
- (2) *If the immersion surface is a torus, $\sum_v \mu(v) \geq (|\partial S|\Delta(\alpha, \beta) - 4)V$.*

Proof. First assume that Γ_S has no monogon faces. Since its vertices each have valency $|\partial S|\Delta(\alpha, \beta)$ we have $2E = |\partial S|\Delta(\alpha, \beta)V$. Let F_i be the number of i -faces so $F = \sum_i F_i$ and $2E = \sum_i iF_i$. Then

$$(|\partial S|\Delta(\alpha, \beta) - 4)V = 2E - 4V = 4(E - V) - 2E = 4\left(\sum_{\text{faces } f} \chi(f)\right) - \chi(Y) - 2E$$

Since $\chi(f) \leq 1$ for each face f , we have

$$\begin{aligned} (|\partial S|\Delta(\alpha, \beta) - 4)V &\leq 4(F - \chi(Y)) - 2E = \sum (4 - i)F_i - 4\chi(Y) \\ &\leq 2F_2 + F_3 - 4\chi(Y) \\ &= \sum_v \left(\varphi_2(v) + \frac{\varphi_3(v)}{3}\right) - 4\chi(Y) \\ &= \sum_v \mu(v) - 4\chi(Y) \end{aligned}$$

Thus the lemma holds when there are no monogons.

If there are monogons, it is easily verified that there is only one, f say, and that it is a collar on ∂Y when Y is a disk and a once-punctured torus when Y is a torus. In either case, $\overline{Y \setminus f}$ is a disk containing Γ_S without monogons. The first case implies that $\sum_v \mu(v) \geq (|\partial S|\Delta(\alpha, \beta) - 4)V + 4$, which implies the result. \diamond

Corollary 12.4.

- (1) *If the immersion surface is a disk there is a vertex v of Γ_S such that $\mu(v) > |\partial S|\Delta(\alpha, \beta) - 4$.*
- (2) *If the immersion surface is a torus, then either there is a vertex v of Γ_S such that $\mu(v) > |\partial S|\Delta(\alpha, \beta) - 4$ or $\mu(v) = |\partial S|\Delta(\alpha, \beta) - 4$ for each vertex. \diamond*

Proposition 12.5. *Suppose that $\mu(v) = |\partial S|\Delta(\alpha, \beta) - 4$ for each vertex v of Γ_S . Then each face of Γ_S is a disk. Further, if v is a vertex of Γ_S and*

- (1) *valency $_{\overline{\Gamma}_S}(v) = 4$, then $\varphi_4(v) = 4$.*

(2) $\text{valency}_{\overline{\Gamma}_S}(v) = 5$, then $\varphi_3(v) = 3$ and $\varphi_4(v) = 2$.

(3) $\text{valency}_{\overline{\Gamma}_S}(v) = 6$, then $\varphi_3(v) = 6$.

Proof. Corollary 12.4 shows that Y is a torus. Thus $0 = \chi(Y) = \sum_v \chi(v)$ where

$$\chi(v) = 1 - \frac{\text{valency}_{\overline{\Gamma}_S}(v)}{2} + \sum_{v \in \partial f} \frac{\chi(f)}{|\partial f|}$$

and f ranges over the faces of $\overline{\Gamma}_S$ containing v . From Proposition 12.2(2) we see that $\chi(v) \leq 0$ for all v . Hence $\chi(v) = 0$ for all v . This is only possible if the proposition holds. \diamond

13. PROOF OF THEOREM 2.7 WHEN F IS NON-SEPARATING

We show that when F is non-separating and $m \geq 3$, $\Delta(\alpha, \beta) \leq 4$ if $M(\alpha)$ is very small and $\Delta(\alpha, \beta) \leq 5$ otherwise. This follows from the two propositions below. Recall that $|\partial S| = 2m$ when F is non-separating.

Proposition 10.1 shows that $l_S \leq 2m - t_1^+ + 1$, so the weight of each edge in the reduced graph $\overline{\Gamma}_S$ of Γ_S is at most $2m - t_1^+ + 2$. Hence if v is a vertex of $\overline{\Gamma}_S$, $2m\Delta(\alpha, \beta)/\text{valency}_{\overline{\Gamma}_S}(v) \leq 2m - t_1^+ + 2$, so

$$(13.0.1) \quad \Delta(\alpha, \beta) \leq \left(\frac{2m - t_1^+ + 2}{2m} \right) \text{valency}_{\overline{\Gamma}_S}(v)$$

Proposition 13.1. *Suppose that F is non-separating and $t_1^+ > 0$. Then*

$$\Delta(\alpha, \beta) \leq \begin{cases} 4 & \text{if } m \leq 5 \text{ or } M(\alpha) \text{ is very small} \\ 5 & \text{if } m \geq 6 \end{cases}$$

Proof. If there is a vertex of $\overline{\Gamma}_S$ of valency 3 or less, Inequality 13.0.1 yields $\Delta(\alpha, \beta) \leq 3$, so we are done. Suppose then that all vertices are of valency 4 or more.

If $t_1^+ = 2$, then by Lemma 11.6 there are no triangle faces of Γ_S and therefore Lemma 11.5(2) implies that $\overline{\Gamma}_S$ is quadrilateral. Thus Y is a torus, so $M(\alpha)$ is not very small. Further, as all vertices have valency 4, Inequality 13.0.1 implies that $\Delta(\alpha, \beta) \leq 4$. Thus we are done.

If $t_1^+ > 2$, then $2m \geq t_1^+ \geq 4$, so $m \geq 2$. Corollary 12.4 and Proposition 12.2 imply that there is a vertex v of $\overline{\Gamma}_S$ of valency at most 5 if Y is a disk (e.g. if $M(\alpha)$ is very small) and at most 6 if it is a torus. Inequality 13.0.1 then shows that the proposition holds. \diamond

Proposition 13.2. *Suppose that F is non-separating and $t_1^+ = 0$.*

$$(1) \text{ If } M(\alpha) \text{ is very small, then } \Delta(\alpha, \beta) \leq \begin{cases} 4 & \text{if } m \geq 2 \\ 6 & \text{if } m = 1 \end{cases}$$

$$(2) \text{ If } M(\alpha) \text{ is not very small, then } \Delta(\alpha, \beta) \leq \begin{cases} 5 & \text{if } m \geq 3 \\ 6 & \text{if } m = 2 \\ 8 & \text{if } m = 1 \end{cases}$$

Proof. Suppose that $t_1^+ = 0$. By Lemma 11.6, $\bar{\Gamma}_S$ has no triangle face, so $\varphi_3(v) = 0$ for each vertex v of Γ_S . Hence $\bar{\Gamma}_S$ has a vertex v of valency at most 4 by Proposition 11.5(2). If there is a vertex of valency 3 or less, then Inequality 13.0.1 shows $\Delta(\alpha, \beta) \leq 4$ for $m \geq 2$ and $\Delta(\alpha, \beta) \leq 6$ for $m = 1$. If there are no vertices of valency less than 4, Proposition 11.5(2) implies that $\bar{\Gamma}_S$ is rectangular, so Y is a torus and $M(\alpha)$ is not very small. Thus assertion (1) of the lemma holds. By Inequality 13.0.1, $\Delta(\alpha, \beta) \leq 4 + 4/m$. It follows that $\Delta(\alpha, \beta) \leq 5$ if $m \geq 3$, $\Delta(\alpha, \beta) \leq 6$ if $m = 2$, and $\Delta(\alpha, \beta) \leq 8$ if $m = 1$. \diamond

14. PROOF OF THEOREM 2.7 WHEN F IS SEPARATING AND $t_1^+ + t_1^- \geq 4$

Proposition 14.1. *Suppose that F is separating and $t_1^+ + t_1^- \geq 4$. Then*

$$\Delta(\alpha, \beta) \leq \begin{cases} 4 & \text{if } M(\alpha) \text{ is very small} \\ 5 & \text{otherwise.} \end{cases}$$

Proof. Since F is separating, $S = F$ and $|\partial S| = m$.

If $t_1^\epsilon \geq 4$ for some ϵ then Proposition 10.1 shows that $l_S \leq m - 3$. Thus the weight of each edge in $\bar{\Gamma}_S$ is at most $m - 2$. If $t_1^\epsilon = 2$ for both ϵ , then $l_+, l_- \leq m - 2$, so $l_S \leq m - 2$. Thus the weight of each edge in $\bar{\Gamma}_S$ is at most $m - 1$. In either case, it follows that for each vertex v of Γ_S , $\frac{m\Delta(\alpha, \beta)}{\text{valency}_{\bar{\Gamma}_S}(v)} \leq m - 1$. Hence

$$(14.0.1) \quad \Delta(\alpha, \beta) \leq \left(\frac{m-1}{m}\right) \text{valency}_{\bar{\Gamma}_S}(v) < \text{valency}_{\bar{\Gamma}_S}(v)$$

Corollary 12.4 and Proposition 12.2 imply that there is a vertex v of valency 5 or less if Y is a disk, in particular if $M(\alpha)$ is very small, and of valency at most 6 otherwise. Inequality 14.0.1 then shows that the conclusion of the proposition hold. \diamond

15. THE RELATION ASSOCIATED TO A FACE OF Γ_S

The proof of Theorem 2.7 when F is separating and $t_1^+ + t_1^- \leq 2$ necessitates a deeper use of the properties of the intersection graph Γ_S . We begin with a description of the relations associated to its faces.

Recall that the boundary components of F have been indexed (mod m): b_1, b_2, \dots, b_m so that they appear successively around ∂M . For each ϵ we use $\tau_\epsilon(j) (= j \pm 1)$ to be the index such that $\tau_\epsilon(b_j) = b_{\tau_\epsilon(j)}$. Let σ_j be a path which runs from b_j to $b_{\tau_\epsilon(j)}$ in the annular component of $\partial M \cap X^\epsilon$ containing $b_j \cup b_{\tau_\epsilon(j)}$. Fix a base point $x_0 \in F$ and for each j a path η_j in F from x_0 to b_j . The loop $\eta_j * \sigma_j * \eta_{\tau_\epsilon(j)}^{-1}$ determines a class $x_j \in \pi_1(\hat{X}^\epsilon; x_0)$ well-defined up to our choice of the η_j . Clearly $x_j x_{\tau_\epsilon(j)} = 1$. (The use of x_j to describe this class is ambiguous in that it does not specify which value ϵ takes on. Nevertheless, whenever we use it the value of ϵ will be understood from the context.)

Recall that if $t_1^\epsilon = 0$, \hat{X}^ϵ admits a Seifert structure with base orbifold $D^2(p, q)$. There is a projection homomorphism $\pi_1(\hat{X}^\epsilon) \rightarrow \pi_1(D^2(p, q))$ obtained by quotienting out the normal

cyclic subgroup of $\pi_1(\widehat{X}^\epsilon)$ determined by the class of a regular Seifert fibre. We denote the image in $\pi_1(D^2(p, q))$ of an element $x \in \pi_1(\widehat{X}^\epsilon)$ by \bar{x} . Fix generators a, b of $\mathbb{Z}/p, \mathbb{Z}/q$ such that ab represents the class of the boundary circle of $D^2(p, q)$ in $\pi_1(D^2(p, q)) \cong \mathbb{Z}/p * \mathbb{Z}/q$.

Proposition 15.1. *If $t_1^\epsilon = 0$, then no x_j is peripheral in \widehat{X}^ϵ . Indeed, there are integers k, l and $\delta \in \{\pm 1\}$ such that x_j is sent to an element \bar{x}_j of the form $(ab)^k a^\delta (ab)^l$ in $\pi_1(D^2(p, q))$.*

Proof. It follows from the method of proof of Proposition 7.5 that $\widehat{X}^\epsilon = V \cup W$ where V and W are solid tori whose intersection is an essential annulus $(A, \partial A) \subset (\widehat{X}^\epsilon, \widehat{F})$. Further, if K_β is the core of the β filling solid torus, we can assume that $K_\beta \cap X^\epsilon$ is a finite union of arcs properly embedded in A . Consideration of the Seifert structure on \widehat{X}^ϵ then shows that the image of the projection of σ_j to $D^2(p, q)$ is a properly embedded arc which separates the two cone points. Thus there are integers k, l and $\delta \in \{\pm 1\}$ such that $\bar{x}_j = (ab)^k a^\delta (ab)^l \in \pi_1(D^2(p, q))$. Such an element is peripheral if and only if it equals $(ab)^n$ for some n . But then $a = (ab)^{\pm(n-k-l)}$ would be peripheral, which is false. \diamond

Consider an n -gon face f of Γ_S lying to the ϵ -side of F with boundary $c_1 \cup e_1 \cup c_2 \cup \dots \cup c_n \cup e_n$ where each c_i is a corner of f , e_i an edge of f , and they are indexed as they arise around ∂f . In this ordering, let b_{j_i} be the boundary component of F at c_i corresponding to $c_i \cap e_i$ and $b_{j'_i}$ that corresponding to $c_{i+1} \cap e_i$. (See Figure 1.)




FIGURE 1.

The relation

$$\prod_{i=1}^n w_i x_{j'_i} = 1$$

holds in $\pi_1(\widehat{X}^\epsilon)$ where w_i is represented by the loop $\eta_{j_i} * e_i^* * \eta_{j'_i}^{-1}$.

For each boundary component b_j of F , let \widehat{b}_j denote the meridional disk it bounds in \widehat{F} .

Corollary 15.2. *Suppose that e_1 is a negative edge of Γ_S whose end labels are the same. Suppose as well that e_1 is a boundary edge of a triangle face lying on the ϵ -side of F where $t_1^\epsilon = 0$. If the boundary label of e is j , then the loop $\widehat{b}_j \cup e_1^*$ is essential in \widehat{F} .*

Proof. The relation from the given face reads $x_j^{-1} w_1 x_j w_2 x_k w_3 = 1$ where $k \in \{1, 2, \dots, m\}$ and w_1, w_2, w_3 are the peripheral elements of $\pi_1(\widehat{X}^\epsilon)$ defined above. If $\widehat{b}_j \cup e_1^*$ is inessential in \widehat{F} ,

then $w_1 = 1$, so the relation gives $x_k = (w_3 w_2)^{-1}$ is peripheral, which contradicts Proposition 15.1. Thus the corollary holds. \diamond

As an immediate consequence of this corollary we have:

Corollary 15.3. *Suppose that e is a negative edge of Γ_S whose end labels are the same. Suppose as well that e is a boundary edge of a triangle face lying on the ϵ -side of F where $t_1^\epsilon = 0$. If the weight of e in the reduced graph $\bar{\Gamma}$ is $k + 1$, then e^* is contained in a component of $\check{\Phi}_k^{-\epsilon}$ which contains an \widehat{F} -essential annulus. \diamond*

16. PROOF OF THEOREM 2.7 WHEN F IS SEPARATING AND $t_1^+ + t_1^- = 2$

We assume that F is separating and $t_1^+ + t_1^- = 2$ in this section. There is an ϵ such that $t_1^\epsilon = 2$ and $t_1^{-\epsilon} = 0$. Without loss of generality we can suppose that $\epsilon = +$.

Proposition 16.1. *If F is separating and $t_1^+ = 2, t_1^- = 0$, then*

$$\Delta(\alpha, \beta) \leq \begin{cases} 5 & \text{if } m \geq 4 \\ 6 & \text{if } m = 2 \end{cases}$$

Proof. Proposition 10.1 shows that $l_+ \leq m - 2$ and $l_S \leq m - 1$. Thus the weight of each edge in $\bar{\Gamma}_S$ is at most m . Hence if there is a vertex of $\bar{\Gamma}_S$ of valency k , then $m\Delta(\alpha, \beta) \leq km$, so $\Delta(\alpha, \beta) \leq k$. In the case that $\mu(v) > m\Delta(\alpha, \beta) - 4$ for some vertex v of $\bar{\Gamma}_S$, Proposition 12.2(1) implies that $\Delta(\alpha, \beta) \leq 5$. In particular this is true when $M(\alpha)$ is very small by Corollary 12.4(1). By Corollary 12.4(2) we can therefore suppose that Y is a torus and $\mu(v) = m\Delta(\alpha, \beta) - 4$ for all vertices v of $\bar{\Gamma}_S$. Then $\Delta(\alpha, \beta) \leq 6$ by Proposition 12.2(2).

To complete the proof we shall suppose that $\Delta(\alpha, \beta) = 6$ and show that $m = 2$. In this case $\bar{\Gamma}_S$ has no vertices of valency 5 or less. Thus it is hexagonal (Proposition 11.5). As no edge of $\bar{\Gamma}_S$ has weight larger than m , each of its edges has weight m . It follows that $l_+ \geq m - 2$, and since we noted above that $l_+ \leq m - 2$, we have $l_+ = m - 2$. Thus each face of $\bar{\Gamma}_S$ lies on the $+$ -side of F .

Note that $t_{m-3}^+ < m$ since $l_+ = m - 2$. The fact that t_{2j+1}^+ is even couples with Proposition 6.3 to show that $t_{m-3}^+ = m - 2$. Thus $\check{\Phi}_{m-3}^+$ has at least $m - 2$ components. If some such component ϕ_0 contains at least three boundary components of F , $\check{\Phi}_{m-3}^+$ has at most $m - 3$ other components. But then ϕ_0 is tight, so there must be another component of $\check{\Phi}_{m-3}^+$ containing at least three boundary components and therefore $m - 2 \leq |\check{\Phi}_{m-3}^+| \leq m - 4$, a contradiction. Thus each component of $\check{\Phi}_{m-3}^+$ contains at most two boundary components of F .

Let b_1, b_2, \dots, b_m be the boundary components of F numbered in successive fashion around ∂M . Fix a triangle face f of $\bar{\Gamma}_S$ and let v_1, v_2 be two of its vertices. They are connected by a family e_1, e_2, \dots, e_m of mutually parallel edges of Γ_S successively numbered around v_1 so that e_1 is the boundary edge of f thought of as a face in Γ_S .

We can suppose that the tail of each e_i lies on v_1 and is labeled i . Let j be the label of the head of e_2 . If v_1 and v_2 are like-oriented, then j is odd and $b_2 \cup e_2^* \cup b_j$ is contained in a component ϕ

of $\dot{\Phi}_{m-3}^+$. From above, b_2 and b_j are the only boundary components of F ϕ contains. Similarly $b_{m-1} \cup e_{m-1}^* \cup b_{j+3}$ is contained in a component of $\dot{\Phi}_{m-3}^+$ and b_{m-1} and b_{j+3} are the only boundary components of F it contains. Let v_3 be the third vertex of f and consider the family of m edges of Γ_S parallel to the edge of f connecting v_1 and v_3 . The second edge from f in this family has label $m-1$ at v_1 and its label at v_3 must be $j+3$ if v_1 and v_3 are like-oriented and $m-1$ otherwise. Similarly, the second edge from f in the family of parallel edges corresponding to the edge of f connecting v_2 and v_3 has label $m-1$ at v_3 if v_2 and v_3 are like-oriented and $j-3$ otherwise. Since the orientations of v_1 and v_3 coincide if and only if those of v_2 and v_3 do, it follows that $j = m-1$ whatever the relative orientations of v_1 and v_3 . This implies that the head of each e_i has label $m+1-i$.

A similar argument shows that the head of e_i is labeled i if v_1 and v_2 are oppositely-oriented.

Suppose that $m \geq 4$ and fix a triangle face f of Γ_S with one positive boundary edge and two negative ones (Proposition 11.2). Let v_1, v_2, v_3 be the vertices of f chosen so that the edge between v_1 and v_2 is positive. Number the family of m parallel edges of Γ_S connecting v_1 and v_2 as in the previous paragraph. In particular e_1 is an edge of f . Let ϕ be the component of $\dot{\Phi}_{m-3}^+$ containing $b_2 \cup b_{m-1}$. Consideration of the $m-2$ successive bigons connected by e_3, e_4, \dots, e_{m-2} shows that $h_{m-3}^+(\phi) = \phi$ (cf. the end of §3.2). Equivalently, if $\epsilon = (-1)^{\frac{m}{2}}$ and $\phi' = (\tau_{-\epsilon} \circ \tau_\epsilon \circ \tau_{-\epsilon} \circ \dots \circ \tau_+)(\phi)$ (a composition of $\frac{m}{2} - 2$ factors), then $\tau_\epsilon(\phi') = \phi'$. Hence ϕ' , and therefore $\phi \subset \dot{\Phi}_{m-3}^+$ contains an \widehat{F} -essential annulus. It follows that the same is true for $\dot{\Phi}_j^+$ for each $j \leq m-3$. (See 3.2.1.) Proposition 7.1 now implies that \widehat{X}^+ admits a Seifert structure with base orbifold of the form $D^2(a, b)$ where $a, b \geq 2$. Furthermore, Proposition 8.2 implies that $m-3 \leq 2$. Thus $m \leq 4$. We assume now that $m = 4$ and show that this leads to a contradiction. This will complete the proof.

Consideration of the family of parallel positive edges adjacent to f shows that there is an S -cycle in Γ_S lying on the $+$ -side of F . Hence Lemma 11.8 implies that \widehat{X}^+ admits a Seifert structure with base orbifold $D^2(2, b)$ and $\dot{\Phi}_1^+$ has a unique component which completes to an \widehat{F} -essential annulus. It's not hard to see then that $\check{\Phi}_1^+$ has three components: two boundary parallel annuli and a 4-punctured sphere with two inner boundary components and two outer ones. If φ_+ denotes the slope on \widehat{F} of the latter component, it is the slope of the Seifert structure on \widehat{X}^+ .

Since $\Delta(\alpha, \beta) > 3$, β is not a singular slope and therefore $M(\beta)$ is not Seifert with base orbifold $S^2(a, b, c, d)$ where $(a, b, c, d) \neq (2, 2, 2, 2)$. Hence as $\dot{\Phi}_3^-$ contains an \widehat{F} -essential annulus, Proposition 8.1 implies that X^- is a twisted I -bundle. In particular, $\check{\Phi}_3^- = \tau_-(\check{\Phi}_1^+)$. Hence if A is an \widehat{F} -essential annulus containing $\check{\Phi}_3^-$, its slope φ_- is given by $(\tau_-)_*(\varphi_+)$. It follows that $\Delta(\varphi_+, \varphi_-) \equiv 0 \pmod{2}$. Thus either $\Delta(\varphi_+, \varphi_-) = 0$ and $\widehat{X}^+(\varphi_-)$ is the connected sum of two non-trivial lens spaces or $\Delta(\varphi_+, \varphi_-) \geq 2$ and $\widehat{X}^+(\varphi_-)$ is a Seifert manifold with base orbifold $S^2(2, b, \Delta(\varphi_+, \varphi_-))$. In either case, $\pi_1(\widehat{X}^+(\varphi_-))$ is non-abelian.

Let $H_{(14)}$ be the component of $(\overline{M(\beta)} \setminus \overline{M}) \cap \widehat{X}^+$ containing $b_1 \cup b_4$ and $\partial_0 H_{(14)}$ the annulus $H_{(14)} \cap X^+$. Then the image in X^+ of ∂f lies in $A \cup \partial_0 H_{(14)}$. Moreover, once oriented, ∂f intersects $\partial_0 H_{(14)}$ in three disjoint arcs exactly two of which are like-oriented. By an application of the Loop Theorem (see [He, Theorem 4.1]), there is a properly embedded disk $(D, \partial D) \subseteq$

$(X^+, A \cup \partial_0 H_{(14)})$ such that $\partial D \cap \partial_0 H_{(14)} \subseteq \partial f \cap \partial_0 H_{(14)}$ and ∂D algebraically intersects a core of $\partial_0 H_{(14)}$ a non-zero number of times (mod 3). There are two possibilities,

- (1) $\partial D \cap \partial_0 H_{(14)}$ consists of two like-oriented arcs, or
- (2) $\partial D \cap \partial_0 H_{(14)}$ consists of three arcs, two like-oriented and one oppositely-oriented.

Suppose that (1) arises. Then $\partial D = e_1 \cup a_1 \cup e_2 \cup a_2$ where a_1, e_1, a_2, e_2 are arcs arising successively around ∂D and a_1, a_2 are properly embedded in $H_{(14)}$ while e_1, e_2 are properly embedded in F . Let \widehat{b}_i be the disk in \widehat{F} with boundary b_i and fix a (fat) basepoint in \widehat{X}^+ to be $\widehat{b}_1 \cup e_1 \cup \widehat{b}_4$. We take η_1, η_4 to be constant paths (see §15). The ‘‘loop’’ e_2 carries a generator t of $\pi_1(A)$ as otherwise $M(\beta)$ would contain a P^3 connected summand. Thus the relation associated to D is $x_1^2 = t$. Further, there is a Möbius band B properly embedded in X^+ whose core carries the class x_1 . Consequently, t represents the class of the slope φ_+ . But by construction, it represents the class of φ_- . It follows that $\varphi_+ = \varphi_-$ so that $\widehat{X}^+(\varphi_-)$ is the connected sum of lens spaces $L(2, 1)$ and $L(n, m)$ for some n, m . Note that x_1 represents a non-trivial class in $H_1(\widehat{X}^+(\varphi_-))$. But the relation associated to f is of the form $t^a x_1 t^b x_1 t^c x_1^{-1} = 1$. In particular, x_1 is trivial in $H_1(\widehat{X}^+)/\langle t \rangle = H_1(\widehat{X}^+(\varphi_-))$. Thus possibility (1) cannot occur.

Suppose that (2) arises. Then $\partial D = e_1 \cup a_1 \cup e_2 \cup a_2 \cup e_3 \cup a_3$ where $a_1, e_1, a_2, e_2, a_3, e_3$ are arcs arising successively around ∂D and a_1, a_2, a_3 are properly embedded in $H_{(14)}$ while e_1, e_2, e_3 are properly embedded in F . We can suppose that the indices are chosen so that e_1 connects b_1 to b_4 , e_2 is a loop based at b_4 and e_3 is a loop based at b_1 . These loops are essential as otherwise we could isotope ∂D so that it intersects a core of $\partial_0 H_{(14)}$ once transversally. This would imply that we could isotope \widehat{F} in $M(\beta)$ to remove two points of intersection with the core of the β -filling solid torus contrary to Assumption 2.2. Fix the basepoint in \widehat{X}^+ to be $\widehat{b}_1 \cup e_1 \cup \widehat{b}_4$ and take η_1, η_4 to be constant paths. The arcs a_1, a_2, a_3 determine triples of distinct points, one on b_1 and one on b_4 , which we denote 1, 2, 3. The reader will verify that these triples are oppositely orientated on A . From this it follows an orientation on ∂D determines the same orientation on the loops $b_1 \cup e_2$ and $b_4 \cup e_3$. In particular they yield the same generator t of $\pi_1(A)$. (See Figure 2.)

Hence the relation associated to D is

$$1 = x_1^2 t x_1^{-1} t = x_1^3 (x_1^{-1} t)^2$$

Let $N(A)$ be a collar of A in \widehat{X}^+ and $N(D)$ a tubular neighbourhood of D in X^+ . Set $Q = N(A) \cup H_{(14)} \cup N(D)$. Then the boundary of Q is a torus and its fundamental group is presented by $\langle x_1, t : x_1^3 (x_1^{-1} t)^2 \rangle$. It follows that Q is a trefoil complement contained in \widehat{X}^+ . Since the latter has a Seifert structure with base orbifold $D^2(2, n)$, Q must be isotopic in $M(\beta)$ to \widehat{X}^+ . It follows from the presentation that t normally generates $\pi_1(\widehat{X}^+)$. Since the slope of A is φ_- we have $\widehat{X}^+(\varphi_-)$ is simply connected, contrary to our observation that it has a non-abelian fundamental group. Thus possibility (2) is also impossible. Therefore $\Delta(\alpha, \beta) \leq 5$ when $m = 4$. \diamond




FIGURE 2.

 17. EXTENDED S -CYCLES IN Γ_S

In this section we examine the implications of the existence of extended and doubly-extended S -cycles in Γ_S when $t_1^+ = t_1^- = 0$. (See §11.)

Proposition 17.1. *Suppose that $t_1^+ = t_1^- = 0$ and $\{e_0, e_1, e_2, e_3\}$ is an extended S -cycle in Γ_S where $\{e_1, e_2\}$ is an S -cycle. Let R be the bigon face between e_1 and e_2 and suppose that R lies on the ϵ -side of \widehat{S} . Then either*

(i) β is a singular slope so $\Delta(\alpha, \beta) \leq 3$.

(ii) $\epsilon = +$, X^- is a twisted I -bundle, and \widehat{X}^+ admits a Seifert structure with base orbifold a disk with two cone points, exactly one of which has order 2.

Proof. Suppose that the S -cycle $\{e_1, e_2\}$ has label pair $\{j, j+1\}$. By Lemma 11.8, $S = F$ is connected and $\dot{\Phi}_1^\epsilon$ has a unique component ϕ which contains an \widehat{S} -essential annulus. Further, ϕ is also τ_ϵ -invariant and \widehat{X}^ϵ admits a Seifert structure with base orbifold a disk with two cone points, at least one of which has order 2. The proof of Lemma 11.8 also shows that a regular neighbourhood N of $b_j \cup e_1^* \cup b_{j+1} \cup e_2^*$ in S is also τ_ϵ -invariant, at least up to isotopy in S , and so there is a Möbius band B properly embedded in $(\widehat{X}^\epsilon, N)$. Thus N contains an \widehat{S} -essential annulus with core ∂B which is vertical in \widehat{X}^ϵ .

Since $\{e_0, e_1, e_2, e_3\}$ is an extended S -cycle, $b_j \cup e_1^* \cup b_{j+1} \cup e_2^*$, and therefore N , is contained in $\dot{\Phi}_1^{-\epsilon}$. Let ϕ' be the component of $\dot{\Phi}_1^{-\epsilon}$ which contains N and Σ' the component of $\dot{\Sigma}_1^{-\epsilon}$ which contains ϕ' . The genera of ϕ and ϕ' cannot both be 1 as otherwise, Proposition 7.1 implies that both \widehat{X}^+ and \widehat{X}^- are twisted I -bundles over the Klein bottle, contrary to Corollary 7.6.

Suppose first that $\text{genus}(\phi) = \text{genus}(\phi') = 0$. Then $\Phi_1^- \neq S$, so X^- cannot be a twisted I -bundle. In particular, $X^{-\epsilon}$ does not admit a properly embedded non-separating annulus (cf. Lemma 4.7). Proposition 7.1(3) then shows that \widehat{X}^- admits a Seifert structure with base

orbifold a disk with two cone points in which $\partial B \subseteq N \subseteq \phi'$ is vertical. Thus $M(\beta)$ admits a Seifert structure with base orbifold the 2-sphere with four cone points. Their orders cannot all be 2 by Corollary 7.6. Hence β is a singular slope ([BGZ1, Theorem 1.7]) so $\Delta(\alpha, \beta) \leq 3$ ([BGZ1, Theorem 1.5]).

Suppose next that $\text{genus}(\phi) = 1$ and $\text{genus}(\phi') = 0$. Then Corollary 6.4 implies that $\phi = S$. Thus X^ϵ is a twisted I -bundle, so $\epsilon = -$. If Σ' is either a product I -bundle which separates $X^{-\epsilon} = X^+$ or a twisted I -bundle, then by Proposition 7.1(3), \widehat{X}^+ admits a Seifert structure with base orbifold a disk with two cone points in which $\partial B \subseteq N \subseteq \phi'$ is vertical. Corollary 7.6 shows that at least one of the cone points has order larger than 2. Thus $M(\beta)$ admits a Seifert structure with base orbifold the 2-sphere with four cone points, at least one of which has order larger than 2. Thus (i) occurs. If, on the other hand, Σ' is a product I -bundle which does not separate X^+ , Proposition 7.1(3) implies that there is a Seifert structure on \widehat{X}^+ for which $\partial B \subseteq \phi'$ contains a fibre and whose base orbifold is a Möbius band with at most one cone point. Since \widehat{X}^+ is not a twisted I -bundle over the Klein bottle (Corollary 7.6), there is exactly one cone point. It follows that $M(\beta)$ admits a Seifert structure with base orbifold a projective plane with three cone points. Thus (i) holds.

Finally suppose that $\text{genus}(\phi) = 0$ and $\text{genus}(\phi') = 1$. Then Corollary 6.4 implies that $X^{-\epsilon}$ is a twisted I -bundle, so $\epsilon = +$. From above, \widehat{X}^+ admits a Seifert structure with base orbifold a disk with two cone points, at least one of which has order 2. They cannot both have order 2 by Corollary 7.6. This is case (ii). \diamond

Proposition 17.2. *Suppose that $t_1^+ = t_1^- = 0$. If Γ_S contains a doubly-extended S -cycle, then $\Delta(\alpha, \beta) \leq 3$.*

Proof. Suppose that $\Delta(\alpha, \beta) > 3$. Proposition 17.1 implies that X^- is a twisted I -bundle, the image of the S -cycle rectangle is contained in X^+ , and \widehat{X}^+ admits a Seifert structure with base orbifold a disk with two cone points, exactly one of which has order 2.

Suppose that the S -cycle $\{e_1, e_2\}$ has label pair $\{j, j+1\}$. It follows from the proof of Lemma 11.8 that there is a τ_+ -invariant regular neighbourhood N of $e_1^* \cup b_j \cup e_2^* \cup b_{j+1}$ contained in a τ_+ -invariant component ϕ_0 of $\dot{\Phi}_1^+$ such that $\widehat{\phi}_0$ is an \widehat{F} -essential annulus. Further, ϕ_0 is the unique component of $\dot{\Phi}_1^+$ to contain an \widehat{F} -essential annulus. Since $\{e_1, e_2\}$ is a doubly-extended S -cycle, $\tau_-(N) \subseteq \dot{\Phi}_1^+$. But $\tau_-(N)$ contains an \widehat{F} -essential annulus, so $\tau_-(N) \subseteq \phi_0$. Since N contains a core of $\widehat{\phi}_0$, it follows that $\widehat{\tau_-(\phi_0)}$ is isotopic to $\widehat{\phi}_0$ in \widehat{F} . In particular, $\widehat{\phi}_0$ is vertical in some Seifert structure on \widehat{X}^- . Thus $M(\beta)$ is Seifert with base orbifold either $P^2(2, n)$ or $S^2(2, 2, 2, n)$ where $n > 2$. Since $\Delta(\alpha, \beta) > 3$, β is not a singular slope ([BGZ1, Theorem 1.5]), so $M(\beta)$ has base orbifold $P^2(2, n)$ ([BGZ1, Theorem 1.7]).

Since N is τ_+ -invariant and connected, it contains a τ_+ -invariant simple closed curve C which is necessarily a core of $\widehat{\phi}_0$. There is a Möbius band B properly embedded in X^+ with boundary C . First suppose that $C \cap \tau_-(C) = \emptyset$. Then there is an annulus A_- properly embedded in X^- with $\partial A_- = C \cup \tau_-(C)$. Since C is vertical in $M(\beta)$, A_- is non-separating in X^- . Hence $C \cup \tau_-(C)$ splits \widehat{F} into two annuli, each containing $m/2$ boundary components of F . Let A_+ be

the properly embedded annulus in X^+ which is the frontier of the component of $\dot{\Sigma}_1^+$ containing ϕ_0 . Since C and $\tau_-(C)$ are disjoint curves in ϕ_0 , each isotopic to a core of $\widehat{\phi}_0$, \widehat{F} is the union of four annuli B_1, B_2, B_3, B_4 with disjoint interiors such that $\widehat{\phi}_0 = B_1 \cup B_2 \cup B_3$, $B_4 = \widehat{F} \setminus \widehat{\phi}_0$, and $\partial B_2 = C \cup \tau_-(C)$. Let $b_j = |B_j \cap \partial F|$. By construction, $b_2 = b_1 + b_3 + b_4 = m/2$. Since C is a τ_+ -invariant curve in ϕ_0 , $m/2 \geq b_1 = b_2 + b_3 = m/2 + b_3$. Hence $b_3 = 0$. There are solid tori $V_1, V_2 \subseteq X^+$ where V_1 is a regular neighbourhood of B and V_2 has boundary $A_+ \cup B_4$. By Lemma 4.2, B_4 has winding number at least 2 in V_2 . It follows that a regular neighbourhood of $V_1 \cup A_- \cup B_3 \cup V_2$ in M is Seifert with incompressible boundary (Lemma 4.1), which is impossible.

Next suppose that $C \cap \tau_-(C) \neq \emptyset$. Since $C \cup \tau_-(C)$ is connected, τ_- -invariant, and contained in ϕ_0 , there is a τ_- -invariant simple closed curve C' in ϕ_0 , necessarily a core of $\widehat{\phi}_0$. In particular C' is vertical in \widehat{X}_+ . It follows that there is a Möbius band B' properly embedded in X^- with boundary C' . Since C' is vertical in the Seifert structure on \widehat{X}^- with base orbifold $D^2(2, 2)$, $M(\beta)$ admits a Seifert structure with base orbifold $S^2(2, 2, 2, n)$, contrary to our previous deductions. This final contradiction completes the proof. \diamond

18. PROOF OF THEOREM 2.7 WHEN X^- IS NOT AN I -BUNDLE AND $t_1^+ = t_1^- = 0$

Throughout this section we assume

(18.0.1) F is separating, $\Delta(\alpha, \beta) > 3$, $t_1^+ = t_1^- = 0$, X^- is not a twisted I -bundle, and $m \geq 4$

By Proposition 9.4 there is a disk $D_\epsilon \subseteq \widehat{F}$ containing $\check{\Phi}_2^\epsilon$. We choose our base point and the images of the paths η_j to lie in $D_\epsilon \cap F$ (cf. §15) when we are interested in a relation associated to a face lying to the $-\epsilon$ -side of F .

18.1. Background results.

Lemma 18.1. *Suppose that conditions 18.0.1 hold and e is a negative edge of Γ_S whose end labels are the same. Suppose as well that e is a boundary edge of a triangle face f of Γ_S . Then the weight of the corresponding edge \bar{e} in $\overline{\Gamma}_S$ is at most 2.*

Proof. Suppose that f lies on the ϵ -side of F . Then if the weight of \bar{e} is at least 3, the image of e in F is contained in $\widehat{\Phi}_2^{-\epsilon} \subseteq D_{-\epsilon}$. Corollary 15.3 then shows the labels at the ends of j are different. \diamond

Proposition 11.7 combines with the fact that Γ_S contains no extended S -cycles (Proposition 17.1) to imply the following lemma.

Lemma 18.2. *Suppose that conditions 18.0.1 hold. Then the weight of a positive edge of $\overline{\Gamma}_S$ is at most $\frac{m}{2} + 2$. In particular, its weight is less than m if $m \geq 6$ and less than or equal to m if $m = 4$. \diamond*

Lemma 18.3. *Suppose that conditions 18.0.1 hold and that $\bar{\Gamma}_S$ has a triangle face f with edges \bar{e}, \bar{e}' where $wt(\bar{e}') > 2$. Then $wt(\bar{e}) \leq m$. Further, if \bar{e} is negative of weight m , the permutation associated to the corresponding family of edges has order $\frac{m}{2}$.*

Proof. Let v be the common vertex of \bar{e} and \bar{e}' , and let v' be the other vertex of \bar{e} . Let e' be the lead edge of \bar{e}' incident to f . Suppose that f lies on the ϵ -side of F .

Suppose otherwise that $wt(\bar{e}) > m$ and let $e_1, e_2, \dots, e_m, e_{m+1}$ be the $m+1$ consecutive edges in \bar{e} -family with e_1 as the lead edge incident to f . We may assume that the labels of $e_1, e_2, \dots, e_m, e_{m+1}$ at v are $1, 2, \dots, m, 1$ respectively. So the label of e' at v is m .

Lemma 18.2 implies that \bar{e} is a negative edge so the parity rule implies that the labels of $e_1, e_2, \dots, e_m, e_{m+1}$ at v' are $1+2k, 2+2k, \dots, m, 1, 2, \dots, 2k, 1+2k$ respectively, for some $0 < k < m/2$ (Lemma 18.1).

As both e_1 and e' are contained in $\widehat{\Phi}_2^{-\epsilon}$ and f is on the ϵ -side of F , f gives the relation

$$(18.1.1) \quad x_{2k}x_m^{-1}x_j \in \pi_1(\widehat{F})$$

for some j . Let B_i be the bigon face between e_i and e_{i+1} for $i = 1, \dots, m$. Note that B_i is on the ϵ -side of F if and only if i is even. Also note that for each even i with $2 < i < m$, the images in F of the two edges of B_i both lie in $\widehat{\Phi}_2^{-\epsilon}$. Also e_3^* is contained in $\widehat{\Phi}_2^{-\epsilon}$. So for each $2 < i = 2p < m$, B_i gives the relation $x_{2p}x_{2p+2k}^{-1} = 1$, so

$$(18.1.2) \quad x_{2p} = x_{2p+2k} \text{ for } 2 < 2p < m$$

Similarly B_2 gives the relation

$$(18.1.3) \quad x_2x_{2+2k}^{-1} = u \in \pi_1(\widehat{F})$$

Now consider the permutation given by the first m edges e_1, \dots, e_m . The orbit of the label $2k$ is $\{2k, 4k, 6k, \dots, m\}$ where we consider the labels (mod m). Applying 18.1.2 successively shows that if 2 is not in this orbit (i.e. the permutation has order less than $m/2$), then

$$x_{2k} = x_{4k} = \dots = x_m$$

Thus $x_{2k}x_m^{-1} = 1$. But comparing with 18.1.1 shows that $x_j \in P$, which contradicts Proposition 15.1. On the other hand, if 2 is in this orbit then by 18.1.3,

$$x_{2k} = x_{4k} = \dots = x_2 = ux_{2k+2} = ux_{4k+2} = \dots = ux_m$$

Thus $x_{2k}x_m^{-1} = u \in \pi_1(\widehat{F})$ which combines with 18.1.1 to yield a similar contradiction. This proves the first assertion of the lemma.

Next suppose that \bar{e} is negative of weight m and let e_1, e_2, \dots, e_m be the m consecutive edges in \bar{e} -family with e_1 as the lead edge incident to f . As above we take the labels of e_1, e_2, \dots, e_m at v to be $1, 2, \dots, m$ respectively and those at v' to be $1+2k, 2+2k, \dots, m, 1, 2, \dots, 2k$ respectively, for some $0 < k < m/2$ (Lemma 18.1). Similar to identity 18.1.2 we have $x_{2p} = x_{2p+2k}$ for $2 < 2p < m-2$ and $x_2x_{2+2k}^{-1} = u \in \pi_1(\widehat{F})$. If the permutation $j \mapsto j+2k \pmod{m}$ does not have order $\frac{m}{2}$ then neither 2 nor $m-2$ lie in the orbit $\{2k, 4k, 6k, \dots, m\}$ of the label $2k$. Thus

$x_{2k} = x_{4k} = \dots = x_m$ so plugging $x_{2k}x_m^{-1} = 1$ into 18.1.1 yields the contradiction $x_j \in \pi_1(\widehat{F})$. This completes the proof of the lemma. \diamond

Lemma 18.4. *Suppose that conditions 18.0.1 hold. If $\Delta(\alpha, \beta) > 5$, then $\overline{\Gamma}_S$ is hexagonal.*

Proof. By Proposition 11.5, it suffices to show that there is no vertex of valency 5 or less.

Suppose otherwise that v is a vertex of $\overline{\Gamma}_S$ of valency 5 or less. Since $m\Delta(\alpha, \beta)/\text{valency}_{\overline{\Gamma}_S}(v) \leq m + 1$ (Proposition 10.1), we have

$$(18.1.4) \quad 6 \leq \Delta(\alpha, \beta) \leq \text{valency}_{\overline{\Gamma}_S}(v) + \frac{\text{valency}_{\overline{\Gamma}_S}(v)}{m}$$

Hence $\text{valency}_{\overline{\Gamma}_S}(v) = 5$, $m = 4$, and $\Delta(\alpha, \beta) = 6$. It follows that the weights of the edges incident to v are 4, 5, 5, 5, 5. Lemma 18.3 implies that there can be no triangle faces incident to v . In other words, $\varphi_3(v) = 0$. Then

$$\mu(v) = \varphi_2(v) = m\Delta(\alpha, \beta) - \text{valency}_{\overline{\Gamma}_S}(v) = m\Delta(\alpha, \beta) - 5$$

Hence by Corollary 12.4 there is a vertex v_0 of $\overline{\Gamma}_S$ with $\mu(v_0) > m\Delta(\alpha, \beta) - 4$. Then Proposition 12.2 shows that v_0 has valency 5 or less. As above we have $\text{valency}_{\overline{\Gamma}_S}(v_0) = 5$ and the weights of the edges incident to v_0 are 4, 5, 5, 5, 5. By Corollary 12.2, $\varphi_3(v_0) \geq 4$, so in particular there is a triangle face of $\overline{\Gamma}_S$ with two edges of weight 5, which is impossible by Lemma 18.3. Thus there is no vertex v of $\overline{\Gamma}_S$ of valency 5 or less, so the lemma holds. \diamond

18.2. Proof. We prove Theorem 2.7 under conditions 18.0.1.

Assume that $\Delta(\alpha, \beta) > 5$ in order to derive a contradiction. Recall that $\overline{\Gamma}_S$ is hexagonal by Lemma 18.4. In particular Y is a torus.

Since X^- is not a twisted I -bundle, Proposition 7.1 implies that for each ϵ , $\dot{\Sigma}_1^\epsilon$ has a unique component and $\check{\Phi}_1^\epsilon$ is the union of at most two components, each an \widehat{F} -essential annulus.

Suppose that there is an edge \bar{e} of weight $m + 1$ incident to a vertex v of $\overline{\Gamma}_S$. Since $\overline{\Gamma}_S$ is hexagonal, Lemma 18.3 implies that the two edges of $\overline{\Gamma}_S$ incident to v which are adjacent to \bar{e} have weights at most 2. Then the sum of the weights of the six edges incident to v is at most $4m + 5$. On the other hand, this is $m\Delta(\alpha, \beta) \geq 6m$. Hence $6m \leq 4m + 5$, which is impossible for $m \geq 4$. Thus the weight of each edge in $\overline{\Gamma}_S$ is at most m . But then $6m \leq m\Delta(\alpha, \beta) \leq 6m$, so each edge of $\overline{\Gamma}_S$ has weight m and $\Delta(\alpha, \beta) = 6$.

As $\overline{\Gamma}_S$ is hexagonal, it has positive edges. Then Lemma 18.2 implies that $m \leq \frac{m}{2} + 2$. Thus $m = 4$ and the weight of any edge in $\overline{\Gamma}_S$ is 4. Proposition 11.2 implies that there is a triangle face f with one positive edge \bar{e}_1 and two negative edges \bar{e}_2, \bar{e}_3 . Let v_1, v_2, v_3 be its vertices where v_1 is determined by \bar{e}_1 and \bar{e}_2 , v_2 is determined by \bar{e}_2 and \bar{e}_3 , and v_3 is determined by \bar{e}_2 and \bar{e}_3 . Let e_1, e_2, e_3 denote the lead edges of $\bar{e}_1, \bar{e}_2, \bar{e}_3$ at f . Without loss of generality we can take the label of e_1 at v_1 to be 1 and that of e_2 to be 4. Lemma 18.1 shows that e_2 has label 2 at v_2 , so e_3 has label 3 there. Lemma 18.1 then shows that the label of e_3 at v_3 is 1, so the label of e_1 at v_3 is 4. But then the four edges of Γ_S parallel to \bar{e}_1 form an extended S -cycle,

which contradicts Proposition 17.1. This final contradiction completes the proof of Theorem 2.7 when X^- is not a twisted I -bundle. \diamond

19. PROOF OF THEOREM 2.7 WHEN X^- IS A TWISTED I -BUNDLE AND $t_1^+ = 0$

We assume throughout this section that

$$(19.0.1) \quad F \text{ is separating, } \Delta(\alpha, \beta) > 3, t_1^+ = 0, X^- \text{ is a twisted } I\text{-bundle, and } m \geq 4$$

Note that $t_1^- = 0$ when X^- is a twisted I -bundle.

19.1. Background results. As X^+ is not a twisted I -bundle, Proposition 7.1 implies that $\hat{\Sigma}_1^+$ has a unique component and $\check{\Phi}_1^+ = \dot{\Phi}_1^+$ is the union of one or two components each of which completes to an \hat{F} -essential annulus. Let ϕ_+ be the slope on \hat{F} of these annuli and note that it is the slope of the Seifert structure on \hat{X}^+ (Proposition 7.1). Set $\alpha_- = \hat{\tau}_-(\phi_+)$, the slope on \hat{F} determined by $\dot{\Phi}_3^- = \tau_-(\dot{\Phi}_1^+)$. As $\hat{\tau}_-$ is a fixed-point free orientation reversing involution, $\Delta(\phi_+, \alpha_-)$ is even.

For the rest of this section we take $A_- = \dot{\Phi}_3^- = \dot{\Phi}_2^- = \tau_-(\dot{\Phi}_1^+) \subset F$. Also we take a disk D in \hat{A}_- containing all \hat{b}_j , choose the paths η_j (defined in Section 15) in D , and define the elements x_j of $\pi_1(\hat{X}^+)$ as in Section 15. It follows that if e is an edge of a face f of Γ_S lying on the $+$ -side of F and the image of e in F lies in $\dot{\Phi}_2^-$, then the associated element of $\pi_1(\hat{F})$ determined by e is a power of t , the element determined by a core of \hat{A}_- .

Lemma 19.1. *Suppose that conditions 19.0.1 hold. Then $\pi_1(\hat{X}^+(\alpha_-))$ is not abelian.*

Proof. The base orbifold of \hat{X}^+ has the form $D^2(p, q)$ for some $p, q \geq 2$. Since $\Delta(\phi_+, \alpha_-)$ is even, it can't be 1. Thus $\hat{X}^+(\alpha_-)$ is either $L_p \# L_q$ or is Seifert fibred with base orbifold $S^2(p, q, \Delta(\phi_+, \alpha_-))$ having three cone points. In either case, its fundamental group is not abelian. \diamond

For each $y \in \pi_1(\hat{X}^+)$ we use \bar{y} to denote its image in $\pi_1(\hat{X}^+(\alpha_-))$.

Definition 19.2. Define $P \leq \pi_1(\hat{X}^+(\alpha_-))$ to be the subgroup generated by $\pi_1(\hat{F})$.

Lemma 19.3. *Suppose that conditions 19.0.1 hold. For no j is the image of x_j in $\pi_1(\hat{X}^+(\alpha_-))$ contained in P .*

Proof. We noted in §15 that there are generators a, b of $\pi_1(D^2(p, q)) \cong \mathbb{Z}/p * \mathbb{Z}/q$ such that ab generates its peripheral subgroup and the image of each x_j in $\pi_1(D^2(p, q))$ is of the form $(ab)^{r_j} a^{\epsilon_j} (ab)^{s_j}$ where $r_j, s_j \in \mathbb{Z}$ and $\epsilon_j \in \{\pm 1\}$. Thus if the image of some x_j in $\pi_1(\hat{X}^+(\alpha_-))$ is contained in P , the image of each x_j in $\pi_1(\hat{X}^+(\alpha_-))$ is contained in P , contrary to Lemma 19.1. \diamond

Lemma 19.4. *Suppose that conditions 19.0.1 hold. Each face of Γ_S which lies on the $--$ -side of F has an even number of edges. In particular, each triangle face lies on the $+-$ -side of F .*

Proof. The boundary of the face intersects the Klein bottle core of \widehat{X}^- transversely in k points where k is the number of edges of the face. Since this curve is homologically trivial, k is even. \diamond

Given this lemma, the fact that Γ_S contains no doubly-extended S -cycles (Proposition 17.2), and the fact that the S -cycle bigon in an extended S -cycles lies to the $+-$ -side of F (Proposition 17.1) we deduce the following lemma.

Lemma 19.5. *Suppose that conditions 19.0.1 hold. The weight of a positive edge of $\overline{\Gamma}_S$ is at most $\frac{m}{2} + 3$ if m is not divisible by 4 and $\frac{m}{2} + 4$ otherwise. In particular, its weight is less than m if $m \geq 10$ and less than or equal to m if $m \geq 6$. \diamond*

Lemma 19.6. *Suppose that conditions 19.0.1 hold. Suppose that e is a negative edge of Γ_S whose end labels are the same. Suppose as well that e is a boundary edge of a triangle face f of Γ_S . Then the weight of the corresponding edge \bar{e} in $\overline{\Gamma}_S$ is at most 2.*

Proof. Suppose otherwise that the weight of \bar{e} is at least 3. Then e^* is contained in $\widehat{\Phi}_2^-$. Let j be the label of e at its two end points. The triangle face f is on the $+-$ -side of F and the associated relation is

$$x_j^{-1} t^a x_j w_2 x_k w_3 = 1$$

in $\pi_1(\widehat{X}^+)$, where t is the class of a loop in the annulus $\widehat{\Phi}_2^-$ corresponding to the slope α_- , and $w_1, w_2 \in \pi_1(\widehat{F})$. Hence the relation implies that the image of x_k is contained in $P \leq \pi_1(\widehat{X}^+(\alpha_-))$, contrary to Lemma 19.3. Thus the lemma holds. \diamond

Lemma 19.7. *Suppose that conditions 19.0.1 hold and that $\overline{\Gamma}_S$ has a triangle face f with edges \bar{e}, \bar{e}' where $wt(\bar{e}') > 2$.*

(1) *If \bar{e} is negative, then $wt(\bar{e}) \leq m$. Further, if $wt(\bar{e}) = m$, then the permutation associated to the \bar{e} -family of parallel edges in Γ_S has order $\frac{m}{2}$.*

(2) *If \bar{e} is positive, then $wt(\bar{e}) \leq m - 1$ for $m > 6$ and $wt(\bar{e}) \leq m$ for $m = 4, 6$.*

Proof. Let v be the common vertex of \bar{e} and \bar{e}' , and let v' be the other vertex of \bar{e} . Let e' be the lead edge of \bar{e}' incident to f .

The proof of assertion (1) mirrors that of Lemma 18.3. The only difference is that we work with the images $\bar{x}_j \in \pi_1(\widehat{X}^+(\alpha_-))$ rather than the $x_j \in \pi_1(\widehat{X}^+)$ and replace the contradiction to Proposition 15.1 with one to Lemma 19.3.

Next we consider assertion (2). Suppose that \bar{e} is a positive edge of $\overline{\Gamma}_S$ with $wt(\bar{e}) > m - 1$. First we show that $m \leq 6$.

Let e_1, e_2, \dots, e_m be the m consecutive edges in the \bar{e} -family with e_1 as the lead edge incident to f . We may assume that the labels of e_1, e_2, \dots, e_m at v are $1, 2, \dots, m$ respectively. The label of e' at v is then m .

By the parity rule, the labels of e_1, e_2, \dots, e_m at v' are $2k, 2k - 1, \dots, 1, m, m - 1, \dots, 2k + 1$ respectively, for some $0 < k \leq m/2$. So the triangle face f gives the relation

$$(19.1.1) \quad \bar{x}_{2k+1}\bar{x}_1\bar{x}_j \in P$$

Again let B_i be the bigon face between e_i and e_{i+1} for $i = 1, \dots, m - 1$. If $2 < 2k < m$, then the image of e_{2k} in F is contained in $\widehat{\Phi}_2^-$ and so the bigon face B_{2k} gives the relation

$$(19.1.2) \quad \bar{x}_{2k+1}\bar{x}_1 \in P$$

Equations 19.1.1 and 19.1.2 imply that \bar{x}_j is peripheral, a contradiction. Thus $2k = 2$ or $2k = m$.

If $m > 6$ and $2k = 2$, then $\{e_{m/2+1}, e_{m/2+2}\}$ is a doubly extended S -cycle, giving a contradiction.

If $m > 4$ and $2k = m$, then $\{e_{m/2}, e_{m/2+1}\}$ is a doubly extended S -cycle, giving a contradiction again.

If $m = 6$, $wt(\bar{e}) > 6$, and $2k = 2$, then $\{e_4, e_5\}$ is a doubly extended S -cycle, giving a contradiction.

Finally suppose $m = 4$ and $wt(\bar{e}) > m = 4$. The label of e at v' is either 2 or 4. If it is 2, $\{e_3, e_4\}$ is an extended S -cycle lying on the $--$ side of F , giving a contradiction. If it is 4, then the face f shows $\bar{x}_1^2\bar{x}_j \in P$ for some j while the bigon between e_4 and e_5 gives $\bar{x}_1^2 \in P$. But then $\bar{x}_j \in P$, which contradicts Lemma 19.3. \diamond

19.2. Proof when $\dot{\Phi}_3^+$ is a union of tight components. The hypothesis $\dot{\Phi}_3^+$ is a union of tight components implies that the edges of $\bar{\Gamma}_S$ have weight bounded above by $m + 2$ (Corollary 11.4). Thus, as in the proof of Lemma 18.4 we have

$$(19.2.1) \quad 6 \leq \Delta(\alpha, \beta) \leq \text{valency}_{\bar{\Gamma}_S}(v) + 2\left(\frac{\text{valency}_{\bar{\Gamma}_S}(v)}{m}\right)$$

Lemma 19.8. *Suppose that conditions 19.0.1 hold and $\dot{\Phi}_3^+$ is a union of tight components. If $\Delta(\alpha, \beta) > 5$, then $\mu(v) = m\Delta(\alpha, \beta) - 4$ for all vertices v of $\bar{\Gamma}_S$.*

Proof. Assume that there is some vertex v with $\mu(v) > m\Delta(\alpha, \beta) - 4$. Proposition 12.2 implies that $\text{valency}_{\bar{\Gamma}_S}(v)$ is at most 5 while inequality 19.2.1 shows that $\text{valency}_{\bar{\Gamma}_S}(v)$ is at least 4 and if it is 4, then $m = 4, \Delta(\alpha, \beta) = 6$, and each edge incident to v has weight 6. Since Proposition 12.2 implies $\varphi_3(v) \geq 1$, Lemma 19.7 shows that this case is impossible. Assume then that $\text{valency}_{\bar{\Gamma}_S}(v) = 5$. Proposition 12.2 implies $\varphi_3(v) \geq 4$ and therefore the sum of the weights of the edges incident to v is at most $5m + 2$ by Lemma 19.7. But this sum is bounded below by $\Delta(\alpha, \beta)m \geq 6m$, which is impossible for $m \geq 4$. Corollary 12.4 then implies the desired conclusion. \diamond

Lemma 19.9. *Suppose that conditions 19.0.1 hold and $\dot{\Phi}_3^+$ is a union of tight components. If $\Delta(\alpha, \beta) > 5$, then $\Delta(\alpha, \beta) = 6$. Further, one of the following two situations occurs.*

- (i) $\bar{\Gamma}_S$ is hexagonal and its edges have weight m .
- (ii) $m = 4$, $\bar{\Gamma}_S$ is rectangular and its edges have weight 6.

Proof. Since $\mu(v) = m\Delta(\alpha, \beta) - 4$ for each vertex v of $\bar{\Gamma}_S$ (Lemma 19.8), Proposition 12.2 implies that the valency of each vertex of $\bar{\Gamma}_S$ is at most 6. First we show that the vertices of $\bar{\Gamma}_S$ have valency 4 or 6.

Let v be a vertex of $\bar{\Gamma}_S$. Inequality 19.2.1 shows that $\text{valency}_{\bar{\Gamma}_S}(v) \geq 4$. Suppose that $\text{valency}_{\bar{\Gamma}_S}(v) = 5$. By Proposition 12.2, $\varphi_3(v) = 3$. Then Lemma 19.7 implies that the sum of the weights of the edges incident to v is at most $5m + 2$. As this sum is the valency of v in Γ_S , we have $6m \leq \Delta(\alpha, \beta) \leq 5m + 2$, which is impossible since $m \geq 4$. Thus no vertex of $\bar{\Gamma}_S$ has valency 5, so each vertex v either has valency 4 and $\varphi_3(v) = 0$ or valency 6 and $\varphi_3(v) = 6$ (Proposition 12.2). In particular, no edge connects a vertex of valency 4 with one of valency 6. It follows that the union of the open star neighbourhoods of the vertices of valency 6 equals the union of the closed star neighbourhoods of these vertices. Thus this union is either \hat{F} or empty. It follows that either each vertex of $\bar{\Gamma}_S$ has valency 6, so $\bar{\Gamma}_S$ is hexagonal (Proposition 11.5), or each has valency 4. In the latter case there are no triangle faces so $\bar{\Gamma}_S$ is rectangular by Proposition 11.5.

Suppose that $\bar{\Gamma}_S$ is rectangular. Then inequality 19.2.1 shows that $m = 4$, $\Delta(\alpha, \beta) = 6$, and therefore each edge of $\bar{\Gamma}_S$ has weight 6. This is case (ii) of the lemma.

Suppose next that $\bar{\Gamma}_S$ is hexagonal. As each of its faces is a triangle, they lie on the $+$ -side of F (Lemma 19.4). Hence each edge of $\bar{\Gamma}_S$ has even weight. Suppose that some such edge \bar{e} has weight $m + 2$. Lemma 19.7 implies that if f is a face of $\bar{\Gamma}_S$ incident to \bar{e} , then each of the two edges of $\partial f \setminus \bar{e}$ has weight 2. Thus there is a vertex of $\bar{\Gamma}_S$ having successive edges of weight 2 incident to it. But then the remaining four edges have weights adding to at least $m\Delta(\alpha, \beta) - 4 \geq 6m - 4$. On the other hand, Proposition 12.2 shows that the maximal weights of these four edges are either m, m, m, m , or $2, m, m, m + 2$, or $2, 2, m + 2, m + 2$. Each possibility implies that $m < 4$. Thus each edge of $\bar{\Gamma}_S$ has weight m or less. Then $6m \leq m\Delta(\alpha, \beta) \leq 6m$. It follows that each edge of $\bar{\Gamma}_S$ has weight m and $\Delta(\alpha, \beta) = 6$. This is case (i). \diamond

Lemma 19.10. *Suppose that conditions 19.0.1 hold and $\dot{\Phi}_3^+$ is a union of tight components. If $\Delta(\alpha, \beta) = 6$, then $m = 4$.*

Proof. By Lemma 19.9 we can suppose that $\bar{\Gamma}_S$ is hexagonal and each of its edges has weight m . Proposition 11.5 implies that it has positive edges. Thus $m \leq \frac{m}{2} + 4$ (Lemma 19.5), so $m \leq 8$.

There are negative edges in $\bar{\Gamma}_S$ (Proposition 11.2) so we can choose a triangle face f of $\bar{\Gamma}_S$ with edges $\bar{e}_1, \bar{e}_2, \bar{e}_3$ where \bar{e}_1 is positive and \bar{e}_2, \bar{e}_3 are negative. Let e_1, e_2, e_3 be the edges of Γ_S incident to f and contained, respectively, in $\bar{e}_1, \bar{e}_2, \bar{e}_3$. Let v_1 be the vertex of Γ_S determined

by e_1 and e_2 , v_2 that determined by e_1 and e_3 , and v_3 that determined by e_2 and e_3 . We can suppose that e_1 has label 1 at the vertex v_1 and e_2 has label m there.

Suppose $m = 8$. Since there are no doubly-extended S -cycles in Γ_S , the permutation associated to any positive edge is of the form $i \mapsto 5 - i \pmod{8}$. As \bar{e}_1 is positive, e_1 has label 4 at v_2 , so e_3 has label 5 there. Then f yields the relation $\bar{x}_5 \bar{x}_1 \bar{x}_j = 1$. But the fourth bigon from f in the \bar{e}_1 family of edges implies that $\bar{x}_5 \bar{x}_1 = 1$. Thus $\bar{x}_j = 1$, which is impossible. Thus $m \neq 8$.

Suppose then that $m = 6$. As \bar{e}_1 is positive and Γ_S has no doubly-extended S -cycles, e_1 has label 2 or 4 at v_2 . We will deal with the first case as the second is similar. Thus e_1 has label 2 at v_2 so e_3 has label 3 there. The label of e_2 at v_3 cannot be 6 by Lemma 19.6 and the same lemma shows that it cannot be 2 as otherwise the label of e_3 at v_3 would be 3. Hence this label must be 4. Examination of the labels of the Γ_S -edges in \bar{e}_2, \bar{e}_3 at v_3 shows that $b_1 \cup b_3 \cup b_5$ and $b_2 \cup b_4 \cup b_8$ lie in components of $\dot{\Phi}_5^-$. But consideration of the lead edge of \bar{e}_1 at f shows that $b_1 \cup b_2$ lie in the same component of $\dot{\Phi}_5^-$. Thus $\dot{\Phi}_5^- = \tau_-(\dot{\Phi}_3^+)$ is connected, contrary to Proposition 9.4. Thus $m \neq 6$, which completes the proof of the lemma. \diamond

The previous two lemmas reduce the proof of Theorem 2.7 under assumption 19.0.1 to the cases described in the following two subsections.

19.2.1. *The case $m = 4$, $\Delta(\alpha, \beta) = 6$ and $\bar{\Gamma}_S$ hexagonal with edges of weight 4.* We consider singular disks D in X^+ , with $D \cap \partial X^+ = \partial D$. We can assume the components of ∂F are labeled so that $\partial X^+ = F \cup A_{23} \cup A_{41}$, where A_{23} and A_{41} are annuli running between boundary components 2,3 and boundary components 4,1 of F , respectively. By a homotopy we may assume that ∂D meets each of A_{23} and A_{41} in a finite disjoint union of essential embedded arcs. We will refer to these arcs as the *corners* of D . More precisely, if we go around ∂D in some direction we get a cyclic sequence of $X_2^{\pm 1}$ and $X_4^{\pm 1}$ -corners, where X_2, X_2^{-1} indicate that ∂D is running across A_{23} from 2 to 3 or from 3 to 2, respectively, and X_4, X_4^{-1} indicate that ∂D is running across A_{41} from 4 to 1 or 1 to 4, respectively. In this way D determines a cyclic word $W = W(X_2^{\pm 1}, X_4^{\pm 1})$, well-defined up to inversion, and we say that D is of type W . (Thus D is of type W if and only if it is of type W^{-1} .) We emphasize that W is an unreduced word; for example X_2 and $X_2 X_4 X_4^{-1}$ are distinct.

Let $\hat{A}_- \subseteq \hat{F}$ be the annulus defined at the beginning of §19.1. If $\partial D \cap F \subseteq A_-$ we say that D is an A_- -disk.

Recall the elements x_2, x_4 of $\pi_1(\hat{X}^+)$ as defined at the beginning of §19.1. We use x_j , respectively \bar{x}_j , to denote the image of x_j in $\pi_1(\hat{X}^+)$, respectively $\pi_1(\hat{X}^+(\alpha_-))$. Clearly, if D is an A_- -disk of type $W(X_2^{\pm 1}, X_4^{\pm 1})$ then D gives the relation $W(\bar{x}_2^{\pm 1}, \bar{x}_4^{\pm 1})$ in $\pi_1(\hat{X}^+(\alpha_-))$.

Note that a triangle face of Γ_S defines an A_- -disk. Note also that there is a one-one correspondence between triangle faces of Γ_S and faces of the reduced graph $\bar{\Gamma}_S$. We therefore say that a face of $\bar{\Gamma}_S$ has type W if and only if the corresponding triangle face of Γ_S has type W .

Let v be a vertex of Γ_S . An endpoint at v of an edge of the reduced graph $\bar{\Gamma}_S$ corresponds to four endpoints of edges of Γ_S , and we can assume that the label sequence (reading around v

anticlockwise if v is positive and clockwise if v is negative) is either 3 4 1 2 or 1 2 3 4. We say that v is of *type I* or *II*, respectively. If v is a positive vertex of type I we will say v is a $(+, I)$ *vertex*, and so on.




FIGURE 3.




FIGURE 4.

Lemma 19.7(1) implies

Lemma 19.11. *No edge of $\bar{\Gamma}_S$ connects vertices of the same type and opposite sign.*

By Proposition 11.2 Γ_S has the same number of positive and negative vertices. In particular, $\bar{\Gamma}_S$ has a face \bar{f}_1 in which not all vertices have the same sign; without loss of generality we may assume that two of the vertices are positive and one negative, and that the negative vertex is of type I. It follows from Lemma 19.11 that the two positive vertices are of type II. Thus the face \bar{f}_1 has type $X_2^{-1}X_4^2$. The configuration $C1$ of Γ_S corresponding to \bar{f}_1 is shown in Figure 3.

Lemma 19.12. $\bar{\Gamma}_S$ has a face of at most one of the types X_4^3 , $X_2X_4^2$, $X_2^2X_4$.

Proof. We consider the relations in $\pi_1(\widehat{X}^+(\alpha_-))$ coming from the corresponding triangle faces of Γ_S . A face of type X_4^3 , $X_2X_4^2$ or $X_2^2X_4$ would give the relation $\bar{x}_4^3 = 1$, $\bar{x}_2\bar{x}_4^2 = 1$ or $\bar{x}_2^2\bar{x}_4 = 1$, respectively. It is easy to see that any two of these, together with the relation $\bar{x}_2 = \bar{x}_4^2$ coming from f_1 , imply $\bar{x}_2 = \bar{x}_4 = 1$, contradicting Lemma 19.3. \diamond

Let $C2$ be the configuration shown in Figure 4.

Proposition 19.13. Γ_S contains the configuration $C2$.

We will assume that Γ_S does not contain such a configuration, and show that this leads to a contradiction. Equivalently, we make the following assumption:

$$(19.2.2) \quad \bar{\Gamma}_S \text{ contains no face with two } (-, I) \text{ vertices and one } (+, II) \text{ vertex.}$$

Let \mathcal{F}_1 be the set of faces of $\bar{\Gamma}_S$ with two $(+, II)$ vertices and one $(-, I)$ vertex, and let \mathcal{F}_2 be the set of faces with three positive vertices, at least two of which are of type II. Note that $\bar{f}_1 \in \mathcal{F}_1$. Let $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2$.

Lemma 19.14. Every face of $\bar{\Gamma}_S$ that shares an edge with a face in \mathcal{F} belongs to \mathcal{F} .

Proof. Let \bar{f} be a face in \mathcal{F} , let \bar{g} be a face of $\bar{\Gamma}_S$ that shares an edge \bar{e} with \bar{f} , and let v be the vertex of \bar{g} that is not a vertex of \bar{f} .

Case (1). $\bar{f} \in \mathcal{F}_1$.

First suppose that the vertices at the endpoints of \bar{e} are $(+, II)$ and $(-, I)$. If v is negative then by Lemma 19.11 it is a $(-, I)$ vertex, contradicting assumption (19.2.2). Therefore v is positive. By Lemma 19.11 it is a $(+, II)$ vertex. Hence $\bar{g} \in \mathcal{F}_1$.

Now suppose that \bar{e} connects the two $(+, II)$ vertices of \bar{f} . If v is negative then by Lemma 19.11 it is of type I, and hence $\bar{g} \in \mathcal{F}_1$. If v is positive then $\bar{g} \in \mathcal{F}_2$.

Case (2). $\bar{f} \in \mathcal{F}_2$.

First suppose that \bar{e} connects two vertices of type II. If v is negative then by Lemma 19.11 it is of type I so $\bar{g} \in \mathcal{F}_1$. If v is positive then $\bar{g} \in \mathcal{F}_2$.

If \bar{e} connects a $(+, I)$ vertex and a $(+, II)$ vertex then v is positive by Lemma 19.11. Also, \bar{f} is of type $X_2X_4^2$, so by Lemma 19.12 \bar{g} is also of type $X_2X_4^2$, and hence $\bar{g} \in \mathcal{F}_2$. \diamond

Now we prove Proposition 19.13. Lemma 19.14 implies that every face of $\bar{\Gamma}_S$ has at least two positive vertices. But this is easily seen to contradict the fact that Γ_S has the same number of positive and negative vertices. We conclude that assumption (19.2.2) is false, i.e. Proposition 19.13 holds. \diamond

If W is a word in $X_2^{\pm 1}$ and $X_4^{\pm 1}$ we denote by $\varepsilon_{X_2}(W)$ and $\varepsilon_{X_4}(W)$ the exponent sum in W of X_2 and X_4 respectively, and if D is a disk in X^+ of type W then we define $\varepsilon_{X_2}(D) = \varepsilon_{X_2}(W)$, $\varepsilon_{X_4}(D) = \varepsilon_{X_4}(W)$.

A disk in X^+ with 1, 2 or 3 corners will be called a *monogon*, *bigon* or *trigon*, respectively.

Lemma 19.15. *There are no monogons.*

Proof. Let D be a monogon. Applying the Loop Theorem to D , among disks with $\varepsilon_{X_2} + \varepsilon_{X_4} \neq 0$, we get an embedded monogon D' of the same type as D . Then D' is a boundary compressing disk for F in M , contradicting the fact that F is essential. \diamond




FIGURE 5.

Lemma 19.16. *There is no A_- -trigon of type W with $|\varepsilon_{X_2}(W)| + |\varepsilon_{X_4}(W)| = 1$.*

Proof. Such a disk would give rise to the relation $\bar{x}_2 = 1$ or $\bar{x}_4 = 1$ in $\pi_1(\widehat{X}^+(\alpha_-))$, contradicting Lemma 19.3. \diamond

Let D be a singular disk in X^+ . We say that an embedded disk E is *nearby* D if ∂E is contained in a small regular neighborhood of ∂D in ∂X^+ .

Lemma 19.17. *If there is an A_- -trigon of type W then there is a nearby embedded A_- -trigon of type W if $W = X_2^{\pm 3}$ or $X_4^{\pm 3}$ and of type W or $W^* = W(X_2^{-1}, X_4)$ otherwise.*

Proof. After possibly interchanging X_2 and X_4 we may assume without loss of generality that $\varepsilon_{X_2}(W) \not\equiv 0 \pmod{2}$. Let D be an A_- -trigon of type W . The Loop Theorem gives an embedded A_- -disk D' with $\varepsilon_{X_2}(D') \not\equiv 0 \pmod{2}$. Lemmas 19.15 and 19.16 now show that D' is of the desired type. \diamond




FIGURE 6.

Let D_1, D_2 be properly embedded disks in X^+ . Putting D_1 and D_2 in general position, $D_1 \cap D_2$ will be a compact 1-manifold. A standard cutting and pasting argument allows us to eliminate the circle components of $D_1 \cap D_2$, without changing ∂D_1 and ∂D_2 . So suppose that $D_1 \cap D_2$ consists of $n \geq 1$ arcs. Let u be one of these arcs. Then u cuts D_i into disks $D'_i, D''_i, i = 1, 2$, and the endpoints of u cut ∂D_1 and ∂D_2 into pairs of arcs α, β and γ, δ respectively; see Figure 5.

Cutting and pasting D_1 and D_2 along u we get four disks $D'_1 \cup D'_2, D'_1 \cup D''_2, D''_1 \cup D'_2,$ and $D''_1 \cup D''_2$, with boundaries $\alpha\gamma^{-1}, \alpha\delta, \beta\gamma$ and $\beta\delta^{-1}$ respectively. See Figure 6.

After a small perturbation, each of these disks E meets each of D_1 and D_2 in less than n double arcs, disjoint from the singularities of E .

The arc u is *trivial* in D_1 if either α or β contains no corner of D_1 , and similarly for D_2 . If u is trivial in D_1 and in D_2 then without loss of generality α contains no corner of D_1 and γ contains no corner of D_2 . Then $D'_1 \cup D'_2$ has the same type as D_1 and $D'_1 \cup D''_2$ has the same type as D_2 . If u is trivial in D_1 but not in D_2 , and D_2 is a bigon or trigon, then at least one of $D'_1 \cup D'_2, D'_1 \cup D''_2, D''_1 \cup D'_2,$ or $D''_1 \cup D''_2$ is a monogon, contradicting Lemma 19.15.

Lemma 19.18. *If there is no A_- -disk of type X_4^3 then there are disjoint embedded A_- -disks of types $X_2^{-1}X_4^2$ and $X_2^{-2}X_4$.*

Proof. The faces f_1 and f_2 of Γ_S in Figures 3 and 4 are A_- -disks of types $X_2^{-1}X_4^2$ and $X_2^{-2}X_4$ respectively. Since $\bar{x}_2^{-1}\bar{x}_4^2 = \bar{x}_2^{-2}\bar{x}_4 = 1$ in $\pi_1(\widehat{X}^+(\alpha_-))$, Lemma 19.3 implies that neither relation $\bar{x}_2\bar{x}_4^2 = 1$ nor $\bar{x}_2^2\bar{x}_4 = 1$ can hold. Lemma 19.17 then gives embedded A_- -disks D_1 and D_2 of type $X_2^{-1}X_4^2$ and $X_2^{-2}X_4$ respectively, which we may assume intersect in double arcs, none of which is trivial in D_1 or D_2 . Let u be a double arc. Ignoring orientations, there are two possibilities for u in each of D_1 and D_2 , shown in Figures 7 and 8 respectively.

Orient u as shown in Figures 7 and 8, so in case (1), $(\alpha, \beta) = (X_4^2, X_2^{-1})$, and in case (2), $(\alpha, \beta) = (X_4, X_4X_2^{-1})$. (Here, we are using the natural convention of labeling the oriented arc α or β by the sequence of corners it contains.) If u is oriented on D_2 as shown in Figure 5,




FIGURE 7.




FIGURE 8.

then in case (1), $(\gamma, \delta) = (X_2^{-2}, X_4)$, and in case (2), $(\gamma, \delta) = (X_2^{-1}, X_2^{-1}X_4)$. Note that if u on D_2 is oriented in the opposite direction to that shown then γ and δ are interchanged, so that in case $(\bar{1})$, $(\gamma, \delta) = (X_4, X_2^{-2})$, and in case $(\bar{2})$, $(\gamma, \delta) = (X_2^{-1}X_4, X_2^{-1})$. This gives eight possibilities (i, j) , where i denotes case (i) for D_1 , $i = 1, 2$, and j denotes case (j) for D_2 , $j = 1, 2, \bar{1}$ or $\bar{2}$. In each case we choose one of the four disks obtained by cutting and pasting

along u . Below we indicate the chosen disk by the arcs in its boundary and record its type:

$$\begin{aligned}
(1, 1) &: \alpha\delta & X_4^3 \\
(1, 2) &: \beta\delta^{-1} & X_2^{-1}X_4^{-1}X_2 \\
(2, 1) &: \beta\delta^{-1} & X_4X_2^{-1}X_4^{-1} \\
(2, 2) &: \alpha\delta & X_2^{-1}X_4^2 \\
(1, \bar{1}) &: \alpha\gamma^{-1} & X_4^2X_4^{-1} \\
(1, \bar{2}) &: \alpha\delta & X_2^{-1}X_4^2 \\
(2, \bar{1}) &: \beta\gamma & X_2^{-1}X_4^2 \\
(2, \bar{2}) &: \alpha\gamma^{-1} & X_4X_4^{-1}X_2
\end{aligned}$$

In case (1, 1) we get an A_- -disk of type X_4^3 , contradicting our assumption.

Cases (1, 2), (2, 1), (1, $\bar{1}$) and (2, $\bar{2}$) contradict Lemma 19.16.

In the remaining three cases, (2, 2), (1, $\bar{2}$) and (2, $\bar{1}$) we get an A_- -disk E of type $X_2^{-1}X_4^2$. By Lemma 19.17 there is a nearby embedded A_- -disk E' of type $X_2X_4^2$ or $X_2^{-1}X_4^2$. The former is impossible as otherwise we would have $\bar{x}_2\bar{x}_4^2 = \bar{x}_2^{-1}\bar{x}_4^2 = \bar{x}_2^{-2}\bar{x}_4 = 1$ in $\pi_1(\widehat{X}^+(\alpha_-))$, which implies $\bar{x}_2 = \bar{x}_4 = 1$, contrary to Lemma 19.3. Thus E' has type $X_2^{-1}X_4^2$. Noting that $|E' \cap D_2| \leq |E \cap D_2| < |D_1 \cap D_2|$, if we continue in this manner we eventually get an embedded A_- -disk of type $X_2^{-1}X_4^2$ disjoint from D_2 . \diamond

Note. It is easy to see that in cases (2, 2), (1, $\bar{2}$) and (2, $\bar{1}$) we also get a disk of type $X_2^{-2}X_4$, so we could equally well have fixed D_1 and obtained an embedded A_- -disk of type $X_2^{-2}X_4$ disjoint from D_1 .

Let D be an embedded disk in X^+ . Recall that the corners of D are the components of $\partial D \cap (A_{23} \cup A_{41})$. We will refer to the components of $\partial D \cap F$ as the *edges* of D . We label the endpoints of the edges of D with the label of the corresponding corner. Thus, if we have a disjoint union Δ of embedded disks whose $X_2^{\pm 1}$ -corners are labeled so that reading clockwise around boundary component 2 of F they appear in the order a, b, c, \dots , then they appear in the same order a, b, c, \dots reading anticlockwise around boundary component 3 of F . Similarly, the clockwise order of the $X_4^{\pm 1}$ -corners of Δ at boundary component 4 is the same as their anticlockwise order at boundary component 1. This ordering condition puts constraints on the existence of the disjoint embedded arcs in F that are the edges of Δ .

Lemma 19.19. *If there is an A_- -disk of type X_4^3 then $\alpha_- = \varphi_+$.*

Proof. If there is an A_- -disk of type X_4^3 then there is an embedded A_- -disk D of type X_4^3 by Lemma 19.17. Let the corners of D be a, b and c ; see Figure 9.

Then without loss of generality the edges of D appear in \widehat{A}_- as shown in Figure 10.

Let $V = \widehat{A}_- \times I \cup H_{(41)} \cup N(D) \subseteq \widehat{X}^+$. Then, taking as ‘‘base-point’’ a disk in \widehat{A}_- containing the two left-hand edges in Figure 10 together with fat vertices v_1 and v_4 , we get $\pi_1(V) \cong \langle x_4, t : x_4^3 = t \rangle \cong \mathbb{Z}$, where t is represented by α_- , the core of \widehat{A}_- . Hence V is a solid torus and α_- has winding number 3 in V . Let A' be the annulus $\partial V - \text{int } \widehat{A}_-$. By Assumption 2.2, the torus




FIGURE 9.




FIGURE 10.

$(\widehat{F} - \widehat{A}_-) \cup A'$ bounds a solid torus V' in \widehat{X}^+ . Therefore $\widehat{X}^+ = V \cup_{A'} V'$ is a Seifert fibre space with base orbifold $D^2(3, b)$, and α_- is the slope of the Seifert fibre φ_+ . \diamond

Lemma 19.20. *If there are disjoint embedded A_- -disks of types $X_2^{-1}X_4^2$ and $X_2^{-2}X_4$ then $\alpha_- = \varphi_+$.*

Proof. Let D_1, D_2 be disjoint embedded A_- -disks of types $X_2^{-1}X_4^2$ and $X_2^{-2}X_4$ respectively.

First note that the union of the edges of D_1 and D_2 and the fat vertices v_1, v_2, v_3, v_4 cannot be contained in a disk in \widehat{A}_- . For this would give relations $x_2 = x_4^2, x_4 = x_2^2$ in $\pi_1(\widehat{X}^+)$, implying $x_2^3 = x_4^3 = 1$. But x_2 and x_4 are non-trivial (Lemma 19.3), so $\pi_1(\widehat{X}^+)$ would have non-trivial torsion, contradicting the fact that \widehat{X}^+ is a Seifert fibre space with base orbifold $D^2(2, b)$.

Let the corners of D_1 and D_2 be a, b, c and p, q, r respectively; see Figure 11.




FIGURE 11.

Without loss of generality the labels c, p, q appear in this order anticlockwise around v_3 . The possible arrangements of the edges of D_1 and D_2 in \hat{A}_- are then shown in Figure 12 (1)–(6). (For simplicity we have labeled the corners a, b, c, p, q, r only in Figure 12 (1).)




FIGURE 12.

Let $V = \widehat{A}_- \times I \cup H_{(23)} \cup H_{(41)} \cup N(D_1) \cup N(D_2) \subseteq \widehat{X}^+$. Then $\pi_1(V)$ is generated by x_2, x_4, t , where t is represented by α_- , the core of \widehat{A}_- . We take as “base-point” a disk in \widehat{A}_- containing the edges of D_1 together with the vertices v_1, v_2, v_3, v_4 . Then the disk D_1 gives the relation $x_2 = x_4^2$ in $\pi_1(V)$. The relation determined by D_2 is as follows in cases (1)–(6):

- (1) $x_2^{-1} t x_2^{-1} x_4 = 1$
- (2) $x_2^{-1} t x_2^{-1} x_4 t = 1$
- (3) $x_2^{-1} t x_2^{-1} t x_4 = 1$
- (4) $x_2^{-1} t x_2^{-1} t x_4 t = 1$
- (5) $x_2^{-2} t x_4 = 1$
- (6) $x_2^{-2} x_4 t = 1$

In case (1) we get $x_4 = x_2 t^{-1} x_2 = x_4^2 t^{-1} x_4^2$, and hence $t = x_4^3$. Therefore $\pi_1(V) \cong \mathbb{Z}$, generated by x_4 . It follows that V is a solid torus and α_- has winding number 3 in V . Hence (see the proof of Lemma 19.19) $\alpha_- = \varphi_+$.

In case (2) $x_4 = (x_2 t^{-1})^2 = z^2$, where $z = x_2 t^{-1}$. Thus $\pi_1(V)$ is generated by x_2, x_4, z with relations $x_2 = x_4^2, x_4 = z^2$. Therefore $\pi_1(V) \cong \mathbb{Z}$, generated by z . Also $t = z^{-1} x_2 = z^{-1} z^4 = z^3$. Hence again $\alpha_- = \varphi_+$, as in case (1).

Cases (3), (5) and (6) are similar and are left to the reader.

In case (4) we have $x_4 = t^{-1} x_2 t^{-1} x_2 t^{-1}$, so $x_4 x_2 = z^3$, where $z = t^{-1} x_2$. Since $x_2 = x_4^2$, $\pi_1(V)$ has presentation $\langle x_4, z : x_4^3 = z^3 \rangle$. But this contradicts the fact that \widehat{X}^+ is a Seifert fibre space with base orbifold $D^2(2, b)$. \diamond

Corollary 19.21. $\alpha_- = \varphi_+$.

Proof. This follows from Lemmas 19.18, 19.19 and 19.20. \diamond

We complete the analysis by showing that Corollary 19.21 implies that $\dot{\Phi}_3^+$ contains an \widehat{F} -essential annulus, contrary to our assumptions.

Lemma 19.22. *There is no pair of disjoint embedded A_- -disks of types $X_2^{-1} X_4^2$ and $X_2^{-2} X_4$.*

Proof. The manifold V given in the proof of Lemma 19.20 is a solid torus such that \widehat{A}_- is contained in ∂V with winding number 3 and thus the annulus $A = \partial V \setminus \widehat{A}_-$ is a vertical annulus in the Seifert fibred structure of \widehat{X}^+ . But A is contained in X^+ and thus it is an essential annulus in X^+ . So $\partial A = \partial \widehat{A}_-$ can be isotoped in F into the interior of $\dot{\Phi}_1^+$. Therefore $\dot{\Phi}_1^+ \cap A_- = \dot{\Phi}_1^+ \cap \tau_-(\dot{\Phi}_1^+) = \dot{\Phi}_1^+ \cap \dot{\Phi}_2^-$ is a pair of \widehat{F} -essential annulus components. Hence $\dot{\Phi}_3^+$ is a pair of \widehat{F} -essential annulus components. But this contradicts our assumption that $\dot{\Phi}_3^+$ is a set of tight components. Thus there is no such pair of embedded disks. \diamond

Thus the situation given by Lemma 19.20 cannot arise. Then by Lemma 19.18, there is an A_- -disk of type X_4^3 . We also have a A_- disk of type $X_2^{-1} X_4^2$ given by configuration $C1$. By Lemma 19.17, there is an embedded A_- -disks D_1 of type X_4^3 and another D_2 of type $X_2^{-1} X_4^2$

or $X_2X_4^2$. In the latter case, we have the relations $\bar{x}_4^3 = \bar{x}_2^{-1}\bar{x}_4^2 = \bar{x}_2\bar{x}_4^2 = 1$ in $\pi_1(\widehat{X}^+(\alpha_-))$, which imply that $\bar{x}_2 = \bar{x}_4 = 1$, contrary to Lemma 19.3. Thus D_2 has type $X_2^{-1}X_4^2$.

Lemma 19.23. *There are disjoint embedded A_- -disks D_1 and D_2 of types X_4^3 and $X_2^{-1}X_4^2$ respectively.*

Proof. The proof is similar to that of Lemma 19.18. We may assume that among all such pairs of embedded A_- -disks of types X_4^3 and $X_2^{-1}X_4^2$ respectively, D_1 and D_2 have been chosen to have the minimal number of intersection components. If the disks D_1 and D_2 are disjoint then we are done. So suppose they intersect. We may assume that they intersect transversely in double arcs, none of which is trivial in D_1 or D_2 .

Let u be an oriented double arc which is outermost in D_1 with respect to the corner it cuts off (i.e. the interior of the corner is disjoint from D_2) as shown in Figure 13 (a). Then there are six possibilities for the oriented arc u in D_2 , as shown in Figure 13 (b1)-(b6) respectively.

If case (b1) of Figure 13 occurs, then cutting and pasting D_1 and D_2 will produce an embedded A_- -disk of type $X_2^{-1}X_4^2$ having fewer intersection components with D_1 than does D_2 , contradicting our assumption on D_1 and D_2 .

If case (b2) of Figure 13 occurs, then cutting and pasting will produce an A_- -disk of type X_4^2 . So in $\pi_1(\widehat{X}^+(\alpha_-))$ we have the relation $\bar{x}_4^2 = 1$. Together with the relations $\bar{x}_4^3 = 1$ and $\bar{x}_2^{-1}\bar{x}_4^2 = 1$ this implies that $\bar{x}_2 = \bar{x}_4 = 1$ in $\pi_1(\widehat{X}^+(\alpha_-))$, a contradiction.

Cases (b3) and (b4) of Figure 13 can be treated similarly to the cases (b1) and (b2) respectively.

If case (b5) of Figure 13 occurs, then cutting and pasting D_1 and D_2 will produce an A_- -disk of type $X_2^{-1}X_4$. Thus in $\pi_1(\widehat{X}^+(\alpha_-))$ we have $\bar{x}_2^{-1}\bar{x}_4 = 1$. Since $\bar{x}_4^3 = 1$ and $\bar{x}_2^{-1}\bar{x}_4^2 = 1$, we deduce $\bar{x}_2 = \bar{x}_4 = 1$ in $\pi_1(\widehat{X}^+(\alpha_-))$, a contradiction.

Finally, if case (b6) of Figure 13 occurs, then cutting and pasting will produce an embedded A_- -disk of type X_4^3 which is disjoint from D_2 , giving an obvious contradiction. \diamond

Now let V be a regular neighborhood in \widehat{X}^+ of the set $\widehat{A}_- \cup H_{(23)} \cup H_{(41)} \cup D_1 \cup D_2$. As in the proof of Lemma 19.20, the union of the edges of D_1 and the fat vertices v_1 and v_4 cannot lie in a disk in \widehat{A}_- and can be assumed to appear as shown in Figure 10. Thus two edges of D_1 connect the fat vertices v_1 and v_4 from the left hand side and one edge of D_1 connects v_1 and v_4 from the right hand side.

Let e_1 be the edge of D_2 connecting v_1 and v_4 , e_2 the edge of D_2 connecting v_1 and v_3 , and e_3 the edge of D_2 connecting v_2 and v_4 .

Now take as ‘‘base-point’’ a disk in \widehat{A}_- containing the union of the two left-hand side edges of D_1 , the fat vertices v_1, v_2, v_3, v_4 , and the edges e_2, e_3 . Then the disk D_1 will give the relation

$$x_4^3 = t,$$

and the disk D_2 will give either the relation

$$x_2^{-1}x_4^2 = 1$$




FIGURE 13.

(when e_1 connects v_1 and v_4 from the left hand side, cf. Figure 10) or the relation

$$x_2^{-1}x_4t^{-1}x_4 = 1$$

(when e_1 connects v_1 and v_4 from the right hand side, cf. Figure 10), where t is represented by α_- . In either case we see that V has the fundamental group

$$\pi_1(V) = \langle x_4, t : x_4^3 = t \rangle.$$

So the manifold V is a solid torus such that \widehat{A}_- is contained in ∂V with winding number 3. Now argue as in the proof of Lemma 19.22 to see that Φ_3^+ cannot be a set of tight components, yielding the final contradiction. \diamond

19.2.2. *The case $m = 4$, $\Delta(\alpha, \beta) = 6$ and $\overline{\Gamma}_S$ rectangular with edges of weight 6.* As $m = 4$, we may assume:

- both $\dot{\Phi}_3^+$ and $\dot{\Phi}_5^-$ consist of a pair of tight components, each a twice-punctured disk;
- $\dot{\Phi}_5^+$ is a collar on ∂F , and so contains no large components.

Recall that b_1, \dots, b_4 denote the four boundary components ∂F appearing successively along ∂M . These four circles cut ∂M into four annuli $A_{i,i+1}$, $i = 1, \dots, 4$, such that $\partial A_{i,i+1} = b_i \cup b_{i+1}$ (indexed mod (4)). We may assume that $\partial X^+ = F \cup A_{2,3} \cup A_{4,1}$.

As in §19.2.1, an n -gon (disk) in X^+ means a singular disk D with $\partial D \subseteq \partial X^+$ such that $\partial D \cap (A_{2,3} \cup A_{4,1})$ is a set of n embedded essential arcs in $A_{2,3} \cup A_{4,1}$, called the *corners* of D , and $\partial D \cap F$ is a set of n singular arcs, called the *edges* of D . Recall that a 1-gon, 2-gon or 3-gon will be called a monogon, bigon or trigon.

There are no monogons in X^+ (cf. Lemma 19.15).

Lemma 19.24. *There is no bigon D in X^+ whose edges e_1, e_2 are essential paths in $(\dot{\Phi}_5^-, \partial F)$ and for which the inclusion $(D, e_1 \cup e_2) \rightarrow (X^+, \dot{\Phi}_5^-)$ is essential as a map of pairs.*

Proof. Suppose otherwise that such a bigon D exists. Then D gives rise to an essential homotopy between its two edges and thus the edges of D can be homotoped, relative to their end points, into $\dot{\Phi}_1^+$. Then the essential intersection $\dot{\Phi}_5^- \wedge \dot{\Phi}_1^+$ contains a large component and therefore so does $\dot{\Phi}_6^+ = \tau_+(\dot{\Phi}_5^- \wedge \dot{\Phi}_1^+)$, contrary to our assumption that $\dot{\Phi}_5^+$ has no large components. \diamond

Recall that h is the π_1 -injective map from the torus T into $M(\alpha)$ which induces the graph Γ_S in T . For a subset s of T we use s^* to denote its image under the map h .

The image under h of every edge of a rectangular face of Γ_S is contained in $\dot{\Phi}_5^-$. The images of the middle two edges of every family of six parallel edges of Γ_S are contained in A_- .

As before the classes $x_j \in \pi_1(\widehat{X}^+)$ are defined and we use \bar{x}_j to denote their images in $\pi_1(\widehat{X}^+(\alpha_-))$.

For notational simplicity, let us write $\dot{\Phi}_5^- = Q$, a pair of twice-punctured disks. A singular disk $D \subset X^+$ whose edges are contained in Q will be called a Q -disk. An *essential* Q -disk is a Q -disk D such that ∂D is essential in ∂X^+ . The following two lemmas are key to our analysis.

Lemma 19.25. *An essential Q - n -gon, $n \leq 4$, is a 4-gon of type $X_2X_4^{-1}X_4X_4^{-1}$ or $X_4X_2^{-1}X_2X_2^{-1}$.*

Lemma 19.26. *There cannot be essential Q -4-gons of both types $X_2X_4^{-1}X_4X_4^{-1}$ and $X_4X_2^{-1}X_2X_2^{-1}$.*

The proofs of these two lemmas will be given after we develop several necessary background results.

All the edges of $\bar{\Gamma}_S$ have weight 6. We may assume without loss of generality that there is a family of parallel edges of Γ_S at one end of which the label sequence is 1 2 3 4 1 2.

Lemma 19.27. *b_1 and b_2 belong to different components of Q .*

Proof. Suppose otherwise so that there is a rectangle face of Γ_S as depicted in Figure 14.




FIGURE 14.

Here $e_1, e_2, e_3, e_4, e_5, e_6$ is a family of six successive parallel edges which connect vertices v_1 and v_2 and whose label-permutation is the identity. Let R_i be the bigon face between e_i, e_{i+1} for $i = 1, \dots, 5$ and R the disk $R_1 \cup \dots \cup R_5$. We know $R_2^*, R_4^*, f^* \subseteq X^+$ while $R_1^*, R_3^*, R_5^* \subseteq X^-$.

There is a product structure $(R_i, e_i, e_{i+1}) = (e_i \times I, e_i \times \{0\}, e_i \times \{1\})$ such that for each $x \in e_i$, $(\{x\} \times I)^*$ is an I -fibre of $\Sigma_1^{(-1)^i}$. Thus $\tau_{(-1)^i}(e_i^*) = e_{i+1}^*$, so the free involution $h_5^- : \dot{\Phi}_5^- \rightarrow \dot{\Phi}_5^-$ (see the end of §3.2) sends $e_1^* \cup b_1$ to $e_6^* \cup b_2$. Proposition 4.5 then shows that b_1 and b_2 lie in different components of $\dot{\Phi}_5^-$. Hence b_3 and b_4 also lie in different components of $\dot{\Phi}_5^-$. \diamond

It follows that b_3 and b_4 also belong to different components of Q .

If $(D, \partial D) \subset (X^+, \partial X^+)$, we will denote the type of D (see §19.2.1) by $W(D)$. Recall that $W(D)$ is defined up to cyclic permutation and inversion.

Corollary 19.28. *Let D be a Q -disk. Then $W(D)$ does not contain the syllable X_2X_4 or X_4X_2 .*

Proof. These give rise to a 34- or 12-edge in ∂D , respectively. \diamond

Lemma 19.29. *Let D be a Q -disk. Then in $W(D)$ no $Z \in \{X_2^{\pm 1}, X_4^{\pm 1}\}$ can be followed or preceded by two distinct letters $\neq Z^{-1}$.*

Proof. If Z were followed by two distinct letters $\neq Z^{-1}$ the same component of Q would contain three boundary components b_i . For example, if $W(D)$ contained syllables X_2X_2 and $X_2X_4^{-1}$ then ∂D would contain a 32-edge and a 31-edge, implying that b_1, b_2 and b_3 belong to the same component of Q . \diamond

Proof of Lemma 19.25. Let E be an essential Q - n -gon, $n \leq 4$. By the Loop Theorem we get an essential embedded Q -disk D , with $\{\text{corners of } D\} \subset \{\text{corners of } E\}$.

Lemma 19.30. *D is a 4-gon.*

Proof. D cannot be a monogon, since then D would be a boundary-compressing disk for F .

D cannot be a bigon by Lemma 19.24.

So suppose D is a trigon. It is easy to see that Corollary 19.28 and Lemma 19.29 imply that D contains only, say, X_2 -corners. By Lemma 19.18 $|\varepsilon_{X_2}(D)| \neq 1$, so $W(D) = X_2^3$.

Let $U = \widehat{F} \times I \cup H_{(23)} \cup N(D) \subset \widehat{X}^+$. Then $\pi_1(U) \cong \pi_1(\widehat{F}) * \mathbb{Z}/3$. It follows that U , and hence \widehat{X}^+ , has a closed summand with fundamental group $\mathbb{Z}/3$, a contradiction. \diamond

There are three possibilities: D has either

- (A) all X_2 -corners (or all X_4 -corners);
- (B) two X_2 -corners and two X_4 -corners;
- (C) one X_2 -corner and three X_4 -corners (or vice versa).

Lemma 19.31. *Case (A) is impossible.*

Proof. We may suppose that D has all X_2 -corners. Note that $|\varepsilon_{X_2}(D)|$ is not 1 by Lemma 19.18 and if it is > 1 then we get a contradiction as in the last part of the proof of Lemma 19.30. Hence $\varepsilon_{X_2}(D) = 0$. Thus $W(D) = X_2^2 X_2^{-2}$ or $X_2 X_2^{-1} X_2 X_2^{-1}$.

In the first case, ∂D contains a 23-edge, and hence b_2 and b_3 belong to the same component of Q . But ∂D also contains a 2-loop and a 3-loop, which clearly must intersect, contradicting the fact that D is embedded.

In the second case, label the corners of D a, b, c, d as shown in Figure 15.




FIGURE 15.

Then ∂D is as shown in Figure 16. Let $V = \widehat{F} \times I \cup H_{(23)}$. Note that $\partial V = \widehat{F} \times \{0\} \amalg G$, where G is a surface of genus 2. We see from Figure 16 that ∂D is isotopic in G to a meridian of $H_{(23)}$, and so bounds a non-separating disk $D' \subset V$. Then $D \cup D'$ is a non-separating 2-sphere $\subset V \cup N(D) \subset \widehat{X}^+$, a contradiction. \diamond

Lemma 19.32. *Case (B) is impossible.*

Proof. By Corollary 19.28 and Lemma 19.29, the only possibilities for $W(D)$ are $X_2 X_2^{-1} X_4 X_4^{-1}$ and $X_2 X_4^{-1} X_2 X_4^{-1}$.




FIGURE 16.

In the first case, ∂D contains a 24-edge. Therefore b_2 and b_4 belong to the same component of Q , and hence b_1 and b_3 belong to the same component of Q . But ∂D also contains a 1-loop and a 3-loop, which must intersect.

In the second case, let $U = \widehat{F} \times I \cup H_{(23)} \cup H_{(41)} \cup N(D) \subset \widehat{X}^+$. Then $\pi_1(U) \cong \pi_1(\widehat{F}) * \mathbb{Z} * \mathbb{Z}/2$, implying that \widehat{X}^+ has a closed summand with fundamental group $\mathbb{Z}/2$, a contradiction. \diamond

By Lemmas 19.31 and 19.32, Case (C) must hold; so suppose that D has one X_2 -corner and three X_4 -corners. Since $\{\text{corners of } D\} \subset \{\text{corners of } E\}$, E is also a 4-gon with one X_2 -corner and three X_4 corners. Corollary 19.28 rules out all possibilities for $W(E)$ except $X_2X_4^{-3}$ and $X_2X_4^{-1}X_4X_4^{-1}$, and the first is ruled out by Lemma 19.29.

This completes the proof of Lemma 19.25. \diamond

Lemma 19.33. *There do not exist disjoint Q -disks of types $X_2X_4^{-1}X_4X_4^{-1}$ and $X_4X_2^{-1}X_2X_2^{-1}$.*

Proof. Let D_1, D_2 be Q -disks of types $X_2X_4^{-1}X_4X_4^{-1}$ and $X_4X_2^{-1}X_2X_2^{-1}$, respectively. Since ∂D_1 contains a 31-edge, b_1 and b_3 must belong to the same component of Q . But ∂D_1 contains a 1-loop and ∂D_2 contains a 3-loop, and these must intersect. \diamond

Proof of Lemma 19.26. Let E_1, E_2 be Q -disks of types $X_2X_4^{-1}X_4X_4^{-1}$ and $X_4X_2^{-1}X_2X_2^{-1}$ respectively. By the Loop Theorem and Lemma 19.25 we get embedded Q -disks D_1 and D_2 of these types. By Lemma 19.33, D_1 and D_2 must intersect; consider an arc of intersection, coming from the identification of arcs $u_i \subset D_i$, $i = 1, 2$. We may assume that the endpoints of u_i lie on distinct edges of D_i , $i = 1, 2$. Then u_i separates D_i into two disks, D'_i and D''_i , say, where D'_i contains either one or two corners of D_i .

If D'_1 and D'_2 each contain a single corner, and these corners are distinct, then $D'_1 \cup D'_2$ is a Q -bigon with one X_2 - and one X_4 -corner, contradicting Lemma 19.24.

If D'_1 and D'_2 both contain, say, a single X_2 -corner, then u_1 is as shown in Figure 17, which also shows one of the three possibilities for u_2 . Since b_1 and b_3 lie in one component of Q , say Q_1 , and b_2 and b_4 lie in the other component, say Q_2 , and each of the arcs u_1 and u_2 has one endpoint in Q_1 and one in Q_2 , u_1 and u_2 must be identified as shown in Figure 17. Then $D_1^* = D'_1 \cup D'_2$ is a Q -disk of type $X_2X_4^{-1}X_4X_4^{-1}$ having fewer intersections than D_1 with D_2 .




FIGURE 17.




FIGURE 18.

If each of D'_i and D''_i contains two corners, $i = 1, 2$, the two possibilities for u_1 and u_2 are illustrated in Figure 18, (a) and (b). In both cases, $D_1^* = D'_1 \cup D'_2$ is again a Q -disk of type $X_2X_4^{-1}X_4X_4^{-1}$ having fewer intersections with D_2 .

Applying the Loop Theorem to the disk D_1^* constructed above, and using Lemma 19.25, we get an embedded Q -disk of type $X_2X_4^{-1}X_4X_4^{-1}$ having fewer intersections with D_2 than D_1 . Continuing, we eventually get disjoint embedded Q -disks of types $X_2X_4^{-1}X_4X_4^{-1}$ and $X_4X_2^{-1}X_2X_2^{-1}$, contradicting Lemma 19.33 .

This completes the proof of Lemma 19.26. \diamond

Since each edge of $\bar{\Gamma}_S$ has weight 6, consecutive 4-gon corners of Γ_S at a given vertex are distinct. Hence the total number of X_2 -corners in the 4-gon faces of Γ_S is the same as the total number of X_4 -corners. Since a 4-gon face of Γ_S is an essential Q -disk, this contradicts Lemmas 19.25 and 19.26.

This completes the proof for the case where $\bar{\Gamma}_S$ is rectangular.

19.3. Proof when $\dot{\Phi}_3^+$ is not a union of tight components. In this section we suppose that X^- is a twisted I -bundle and $\dot{\Phi}_3^+$ is not a union of tight components. Proposition 8.2 implies that

- $M(\beta)$ is Seifert with base orbifold $P^2(2, n)$ for some $n > 2$;
- \widehat{F} is vertical in $M(\beta)$;
- $\check{\Phi}_1^+$ is connected and completes to an \widehat{F} -essential annulus;
- $\check{\Phi}_3^+$ completes to the union of two \widehat{F} -essential annuli.

By Corollary 11.4, the edges of $\bar{\Gamma}_S$ have weight bounded above by $m + 4$. Hence for any vertex v of $\bar{\Gamma}_S$ we have

$$(19.3.1) \quad \Delta(\alpha, \beta) \leq \text{valency}_{\bar{\Gamma}_S}(v) + 4 \left(\frac{\text{valency}_{\bar{\Gamma}_S}(v)}{m} \right)$$

As the Seifert structure on \widehat{X}^+ is unique, it is the restriction of the Seifert structure of $M(\beta)$ and therefore its base orbifold is $D^2(2, n)$. Recall from §19.1 that ϕ_+ is the fibre slope on \widehat{F} of this structure. By hypothesis, it is also the fibre slope of the Seifert structure on \widehat{X}^- , a twisted I -bundle over the Klein bottle with base orbifold a Möbius band. Hence $\phi_+ = \tau_-(\phi_+) = \alpha_-$, so the class $t \in \pi_1(\widehat{F}) \leq \pi_1(\widehat{X}^+)$ is the fibre class.

Proposition 19.34. *Suppose that conditions 19.0.1 hold and $\dot{\Phi}_3^+$ is not a union of tight components. If $m = 4$ there is a presentation $\langle a, b, z : a^2, b^n, abz^{-2} \rangle$ of $\Gamma = \pi_1(P^2(2, n))$ such that the image in Γ of the core K_β of the β -filling solid torus in $M(\beta)$ represents the element $\kappa = az^{-1}b^{-1}z \in \Gamma$, at least up to conjugation and taking inverse.*

Proof. Let E_0 be the \widehat{F} -essential annulus $\check{\Phi}_1^+$. Then ∂E_0 is a pair of \widehat{F} -essential curves c_1, c_2 . By Proposition 8.2, $\dot{\Phi}_3^+$ is the union of two \widehat{F} -essential annuli, and there are disjoint, non-separating annuli A_1^-, A_2^- properly embedded in X^- such that $\partial A_1^- \cup \partial A_2^- \subseteq \dot{\Phi}_1^+$ and for each j , $\partial \dot{\Phi}_1^+ \cap \partial A_j^-$ is a boundary component of $\dot{\Phi}_1^+$ which we can take to be c_j .

We can assume that each A_j^- is τ_- -invariant. Then $A_1^- \cup A_2^-$ splits \widehat{X}^- into two τ_- -invariant solid tori V_1, V_2 where $V_1 \cap M \supset \partial M \cap X^-$ and $V_2 \subset M$. The reader will verify that $\check{\Phi}_1^+ \cap V_1$ is the union of disjoint \widehat{F} -essential annuli E_1, E_2 where $\tau_-(E_1) = E_2$ while $F \cap V_2$ is the union of disjoint \widehat{F} -essential annuli E_3, E_4 such that $\tau_-(E_3) = E_4$. Without loss of generality we can suppose that $c_j \subset E_j$ ($j = 1, 2$) and $\widehat{\Phi}_1^+ = E_1 \cup E_2 \cup E_3$. Then $\check{\Phi}_3^- = \tau_-(\check{\Phi}_1^+) = E_1 \cup E_2 \cup E_4$.

Number the components of ∂F so that $\partial M \cap X^+$ consists of two annuli, one with boundary $b_1 \cup b_4$, the other with boundary $b_2 \cup b_3$. Let $x = x_4$ and $y = x_2$ be the elements of $\pi_1(\widehat{X}^+) \leq \pi_1(M(\beta))$ defined using the disk $D \subset A_- = \check{\Phi}_3^-$.

The intersection of ∂M with X^- consists of two annuli, one with boundary $b_1 \cup b_2$ and the other with boundary $b_3 \cup b_4$. Let w_1, w_3 be the associated elements of $\pi_1(\widehat{X}^-) \leq \pi_1(M(\beta))$ determined by D . Since $V_1 \cap M$ is a twice-punctured annulus cross an interval we see that $w_1 = w_3^{\pm 1}$. We claim that $w_1 = w_3^{-1}$. To see this, exchange E_1 and E_2 , if necessary, so that $b_j \subset E_j$ for $j = 1, 2$. We will be done if $b_4 \in E_1$ and $b_3 \in E_2$. Suppose otherwise that $b_3 \in E_1$ and $b_4 \in E_2$. Then $\tau_+(E_1)$ is an \widehat{F} -essential annulus in E_0 containing $c_2 \cup b_2 \cup b_4$ while $\tau_+(E_2)$ is an \widehat{F} -essential annulus in E_0 containing $c_1 \cup b_1 \cup b_3$. It follows that $\partial\tau_+(E_1) \setminus c_2$ is an \widehat{F} -essential curve in $\check{\Phi}_1^+$ which separates $b_2 \cup b_4$ from $b_1 \cup b_3$. A similar conclusion holds for $\partial\tau_+(E_2) \setminus c_1$. It follows that up to isotopy we can assume $\tau_+(E_1 \cap F) = E_2 \cap F$. On the other hand, by construction we have $\tau_-(E_1 \cap F) = E_2 \cap F$ and therefore $(\tau_- \circ \tau_+)(E_1 \cap F) = E_1 \cap F$. Hence the inclusion of $E_2 \cap F$ in F admits essential homotopies of arbitrarily large length, contrary to the results of §10. Thus $w_1 = w_3^{-1}$. Let z be the image of w_1 in Γ .

The class of $\pi_1(M(\beta))$ carried by K_β is given by $xw_1yw_1^{-1}$. Let κ be its image in Γ .

The base orbifold of \widehat{X}^+ is $D^2(2, n)$ with fundamental group $\pi_1(D^2(2, n)) = \langle a, b : a^2 = 1, b^n = 1 \rangle$. Here a, b are chosen to be represented by oriented simple closed curves in the complement P of the cone points of $D^2(2, n)$.

We can assume that the E_i are vertical in the Seifert structure on $M(\beta)$. Since $D \subset E_1 \cup E_2 \cup E_4$, it projects to a proper subarc of the circle in $P^2(2, n)$ given by the image of the vertical torus \widehat{F} . Thus the images of x and y in $\pi_1(D^2(2, n))$ lie in $\{a^{\pm 1}, b^{\pm 1}\}$ (cf. the proof of Proposition 15.1). Further $z^2 \in \{ab, ab^{-1}, ba, b^{-1}a\} \subset \Gamma$. By construction $b_1 \cup b_4 \subset E_1$ and $b_2 \cup b_3 \subset E_2$ and so as w_1 is obtained by concatenating an arc in $\partial M \cap X^-$ from b_1 to b_2 with an arc in D from b_2 to b_1 , it follows that one of the following four possibilities arises:

- (1) $x \mapsto a, y \mapsto b, z^2 = ba$ and $\kappa = abz^{-1}$.
- (2) $x \mapsto a, y \mapsto b^{-1}, z^2 = b^{-1}a$ and $\kappa = azb^{-1}z^{-1}$.
- (3) $x \mapsto b, y \mapsto a, z^2 = ab$ and $\kappa = bza z^{-1}$.
- (4) $x \mapsto b^{-1}, y \mapsto a, z^2 = ab^{-1}$ and $\kappa = b^{-1}zaz^{-1}$.

In case (3) we have $\Gamma = \langle a, b, z : a^2, b^n, ab = z^2 \rangle$ where $\kappa = bza z^{-1} = z(az^{-1}b^{-1}z)^{-1}z^{-1}$. In case (4) we have $\Gamma = \langle a, b, z : a^2, b^n, ab^{-1} = z^2 \rangle$ where $\kappa = b^{-1}zaz^{-1}$. Replacing b by b^{-1} gives the presentation stated in the proposition and $\kappa = bza z^{-1} = z(az^{-1}b^{-1}z)^{-1}z^{-1}$ as before. In case (2) we replace z by z^{-1} and note that then $\kappa = az^{-1}b^{-1}z$. Finally in case (1) we replace b by b^{-1} and z by z^{-1} after which again we have $\kappa = az^{-1}b^{-1}z$. \diamond

Proposition 19.35. *Suppose that conditions 19.0.1 hold and $\check{\Phi}_3^+$ is not a union of tight components. If $m \equiv 2 \pmod{4}$ and $\Delta(\alpha, \beta)$ is even, then $\Delta(\alpha, \beta) = 2$.*

Proof. Suppose otherwise. Consider the 2-fold cover of $\widetilde{M} \rightarrow M$ which restricts to the cover $F \times I \rightarrow X^-$ on the $-$ -side of F and the trivial double cover on the $+$ -side of F . Since $m \equiv 2$

(mod 4) the boundary of \widetilde{M} is connected. Now β lifts to a slope β' on $\partial\widetilde{M}$ with associated filling a Seifert manifold with base orbifold $S^2(2, n, 2, n) \neq S^2(2, 2, 2, 2)$. Hence β' is a singular slope of some closed essential surface $S \subseteq \widetilde{M}$. Since the distance of α to β is even, α also lifts to a slope α' on $\partial\widetilde{M}$ with the associated filling Seifert with base orbifold a 2-sphere with three or four cone points. It's easy to see that the distance between α' and β' is $\Delta(\alpha, \beta)/2$. Hence as β' is a singular slope for S , S is incompressible in $\widetilde{M}(\alpha')$. As \widetilde{M} is hyperbolic, S cannot be a torus and therefore must be horizontal in $\widetilde{M}(\alpha')$. It cannot be separating as the base orbifold of $\widetilde{M}(\alpha')$ is orientable. Thus it is non-separating. But then [BGZ1, Theorem 1.5] implies the distance between α' and β' is at most 1, so $\Delta(\alpha, \beta) = 2$. \diamond

19.3.1. *M*(α) is very small. We assume that *M*(α) is very small in this subsection and prove $\Delta(\alpha, \beta) \leq 3$.

Lemma 19.36. *M*(β) contains no horizontal essential surfaces. Thus every closed orientable incompressible surface in *M*(β) is a vertical torus.

Proof. Suppose *M*(β) contains a horizontal essential surface *G*. Then for each ϵ , the components of $G \cap \widehat{X}^\epsilon$ are horizontal incompressible surfaces in \widehat{X}^ϵ . Hence if λ denotes the slope on \widehat{F} of the curves $G \cap \widehat{F}$, then λ is the fibre slope of the Seifert structure on \widehat{X}^- with base orbifold $D^2(2, 2)$. In particular, $\Delta(\lambda, \phi_+) = \Delta(\lambda, \alpha_-) = 1$. Then $\widehat{X}^+(\lambda)$ is a Seifert manifold with base orbifold $S^2(2, n)$ which admits a horizontal surface. Thus it must be $S^1 \times S^2$. But then $n = 2$ and therefore X^+ is a twisted *I*-bundle (Proposition 7.5), contrary to our assumptions. \diamond

Note that closed, essential surfaces in *M* have genus 2 or larger. Hence we deduce the following corollary.

Corollary 19.37. *If M contains a closed orientable essential surface, then the surface must compress in M*(β). \diamond

Lemma 19.38. *If β is not a singular slope, then any orientable essential surface H in M with boundary slope β has at least 4 boundary components.*

Proof. We may assume that $|\partial H|$ is minimal among all such surfaces. Then by [CGLS, Theorem 2.0.3], either β is a singular slope or \widehat{H} is incompressible in *M*(β). So by our assumption \widehat{H} is incompressible in *M*(β). Thus by Lemma 19.36, \widehat{H} is an incompressible torus in *M*(β). Hence $|\partial H| \geq m$ and so is at least 4. \diamond

We complete this part of the proof of Theorem 2.7 using $PSL_2(\mathbb{C})$ -character variety methods. We refer the reader to §6 of [BCSZ2] for the explanations of the relevant notation, background results, and references.

Now let $X_0 \subseteq X_{PSL_2}(M(\beta)) \subseteq X_{PSL_2}(M)$ be an irreducible curve which contains a character of a non-virtually-reducible representation. Let x be any ideal point of \widetilde{X}_0 . If \tilde{f}_α has finite value at x , then [BZ1, Proposition 4.10] and Corollary 19.37 imply that β is a singular slope, in which case we would have $\Delta(\alpha, \beta) \leq 1$. So every ideal point of \widetilde{X}_0 is a pole of \tilde{f}_α . In particular

X_0 provides a non-zero Culler-Shalen seminorm $\|\cdot\|_{X_0}$ on $H_1(\partial M; \mathbb{R})$ with β the unique slope with $\|\beta\|_{X_0} = 0$.

By [BCSZ2, Proposition 10.2] and [BZ1] we have

$$\|\alpha\|_{X_0} \leq s_{X_0} + 5$$

Let H be an essential surface associated to an ideal point x of \tilde{X}_0 . As x is a pole of \tilde{f}_α , H has boundary slope β . By Lemma 19.38, $|\partial H| \geq 4$. This implies, by the arguments in [BCSZ2, Proposition 6.6], that $s_{X_0} \geq 2$. Thus

$$\Delta(\alpha, \beta) = \frac{\|\alpha\|_{X_0}}{s_{X_0}} \leq 1 + 5/2 = 3.5$$

Thus $\Delta(\alpha, \beta) \leq 3$, which completes the proof when $M(\alpha)$ is very small.

19.3.2. *$M(\alpha)$ is not very small.* We suppose that $M(\alpha)$ is not very small in this subsection and that Y is a torus.

Lemma 19.39. *Suppose that conditions 19.0.1 hold and $\dot{\Phi}_3^+$ is not a union of tight components. If $\Delta(\alpha, \beta) > 5$ then and there is a vertex v of $\bar{\Gamma}_S$ such that $\mu(v) > m\Delta(\alpha, \beta) - 4$, then $m = 4$.*

Proof. Proposition 12.2 and Inequality 19.3.1 show that $3 \leq \text{valency}_{\bar{\Gamma}_S}(v) \leq 5$ and if v has valency 3, then $\Delta(\alpha, \beta) \leq 6$ with equality only if $m = 4$. If it has valency 4, Proposition 12.2 shows that $\varphi_3(v) \geq 1$. Lemma 19.7 then implies that $\Delta(\alpha, \beta)m$, the sum of the weights of the edges incident to v , is bounded above by $\max\{3m + 14, 4m + 4\}$. Hence if $\Delta(\alpha, \beta) > 5$, then $m = 4$ and $\Delta(\alpha, \beta) = 6$. Finally suppose that v has valency 5. In this case $\varphi_3(v) \geq 4$ (cf. Corollary 12.2) so Lemma 19.7 implies that $\Delta(\alpha, \beta)m \leq \max\{3m + 16, 4m + 6, 5m\}$. Hence if $\Delta(\alpha, \beta) > 5$, then $m = 4$. \diamond

In the absence of vertices v of $\bar{\Gamma}_S$ for which $\mu(v) > m\Delta(\alpha, \beta) - 4$, Corollary 12.4 implies that $\mu(v) = m\Delta(\alpha, \beta) - 4$ for all vertices.

Lemma 19.40. *Suppose that conditions 19.0.1 hold and $\dot{\Phi}_3^+$ is not a union of tight components. Assume moreover that $\mu(v) = m\Delta(\alpha, \beta) - 4$ for all vertices v of $\bar{\Gamma}_S$. If $\Delta(\alpha, \beta) > 5$, then either*

(i) $m = 4$, or

(ii) $m = 8$, $\Delta(\alpha, \beta) = 6$, each edge has weight 12, and $\bar{\Gamma}_S$ is rectangular.

Proof. Proposition 12.2 shows that $4 \leq \text{valency}_{\bar{\Gamma}_S}(v) \leq 6$ for all vertices of $\bar{\Gamma}_S$. Further, Proposition 12.5 shows that if

$$(19.40.1) \quad \begin{cases} \text{valency}_{\bar{\Gamma}_S}(v) = 4, \text{ then } \varphi_3(v) = 0, \varphi_4(v) = 4, \text{ and } \varphi_j(v) = 0 \text{ for } j > 4 \\ \text{valency}_{\bar{\Gamma}_S}(v) = 5, \text{ then } \varphi_3(v) = 3 \text{ and } \varphi_4(v) = 2, \text{ and } \varphi_j(v) = 0 \text{ for } j > 4 \\ \text{valency}_{\bar{\Gamma}_S}(v) = 6, \text{ then } \varphi_3(v) = 6, \text{ and } \varphi_j(v) = 0 \text{ for } j > 3 \end{cases}$$

Let v be a vertex of valency 6. Since the weight of each edge of $\bar{\Gamma}_S$ is at most $m + 4$, Lemma 19.7 implies that if some edge incident to v has weight larger than m then $\Delta(\alpha, \beta)m$, the sum of the weights of the edges incident to the vertex, is bounded above by $\max\{3m + 18, 4m + 8\}$. Hence Proposition 19.35 implies that $m = 4$ and $\Delta(\alpha, \beta) = 6$. If, on the other hand, each edge incident to v has weight m or less, then Inequality 19.3.1 shows that $\Delta(\alpha, \beta) = 6$ and each such edge has weight m . If some edge incident to v connects it to a vertex v_1 of valency less than 6, 19.40.1 implies that the valency of v_1 is 5 and $\varphi_3(v_1) = 3$. Then Lemma 19.7 shows that $6m$, the sum of the weights of the edges incident to v_1 , is bounded above by $\max\{4m + 10, 5m + 4\}$. In either case, $m = 4$. Assume then that each edge incident to v connects it to a vertex v_1 of valency 6. Proceeding inductively we see that if $m > 4$, then each vertex in the component of $\bar{\Gamma}_S$ containing v has valency 6. It follows that $\bar{\Gamma}_S$ is hexagonal (cf. the proof of Lemma 19.9) and each edge of $\bar{\Gamma}_S$ has weight m . Since $\bar{\Gamma}_S$ must have a positive edge, Lemma 19.5 shows that $m = 6$. But this is impossible by Proposition 19.35. Thus $m = 4$.

Next let v be a vertex of valency 5. Then $\Delta(\alpha, \beta)m$, the sum of the weights of the edges incident to v , is bounded above by $\max\{3m + 16, 4m + 10, 5m\}$. Since $\Delta(\alpha, \beta) > 5$, the only possibility is for $m = 4$.

Finally if there are no vertices of valency 5 or 6, each vertex of $\bar{\Gamma}_S$ has valency 4 and thus Identities 19.40.1 implies that it has no triangle faces. Lemma 11.5 then shows that $\bar{\Gamma}_S$ is rectangular. Inequality 19.3.1 shows that $m \leq 8$ and

$$\Delta(\alpha, \beta) \leq \begin{cases} 8 & \text{if } m = 4 \\ 6 & \text{if } m = 6, 8 \end{cases}$$

Since $\Delta(\alpha, \beta) \geq 6$, Proposition 19.35 implies that $m \neq 6$. If $m = 8$, it is easy to see that each edge of $\bar{\Gamma}_S$ has weight 12. This completes the proof. \diamond

By the last two results, the proof of Theorem 2.7 when $\dot{\Phi}_3^+$ is not a union of tight components reduces to proving the following two propositions.

Proposition 19.41. *If $m = 8, \Delta(\alpha, \beta) = 6, \bar{\Gamma}_S$ is rectangular, each of its edges has weight 12, and $\dot{\Phi}_3^+$ is not a union of tight components, then $\Delta(\alpha, \beta) \leq 5$.*

Proposition 19.42. *If $m = 4$ and $\dot{\Phi}_3^+$ is not a union of tight components, then $\Delta(\alpha, \beta) \leq 5$.*

Proof of Proposition 19.41. Each component of $\dot{\Phi}_j^-$ is tight for $j \geq 5$ (Proposition 9.4) and so $\dot{\Phi}_{11}^-$ has at least six tight components (Proposition 6.3(2)). On the other hand, since the weight of each edge of $\bar{\Gamma}_S$ is 12, at least two components of $\dot{\Phi}_{11}^-$ have two or more outer boundary components. It follows that $\dot{\Phi}_{11}^-$ has two components, each having two outer boundary components. We shall call the union of these two large components Q . By Lemma 19.5 each edge of $\bar{\Gamma}_S$ is negative. Without loss of generality we may assume that there is a parallel family of edges \bar{e} of Γ_S whose label sequence at one of the vertices v adjacent to \bar{e} is 1 2 3 4 5 6 7 8 1 2 3 4. Therefore b_1 and b_4 belong to Q , and by looking at the corners of the 4-gons of Γ_S contiguous to \bar{e} at v we see that b_5 and b_8 also belong to Q . As in Lemma 19.27, b_1 and b_4 belong to different components of Q , as do b_5 and b_8 .

This case is now ruled out exactly as in §19.2.2, with the corners (45) and (81) replacing (23) and (41). \diamond

The proof of Proposition 19.42 requires a certain amount of preparatory work. We use Δ to denote $\Delta(\alpha, \beta)$ and assume it is at least 6.

Let $\gamma_\beta \in \pi_1(M(\beta))$ be the element represented by the core K_β of the Dehn filling solid torus. Then $[\alpha] \in \pi_1(M)$ is sent to $\gamma_\beta^\Delta \in \pi_1(M(\beta)) = \pi_1(M)/\langle\langle[\beta]\rangle\rangle$. Hence $\pi_1(M)/\langle\langle[\alpha], [\beta]\rangle\rangle \cong \pi_1(M(\beta))/\langle\langle\gamma_\beta^\Delta\rangle\rangle$. Note that this group is a quotient of $\pi_1(M)/\langle\langle[\alpha]\rangle\rangle \cong \pi_1(M(\alpha))$.

The quotient of $\pi_1(M(\beta))$ by the fibre-class is $\Gamma = \pi_1(P^2(2, n))$. As before, denote the image of γ_β in Γ by κ . By Proposition 19.34 Γ admits a presentation $\langle a, b, z : a^2, b^n, abz^{-2} \rangle$ such that up to conjugation and taking inverse, $\kappa = az^{-1}b^{-1}z$. Thus if we set $G = \Gamma/\langle\langle\kappa^\Delta\rangle\rangle$, then G has a presentation

$$G = \langle a, b, z : a^2, b^n, abz^{-2}, (az^{-1}b^{-1}z)^\Delta \rangle$$

Since $\pi_1(M(\beta))/\langle\langle\gamma_\beta^\Delta\rangle\rangle$ is a quotient of $\pi_1(M(\alpha))$, the same is true for G . We will show that this is impossible when $\Delta \geq 6$.

First we give an alternate presentation of G which will be useful in the sequel.

Lemma 19.43. $G \cong \langle a, d, z : a^2, d^n, (ad)^\Delta, az^3dz^{-1} \rangle$.

Proof. Let $d = z^{-1}b^{-1}z$ and eliminate $b = zd^{-1}z^{-1}$. This gives the stated presentation. \diamond

Lemma 19.43 shows that G is obtained from the triangle group $T = T(2, n, \Delta)$ by adding a new generator z and the relation $az^3dz^{-1} = 1$. Such *relative presentations* ([BP]) have been studied extensively. In particular, since T is residually finite, a result of Gerstenhaber and Rothaus [GR] implies

Lemma 19.44. *The natural map $T \rightarrow G$ is injective.* \diamond

The specific relation az^3dz^{-1} is analysed by Edjvet and Howie in [EH], in the more general setting where T is replaced by an arbitrary group H generated by a and d . They show, using the method of Dehn (or Van Kampen) diagrams, that the natural map $H \rightarrow G$ is injective [EH, Proposition 1]. Combining this proof with a result of Bogley and Pride [BP] gives us the following.

Lemma 19.45. *Any finite subgroup of G is contained in a conjugate of T .*

Proof. Proposition 1 in [EH] is proved by showing that the relative presentation in question admits no non-empty spherical diagram, except for some special cases where the group H generated by a and d is small. We observe that these do not arise in our situation where $H = T(2, n, \Delta)$. The part of the proof of Proposition 1 that is relevant there is Case 2 ([EH, page 353]). In the exceptional cases that arise either H is finite, or there is a relation in H , other than a^2 , which contains at most three occurrences each of a and d , or a relation which is a product of between one and five words of the form $(ad^{\pm 1})^{\pm 1}$. Since none of these hold in our

case ($H = T(2, n, \Delta)$ where $n \geq 3$ and $\Delta \geq 6$), we conclude that our relative presentation of G admits no non-empty spherical diagram. In the dual language of *pictures*, this says that it admits no reduced spherical picture [BP]. Since the element $az^3dz^{-1} \in T * \langle z \rangle$ is not a proper power, Lemma 19.45 follows from [BP, (0.4)]. \diamond

Lemma 19.46. *The centre of G is finite.*

Proof. The orbifold Euler characteristic

$$\chi(\Gamma) = \chi^{orb}(P^2(2, n)) = 1 - \left(\frac{1}{2} + \frac{n-1}{n}\right) = \frac{1}{n} - \frac{1}{2}$$

Hence, unless $n = 3$ and $\Delta = 6$, $\chi(\Gamma) + \frac{1}{\Delta} < 0$, and so by [BZ2, Theorem 1.2] G has a normal subgroup G_0 of finite index with deficiency $\text{def}(G_0) \geq 2$ as long as there is a representation $\rho : \Gamma \rightarrow PSL_2(\mathbb{C})$ which preserves the orders of the torsion elements of Γ and which sends ad to an element of order Δ . This is easy to do by hand in our case, but we can also appeal to [DT, Lemma 8.1] where the result is proven in a broader context. By [Hil2, Corollaries 2.3.1 and 2.4.1], the centre $Z(G_0)$, and hence $Z(G)$, is finite.

Suppose then that $n = 3$ and $\Delta = 6$. In this case $\chi(\Gamma) + \frac{1}{\Delta} = 0$ and by [BZ2, Theorem 1.2] and [DT, Lemma 8.1] G has a normal subgroup G_0 of finite index with deficiency $\text{def}(G_0) \geq 1$. If $\text{def}(G_0) > 1$ we argue as above. If $\text{def}(G_0) = 1$, [Hil1, Corollary 1, page 38] implies that if $Z(G_0)$ is infinite then the commutator subgroup $[G_0, G_0]$ is free. But $[G, G]$ contains $[T, T]$ (by Lemma 19.44), which is isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$, and hence $[G_0, G_0]$ contains a copy of $\mathbb{Z} \oplus \mathbb{Z}$. It follows that $Z(G_0)$, and therefore $Z(G)$, is finite in this case also. \diamond

Since the triangle group T has trivial centre, Lemmas 19.45 and 19.46 give

Proposition 19.47. *The centre of G is trivial.* \diamond

Proof of Proposition 19.42. Suppose that $\Delta(\alpha, \beta) > 5$ and let $\varphi : \pi_1(M(\alpha)) \rightarrow G$ be the epimorphism described above. Recall that $M(\alpha)$ is a small Seifert fibred manifold with hyperbolic base orbifold $S^2(a, b, c)$. Let Z be the (infinite cyclic) center of $\pi_1(M(\alpha))$. By Proposition 19.47 $\varphi(Z) = \{1\}$, and hence φ factors through $\pi_1(M(\alpha))/Z \cong T(a, b, c) = T'$. Since T' is generated by elements of finite order, its image under the induced homomorphism is contained in $\langle\langle T \rangle\rangle$ by Lemma 19.45. Since $G/\langle\langle T \rangle\rangle \cong \mathbb{Z}/2$, this is a contradiction. \diamond

REFERENCES

- [Ag] I. Agol, *Bounds on exceptional Dehn filling II*, *Geom. Topol.* **14** (2010), -1921–1940
- [BP] W. A. Bogley and S. J. Pride, *Aspherical relative presentations*, *Proc. Edinburgh Math. Soc.* **35** (1992), 1–39.
- [BCSZ1] S. Boyer, M. Culler, P. Shalen and X. Zhang, *Characteristic subsurfaces and Dehn filling*, *Trans. Amer. Math. Soc.*, 357 (2005) 2389–2444.
- [BCSZ2] —, *Characteristic subsurfaces, character varieties and Dehn fillings*, *Geometry & Topology* **12** (2008) 233–297.

- [BGZ1] S. Boyer, C. McA. Gordon, and X. Zhang, *Dehn fillings of large hyperbolic 3-manifolds*, J. Diff. Geom., **58** (2001) 263-308.
- [BGZ2] ———, *Reducible and finite Dehn fillings*, J. Lond. Math. Soc. **79** (2009), 72–84.
- [BGZ3] ———, *Dehn fillings of knot manifolds containing essential once-punctured tori*, to appear in Trans. Amer. Math. Soc.
- [BGZ4] ———, *Dehn fillings of knot manifolds containing essential twice-punctured tori*, in preparation.
- [BZ1] S. Boyer and X. Zhang, *On Culler-Shalen seminorms and Dehn fillings*, Ann. Math. **148** (1998), 737–801.
- [BZ2] ———, *Virtual Haken 3-manifolds and Dehn filling*, Topology **39** (2000), 103–114.
- [CL] D. Cooper and D. Long, *Virtually Haken Dehn-filling*, J. Diff. Geom. **52** (1999), 173–187.
- [CGLS] M. Culler, C. M. Gordon, J. Luecke and P. Shalen, *Dehn surgery on knots*, Ann. of Math. **125** (1987) 237–300.
- [DT] N. Dunfield and W. Thurston, *The virtual Haken conjecture: Experiments and Examples*, Geom. & Top. **7** (2003), 399–441.
- [EH] M. Edjvet and J. Howie, *The solution of length four equations over groups*, Trans. Amer. Math. Soc. **326**, (1991), 345–369.
- [Ga] D. Gabai, *Foliations and the topology of 3-manifolds II*, J. Diff. Geom. **26** (1987), 461–478.
- [GR] M. Gerstenhaber and O. S. Rothaus, *The solution of sets of equations in groups*, Proc. Nat. Acad. Sci. U.S.A. **48** (1962), 1531–1533.
- [Go] C. McA. Gordon, *Boundary slopes of punctured tori in 3-manifolds*, Trans. Amer. Math. Soc. **350** (1998) 371-415.
- [GL] C. McA. Gordon and J. Luecke, *Reducible manifolds and Dehn surgery*, Topology **35** (1996), 385–410.
- [GW] C. McA. Gordon and Y.-Q. Wu, *Toroidal Dehn fillings on hyperbolic 3-manifolds*, Memoirs Amer. Math. Soc. **194** (2008), no. 909.
- [He] J. Hempel, *3-Manifolds*, Annals of Math. Studies **86**, Princeton University Press and University of Tokyo Press, 1976.
- [Hil1] J. Hillman, *2-knots and their groups*, Austr. Math. Soc. Lecture Series **5**, Cambridge University Press, 1989.
- [Hil2] J. Hillman, *Four-manifolds, geometries and knots*, Geom. Top. Monographs **5**, 2002.
- [JS] Wm. Jaco and P. Shalen, *Seifert Fibered Spaces in 3-Manifolds*, Mem. Amer. Math. Soc. **220**, Providence, 1979.
- [Jo] K. Johansson, *Homotopy Equivalences of 3-Manifolds with Boundaries*, Lecture Notes in Mathematics **761**, Springer-Verlag, Berlin, 1979.
- [LM] M. Lackenby and R. Meyerhoff, *The maximal number of exceptional Dehn surgeries*, preprint (2008), arXiv:0808.1176.
- [Li] T. Li, *Immersed essential surfaces in hyperbolic 3-manifolds*, Comm. Anal. Geom. **10** (2002), 275–290.
- [Oh] S. Oh, *Reducible and toroidal manifolds obtained by Dehn filling*, Top. Appl. **75** (1997), 93–104.
- [Wu1] Y.-Q. Wu, *Incompressibility of surfaces in surgered 3-manifolds*, Topology **31** (1992), 271–279
- [Wu2] ———, *Dehn fillings producing reducible manifolds and toroidal manifolds*, Topology **37** (1998), 95–108.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, 201 AVENUE DU PRÉSIDENT-KENNEDY, MONTRÉAL, QC H2X 3Y7.

E-mail address: boyer.steven@uqam.ca

URL: <http://www.cirget.uqam.ca/boyer/boyer.html>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, 1 UNIVERSITY STATION, AUSTIN, TX 78712, USA.

E-mail address: gordon@math.utexas.edu

URL: <http://www.ma.utexas.edu/text/webpages/gordon.html>

DEPARTMENT OF MATHEMATICS, UNIVERSITY AT BUFFALO, BUFFALO, NY, 14214-3093, USA.

E-mail address: xinzhang@buffalo.edu

URL: <http://www.math.buffalo.edu/~xinzhang/>