

# POSET EDGE-LABELLINGS AND LEFT MODULARITY

PETER MCNAMARA AND HUGH THOMAS

ABSTRACT. It is known that a graded lattice of rank  $n$  is supersolvable if and only if it has an EL-labelling where the labels along any maximal chain are exactly the numbers  $1, 2, \dots, n$  without repetition. These labellings are called  $S_n$  EL-labellings, and having such a labelling is also equivalent to possessing a maximal chain of left modular elements. In the case of an ungraded lattice, there is a natural extension of  $S_n$  EL-labellings, called interpolating labellings. We show that admitting an interpolating labelling is again equivalent to possessing a maximal chain of left modular elements. Furthermore, we work in the setting of an arbitrary bounded poset as all the above results generalize to this case.

RÉSUMÉ. On sait qu'un treillis gradué de rang  $n$  est supersoluble si et seulement si il possède un EL-étiquetage où les étiquettes de n'importe quelle chaîne maximale sont les nombres  $1, 2, \dots, n$  sans répétition. Ces étiquetages sont appelés  $S_n$  EL-étiquetages. L'existence d'un tel étiquetage d'un treillis est aussi équivalente à l'existence d'une chaîne maximale d'éléments modulaires à gauche. Dans le cas d'un treillis non-gradué, il existe une extension naturelle des  $S_n$  EL-étiquetages, appelée étiquetages interpolants. Nous démontrons qu'avoir un étiquetage interpolant est, encore une fois, équivalent à avoir une chaîne maximale d'éléments modulaires à gauche. De plus, tous ces résultats se généralisent aux ensembles ordonnés bornés même s'ils ne sont pas des treillis.

## 1. INTRODUCTION

An *edge-labelling* of a poset  $P$  is a map from the edges of the Hasse diagram of  $P$  to  $\mathbb{Z}$ . Our primary goal is to express certain classical properties of  $P$  in terms of edge-labellings admitted by  $P$ . The idea of studying edge-labellings of posets goes back to [9]. An important milestone was [2], where A. Björner defined EL-labellings, and showed that if a poset admits an EL-labelling, then it is shellable and hence Cohen-Macaulay. We will be interested in a subclass of EL-labellings, known as  $S_n$  EL-labellings. In [10], R. Stanley introduced supersolvable lattices and showed that they admit  $S_n$  EL-labellings. Examples of supersolvable lattices include distributive lattices, the lattice of partitions of  $[n]$ , the lattice of non-crossing partitions of  $[n]$  and the lattice of subgroups of a supersolvable group (hence the terminology). It was shown in [8] that a finite graded lattice of rank  $n$  is supersolvable if and only if it admits an  $S_n$  EL-labelling. In many ways, this characterization of lattice supersolvability in terms of edge-labellings serves as the starting point for our investigations.

For basic definitions concerning partially ordered sets, see [11]. We will say that a poset  $P$  is *bounded* if it contains a unique minimal element and a unique maximal element, denoted  $\hat{0}$  and  $\hat{1}$  respectively. All the posets we will consider will be finite and bounded. A chain of a poset  $P$  is said to be *maximal* if it is maximal under inclusion. We say that  $P$  is *graded* if all the maximal chains of  $P$  have the same length, and we call this length the *rank* of  $P$ . We will write  $x < y$  if  $y$  covers  $x$  in  $P$  and  $x \leq y$  if  $y$  either covers or equals  $x$ . The edge-labelling  $\gamma$  of  $P$  is said to be an *EL-labelling* if for any  $y < z$  in  $P$ ,




FIGURE 1

- (i) there is a unique unrefinable chain  $y = w_0 < w_1 < \cdots < w_r = z$  such that  $\gamma(w_0, w_1) \leq \gamma(w_1, w_2) \leq \cdots \leq \gamma(w_{r-1}, w_r)$ , and
- (ii) the sequence of labels of this chain (referred to as the *increasing chain* from  $y$  to  $z$ ), when read from bottom to top, lexicographically precedes the labels of any other unrefinable chain from  $y$  to  $z$ .

This concept originates in [2]; for the case where  $P$  is not graded, see [3, 4]. If  $P$  is graded of rank  $n$  with an EL-labelling  $\gamma$ , then  $\gamma$  is said to be an  $S_n$  *EL-labelling* if the labels along any maximal chain of  $P$  are all distinct and are elements of  $[n]$ . In other words, for every maximal chain  $\hat{0} = w_0 < w_1 < \cdots < w_n = \hat{1}$  of  $P$ , the map sending  $i$  to  $\gamma(w_{i-1}, w_i)$  is a permutation of  $[n]$ . Note that the second condition in the definition of an EL-labelling is redundant in this case.


**Example 1.1.** Any finite distributive lattice has an  $S_n$  EL-labelling. Let  $L$  be a finite distributive lattice of rank  $n$ . By the Fundamental Theorem of Finite Distributive Lattices [1, p. 59, Thm. 3], that is equivalent to saying that  $L = J(Q)$ , the lattice of order ideals of some  $n$ -element poset  $Q$ . Let  $\omega : Q \rightarrow [n]$  be a linear extension of  $Q$ , i.e., any bijection labeling the vertices of  $Q$  that is order-preserving (if  $a < b$  in  $Q$  then  $\omega(a) < \omega(b)$ ). This labeling of the vertices of  $Q$  defines a labeling of the edges of  $J(Q)$  as follows. If  $y$  covers  $x$  in  $J(Q)$ , then the order ideal corresponding to  $y$  is obtained from the order ideal corresponding to  $x$  by adding a single element, labeled by  $i$ , say. Then we set  $\gamma(x, y) = i$ . This gives us an  $S_n$  EL-labelling for  $L = J(Q)$ . Figure 1 shows a labelled poset and its lattice of order ideals with the appropriate edge-labelling.

A finite lattice  $L$  is said to be *supersolvable* if it contains a maximal chain, called an *M-chain* of  $L$ , which together with any other chain in  $L$  generates a distributive sublattice. We can label each such distributive sublattice by the method described in Example 1.1 in such a way that the M-chain is the unique increasing maximal chain. As shown in [10], this will assign a unique label to each edge of  $L$  and the resulting global labeling of  $L$  is an  $S_n$  EL-labelling.

There is also a characterization of lattice supersolvability in terms left modularity. Given an element  $x$  of a finite lattice  $L$ , and a pair of elements  $y \leq z$ , it is always true that

$$(x \vee y) \wedge z \geq (x \wedge z) \vee y. \quad (1)$$

The element  $x$  is said to be *left modular* if, for all  $y \leq z$ , equality holds in (1). As shown in [10], any M-chain of  $L$  is always a left modular maximal chain, that is, a maximal chain each of whose elements is left modular. Furthermore, it is shown by L. S.-C. Liu [6] that if  $L$  is a finite graded lattice with a left modular maximal chain  $M$ , then  $L$  has an  $S_n$


 FIGURE 2. The Tamari lattice  $T_4$  and its interpolating EL-labelling

EL-labelling with increasing maximal chain  $M$ . Since this implies that  $L$  is supersolvable, we conclude the following.

**Theorem 1.** *Let  $L$  be a finite graded lattice of rank  $n$ . Then the following are equivalent:*

- (1)  $L$  has an  $S_n$  EL-labelling,
- (2)  $L$  has a left modular maximal chain,
- (3)  $L$  is supersolvable.

It is shown in [10] that if  $L$  is upper-semimodular, then possessing a left modular maximal chain and being supersolvable are equivalent. Theorem 1 is a considerable strengthening of this. Here we used  $S_n$  EL-labellings to connect left modularity and supersolvability. It is natural to ask for a direct proof of the equivalence of (2) and (3).

Our goal is to generalize Theorem 1 to the case when  $L$  is not graded and, moreover, to the case when  $L$  is not necessarily a lattice. We now wish to define natural generalizations of  $S_n$  EL-labellings and of maximal left-modular chains.

**Definition 1.2.** An EL-labelling  $\gamma$  of  $P$  is said to be *interpolating* if, for any  $y < u < z$ , either

- (i)  $\gamma(y, u) < \gamma(u, z)$  or
- (ii) the increasing chain from  $y$  to  $z$ , say  $y = w_0 < w_1 < \dots < w_r = z$ , has the properties that its labels are strictly increasing and that  $\gamma(w_0, w_1) = \gamma(u, z)$  and  $\gamma(w_{r-1}, w_r) = \gamma(y, u)$ .

**Example 1.3.** The reader is invited to check that the labelling of the non-graded poset shown in Figure 2 is an interpolating EL-labelling. In fact, the poset shown is the so-called “Tamari lattice”  $T_4$ . For all positive integers  $n$ , there exists a Tamari lattice  $T_n$  with  $C_n$  elements, where  $C_n = \frac{1}{n+1} \binom{2n}{n}$ , the  $n$ th Catalan number. More information on the Tamari lattice can be found in [4, §9], [5, §7] and the references given there, and in [6, §3.2], where this interpolating EL-labelling appears. The Tamari lattice is shown to have an EL-labelling in [4] and is shown to have a left modular maximal chain in [5].

If  $P$  is graded of rank  $n$  and has an interpolating labelling  $\gamma$  in which the labels on the increasing maximal chain reading from bottom to top are  $1, 2, \dots, n$ , then we can check (cf. Lemma 3.2) that  $\gamma$  is an  $S_n$  EL-labelling.

Our next step is to define left modularity in the non-lattice case. Let  $x$  and  $y$  be elements of  $P$ . We know that  $x$  and  $y$  have at least one common upper bound, namely




FIGURE 3

$\hat{1}$ . If the set of common upper bounds of  $x$  and  $y$  has a least element, then we denote it by  $x \vee y$ . Similarly, if  $x$  and  $y$  have a greatest common lower bound, then we denote it by  $x \wedge y$ .

Now let  $w$  and  $z$  be elements of  $P$  with  $w, z \geq y$ . Consider the set of common lower bounds for  $w$  and  $z$  that are also greater than or equal to  $y$ . Clearly,  $y$  is in this set. If this set has a greatest element, then we denote it by  $w \wedge_y z$  and we say that  $w \wedge_y z$  is well-defined (in  $[y, \hat{1}]$ ). We see that  $(x \vee y) \wedge_y z$  is well-defined in the poset shown in Figure 3, even though  $(x \vee y) \wedge z$  is not. Similarly, let  $w$  and  $y$  be elements of  $P$  with  $w, y \leq z$ . If the set  $\{u \in P \mid u \geq w, y \text{ and } u \leq z\}$  has a least element, then we denote it by  $w \vee^z y$  and we say that  $w \vee^z y$  is well-defined (in  $[\hat{0}, z]$ ). We will usually be interested in expressions of the form  $(x \vee y) \wedge_y z$  and  $(x \wedge z) \vee^z y$ . The reader that is solely interested in the lattice case can choose to ignore the subscripts and superscripts on the meet and join symbols.

**Definition 1.4.** An element  $x$  of  $P$  is said to be *viable* if, for all  $y \leq z$  in  $P$ ,  $(x \vee y) \wedge_y z$  and  $(x \wedge z) \vee^z y$  are well-defined. A maximal chain of  $P$  is said to be viable if each of its elements is viable.

**Example 1.5.** The poset shown in Figure 3 is certainly not a lattice but the reader can check that the increasing maximal chain is viable.

**Definition 1.6.** A viable element  $x$  of  $P$  is said to be *left modular* if, for all  $y \leq z$  in  $P$ ,

$$(x \vee y) \wedge_y z = (x \wedge z) \vee^z y.$$

A maximal chain of  $P$  is said to be left modular if each of its elements is viable and left modular.

This brings us to the first of our main theorems.

**Theorem 2.** Let  $P$  be a bounded poset with a viable left modular maximal chain  $M$ . Then  $P$  has an interpolating EL-labelling with  $M$  as its increasing maximal chain.

The proof of this theorem will be the content of the next section. In section 3, we will prove the following converse result.

**Theorem 3.** Let  $P$  be a bounded poset with an interpolating EL-labelling. The unique increasing chain from  $\hat{0}$  to  $\hat{1}$  is a viable left modular maximal chain.

As one consequence, in the lattice case, we have that  $P$  has an interpolating EL-labelling with increasing maximal chain  $M$  if and only if  $M$  is a left modular maximal

chain of  $P$ . As another consequence, in the graded poset case, we have given an answer to the question of when  $P$  has an  $S_n$  EL-labelling. This has ramifications on the existence of a “good 0-Hecke algebra action” on the maximal chains of the poset, as discussed in [8].

These two theorems, when compared with Theorem 1, might lead one to ask about possible supersolvability results for bounded posets that aren’t graded lattices. This problem is discussed in section 4. We obtain a satisfactory result in the graded case but the ungraded case is left as an open problem.

## 2. PROOF OF THEOREM 2

We suppose that  $P$  is a bounded poset with a viable left modular maximal chain  $M : \hat{0} = x_0 \leq x_1 \leq \cdots \leq x_n = \hat{1}$ . We want to show that  $P$  has an interpolating EL-labelling. We begin with some lemmas which build on the viability and left modularity properties.

**Lemma 2.1.** *Suppose that  $y \leq w \leq z$  in  $P$  and let  $x \in M$ . Then  $((x \wedge z) \vee^z y) \vee^z w$  is well-defined and equals  $(x \wedge z) \vee^z w$ . Similarly,  $((x \vee y) \wedge_y z) \wedge_y w$  is well-defined and equals  $(x \vee y) \wedge_y w$ .*

*Proof.* It is routine to check that, in  $[\hat{0}, z]$ ,  $(x \wedge z) \vee^z w$  is the least common upper bound for  $w$  and  $(x \wedge z) \vee^z y$ , and that, in  $[y, \hat{1}]$ ,  $(x \vee y) \wedge_y w$  is the greatest common lower bound lower bound for  $(x \vee y) \wedge_y z$  and  $w$ .  $\square$

**Lemma 2.2.** *Suppose that  $t \leq u$  in  $[y, z]$  and  $x \in M$ . Let  $w = (x \vee y) \wedge_y z = (x \wedge z) \vee^z y$  in  $[y, z]$ . Then  $(w \vee^z t) \wedge_t u$  and  $(w \wedge_y u) \vee^u t$  are well-defined elements of  $[t, u]$  and are equal.*

*Proof.* We see that, by Lemma 2.1,

$$\begin{aligned} (x \vee t) \wedge_t u &= ((x \vee t) \wedge_t z) \wedge_t u = ((x \wedge z) \vee^z t) \wedge_t u \\ &= (((x \wedge z) \vee^z y) \vee^z t) \wedge_t u = (w \vee^z t) \wedge_t u. \end{aligned}$$

Similarly,

$$(x \wedge u) \vee^u t = (w \wedge_y u) \vee^u t$$

But  $(x \vee t) \wedge_t u = (x \wedge u) \vee^u t$ , yielding the result.  $\square$

**Lemma 2.3.** *Suppose  $x$  and  $w$  are viable and that  $x$  is left modular in  $P$ .*

- (a) *If  $x \leq w$  then for any  $z$  in  $P$  we have  $x \wedge z \leq w \wedge z$ .*
- (b) *If  $w \leq x$  then for any  $y$  in  $P$  we have  $w \vee y \leq x \vee y$ .*

Part (b) appears in the lattice case in [6, Lemma 2.5.6] and [7, Lemma 5.3].

*Proof.* We prove (a); (b) is similar. Assume, seeking a contradiction, that  $x \wedge z < u < w \wedge z$  for some  $u \in P$ . Now  $u \leq z$  and  $u \leq w$ . It follows that  $u \not\leq x$ .

Now  $x < x \vee u \leq w$ . Therefore,  $w = x \vee u$ . So

$$u = (x \wedge z) \vee^z u = (x \vee u) \wedge_u z = w \wedge z,$$

which is a contradiction.  $\square$

We now prove a slight extension of [6, Lemma 2.5.7] and [7, Lemma 5.4].

**Lemma 2.4.** *The elements of  $[y, z]$  of the form  $(x_i \vee y) \wedge_y z$  form a viable left modular maximal chain in  $[y, z]$ .*

*Proof.* Lemma 2.2 gives the viability and left modularity properties. By Lemma 2.3(b),  $x_i \vee y \leq x_{i+1} \vee y$ . By Lemma 2.2 with  $z = \hat{1}$ , we have that  $x_i \vee y$  is left modular in  $[y, \hat{1}]$ . Therefore,  $(x_i \vee y) \wedge_y z \leq (x_{i+1} \vee y) \wedge_y z$  by Lemma 2.3(a).  $\square$

We are now ready to specify an edge-labelling for  $P$ . Let  $P$  be a bounded poset with a viable left modular maximal chain  $M : \hat{0} = x_0 \leq x_1 \leq \dots \leq x_n = \hat{1}$ . We choose a label set  $l_1 < \dots < l_n$  of natural numbers. (For most purposes, we can let  $l_i = i$ .) We define an edge-labelling  $\gamma$  on  $P$  by, for  $y \leq z$ ,  $\gamma(y, z) = l_i$  if

$$(x_{i-1} \vee y) \wedge_y z = y \quad \text{and} \quad (x_i \vee y) \wedge_y z = z.$$

It is easy to see that  $\gamma$  is well-defined. We will refer to it as the labelling induced by  $M$  and the label set  $\{l_i\}$ . When  $P$  is a lattice, this labelling appears, for example, in [6] and [12]. As in [6], we can give an equivalent definition of  $\gamma$  as follows.

**Lemma 2.5.** *Suppose  $y \leq z$  in  $P$ . Then  $\gamma(y, z) = l_i$  if and only if*

$$i = \min\{j \mid x_j \vee y \geq z\} = \max\{j + 1 \mid x_j \wedge z \leq y\}.$$

*Proof.* That  $i = \min\{j \mid x_j \vee y \geq z\}$  is immediate from the definition of  $\gamma$ . By left modularity,  $\gamma(y, z) = l_i$  if and only if  $(x_{i-1} \wedge z) \vee^z y = y$  and  $(x_i \wedge z) \vee^z y = z$ . In other words,  $x_{i-1} \wedge z \leq y$  and  $x_i \wedge z \not\leq y$ . It follows that  $i = \max\{j + 1 \mid x_j \wedge z \leq y\}$ .  $\square$

We are now ready for the last, and most important, of our preliminary results. Let  $[y, z]$  be an interval in  $P$ . We call the maximal chain of  $[y, z]$  from Lemma 2.4 the induced left modular maximal chain of  $[y, z]$ . One way to get a second edge-labelling for  $[y, z]$  would be to take the labelling induced in  $[y, z]$  by this induced maximal chain. We now prove that, for a suitable choice of label set, this labelling coincides with  $\gamma$ .

**Proposition 2.6.** *Let  $P$  be a bounded poset,  $\hat{0} = x_0 \leq x_1 \leq \dots \leq x_n = \hat{1}$  a viable left modular maximal chain and  $\gamma$  the corresponding edge-labelling with label set  $\{l_i\}$ . Let  $y < z$ , and define  $c_i$  by saying*

$$\begin{aligned} y &= (x_0 \vee y) \wedge_y z = \dots = (x_{c_1-1} \vee y) \wedge_y z \\ &< (x_{c_1} \vee y) \wedge_y z = \dots = (x_{c_2-1} \vee y) \wedge_y z < \dots \\ &< (x_{c_r} \vee y) \wedge_y z = \dots = (x_n \vee y) \wedge_y z. \end{aligned}$$

*Let  $m_i = l_{c_i}$ . Let  $\delta$  be the labelling of  $[y, z]$  induced by its induced left modular maximal chain and the label set  $\{m_i\}$ . Then  $\delta$  agrees with  $\gamma$  restricted to the edges of  $[y, z]$ .*

*Proof.* Suppose  $t \leq u$  in  $[y, z]$ . Using ideas from the proof of Lemma 2.2,

$$\begin{aligned} \delta(t, u) = m_i &\Leftrightarrow (((x_{c_i-1} \vee y) \wedge_y z) \vee^z t) \wedge_t u = t \quad \text{and} \\ &\quad (((x_{c_i} \vee y) \wedge_y z) \vee^z t) \wedge_t u = u \\ &\Leftrightarrow (x_{c_i-1} \vee t) \wedge_t u = t \quad \text{and} \quad (x_{c_i} \vee t) \wedge_t u = u \\ &\Leftrightarrow \gamma(t, u) = l_{c_i}. \end{aligned}$$

$\square$

*Proof of Theorem 2.* We now know that the induced left modular chain in  $[y, z]$  has (strictly) increasing labels, say  $m_1 < m_2 < \dots < m_r$ . Our first step is to show that it is the only maximal chain with (weakly) increasing labels. Suppose that  $y = w_0 \leq w_1 \leq \dots \leq w_r = z$  is the induced chain and that  $y = u_0 \leq u_1 \leq \dots \leq u_s = z$  is another chain with increasing labels.

If  $s = 1$  then  $y \lessdot z$  and the result is clear. Suppose  $s \geq 2$ . By Proposition 2.6, we may assume that the labelling on  $[y, z]$  is induced by the induced left modular chain  $\{w_i\}$ . In particular, we have that  $\gamma(u_i, u_{i+1}) = m_l$  where  $l = \min\{j \mid w_j \vee^z u_i \geq u_{i+1}\}$ . Let  $k$  be the least number such that  $u_k \geq w_1$ . Then it is clear that  $\gamma(u_{k-1}, u_k) = m_1$ . Note that this is the smallest label that can occur on any edge in  $[y, z]$ . Since the labels on the chain  $\{u_i\}$  are assumed to be increasing, we must have  $\gamma(u_0, u_1) = m_1$ . It follows that  $w_1 \vee^z u_0 \geq u_1$  and since  $y \lessdot w_1$ , we must have  $u_1 = w_1$ . Thus, by induction, the two chains coincide. We conclude that the induced left modular maximal chain is the only chain in  $[y, z]$  with increasing labels.

It also has the lexicographically least set of labels. To see this, suppose that  $y = u_0 \lessdot u_1 \lessdot \cdots \lessdot u_s = z$  is another chain in  $[y, z]$ . We assume that  $u_1 \neq w_1$  since, otherwise, we can just restrict our attention to  $[u_1, z]$ . We have  $\gamma(u_0, u_1) = m_l$ , where  $l = \min\{j \mid w_j \geq u_1\} \geq 2$  since  $w_1 \not\geq u_1$ . Hence  $\gamma(u_0, u_1) \geq m_2 > \gamma(w_0, w_1)$ . This gives that  $\gamma$  is an EL-labelling. (That  $\gamma$  is an EL-labelling was already shown in the lattice case in [6] and [12].)

Finally, we show that it is an interpolating EL-labelling. If  $y \lessdot u \lessdot z$  is not the induced left modular maximal chain in  $[y, z]$ , then let  $y = w_0 \lessdot w_1 \lessdot \cdots \lessdot w_r = z$  be the induced left modular maximal chain. We have that  $\gamma(y, u) = m_l$  where

$$l = \min\{j \mid w_j \vee^z y \geq u\} = \min\{j \mid w_j \geq u\} = r$$

since  $u \lessdot z$ . Therefore,  $\gamma(y, u) = m_r$ . Also,  $\gamma(u, z) = m_l$  where

$$l = \max\{j + 1 \mid w_j \wedge_y z \leq u\} = \max\{j + 1 \mid w_j \leq u\} = 1$$

since  $y \lessdot u$ . Therefore,  $\gamma(y, u) = m_1$ , as required.  $\square$

### 3. PROOF OF THEOREM 3

We suppose that  $P$  is a bounded poset with an interpolating EL-labelling  $\gamma$ . Let  $\hat{0} = x_0 \lessdot x_1 \lessdot \cdots \lessdot x_n = \hat{1}$  be the increasing chain from  $\hat{0}$  to  $\hat{1}$  and let  $l_i = \gamma(x_{i-1}, x_i)$ . We wish to establish some basic facts about interpolating labellings.

Let  $y = w_0 \lessdot w_1 \lessdot \cdots \lessdot w_r = z$ . Suppose that, for some  $i$ , we have  $\gamma(w_{i-1}, w_i) > \gamma(w_i, w_{i+1})$ . Then the “basic replacement” at  $i$  takes the given chain and replaces the subchain  $w_{i-1} \lessdot w_i \lessdot w_{i+1}$  by the increasing chain from  $w_{i-1}$  to  $w_{i+1}$ . The basic tool for dealing with interpolating labellings is the following well-known fact about EL-labellings.

**Lemma 3.1.** *Let  $y = w_0 \lessdot w_1 \lessdot \cdots \lessdot w_r = z$ . Successively perform basic replacements on this chain, and stop when no more basic replacements can be made. This algorithm terminates, and yields the increasing chain from  $y$  to  $z$ .*

*Proof.* At each step, the sequence of labels on the new chain lexicographically precedes the sequence on the old chain, so the process must terminate, and it is clear that it terminates in an increasing chain.  $\square$

We now prove some simple consequences of this lemma.

**Lemma 3.2.** *Let  $m$  be the chain  $y = w_0 \lessdot w_1 \lessdot \cdots \lessdot w_r = z$ . Then the labels on  $m$  all occur on the increasing chain from  $y$  to  $z$  and are all different. Furthermore, all the labels on the increasing chain from  $y$  to  $z$  are bounded between the lowest and highest labels on  $m$ .*

*Proof.* That the labels on the given chain all occur on the increasing chain follows immediately from Lemma 3.1 and the fact that after a basic replacement, the labels on the

old chain all occur on the new chain. Similar reasoning implies that the labels on the increasing chain are bounded between the lowest and highest labels on  $m$ .

That the labels are all different again follows from Lemma 3.1. Suppose otherwise. By repeated basic replacements, one obtains a chain which has two successive equal labels, which is not permitted by the definition of an interpolating labelling.  $\square$

**Lemma 3.3.** *Let  $z \in P$  such that there is some chain from  $\hat{0}$  to  $z$  all of whose labels are in  $\{l_1, \dots, l_i\}$ . Then  $z \leq x_i$ . Conversely, if  $z \leq x_i$ , then all the labels on any chain from  $\hat{0}$  to  $z$  are in  $\{l_1, \dots, l_i\}$ .*

*Proof.* We begin by proving the first statement. By Lemma 3.2, the labels on the increasing chain from  $\hat{0}$  to  $z$  are in  $\{l_1, \dots, l_i\}$ . Find the increasing chain from  $z$  to  $\hat{1}$ . Let  $w$  be the element in that chain such that all the labels below it on the chain are in  $\{l_1, \dots, l_i\}$ , and those above it are in  $\{l_{i+1}, \dots, l_n\}$ . Again, by Lemma 3.2, the increasing chain from  $\hat{0}$  to  $w$  has all its labels in  $\{l_1, \dots, l_i\}$ , and the increasing chain from  $w$  to  $\hat{1}$  has all its labels in  $\{l_{i+1}, \dots, l_n\}$ . Thus  $w$  is on the increasing chain from  $\hat{0}$  to  $\hat{1}$ , and so  $w = x_i$ . But by construction  $w \geq z$ . So  $x_i \geq z$ .

To prove the converse, observe that by Lemma 3.2, no label can occur more than once on any chain. But since every label in  $\{l_{i+1}, \dots, l_n\}$  occurs on the increasing chain from  $x_i$  to  $\hat{1}$ , no label from among that set can occur on any edge below  $x_i$ .  $\square$

The obvious dual of Lemma 3.3 is proved similarly:

**Corollary 3.4.** *Let  $z \in P$  such that there is some chain from  $z$  to  $\hat{1}$  all of whose labels are in  $\{l_{i+1}, \dots, l_n\}$ . Then  $z \geq x_i$ . Conversely, if  $z \geq x_i$ , then all the labels on any chain from  $z$  to  $\hat{1}$  are in  $\{l_{i+1}, \dots, l_n\}$ .*

We are now ready to prove the necessary viability properties.

**Lemma 3.5.**  *$x_i \vee z$  and  $x_i \wedge z$  are well-defined for any  $z \in P$  and for  $i = 1, 2, \dots, n$ .*

*Proof.* We will prove that  $x_i \wedge z$  is well-defined. The proof that  $x_i \vee z$  is well-defined is similar. Let  $w$  be the maximum element on the increasing chain from  $\hat{0}$  to  $z$  such that all labels on the increasing chain between  $\hat{0}$  and  $w$  are in  $\{l_1, \dots, l_i\}$ . Clearly  $w \leq z$  and, by Lemma 3.3,  $w \leq x_i$ .

Suppose  $y \leq z, x_i$ . It follows that all labels from  $\hat{0}$  to  $y$  are in  $\{l_1, \dots, l_i\}$ . Consider the increasing chain from  $y$  to  $z$ . There exists an element  $u$  on this chain such that all the labels on the increasing chain from  $\hat{0}$  to  $u$  are in  $\{l_1, \dots, l_i\}$  and all the labels on the increasing chain from  $u$  to  $z$  are in  $\{l_{i+1}, \dots, l_n\}$ . Therefore,  $u$  is on the increasing chain from  $\hat{0}$  to  $z$  and, in fact,  $u = w$ . Also, we have that  $\hat{0} \leq y \leq u = w \leq z$ . We conclude that  $w$  is the greatest common lower bound for  $z$  and  $x_i$ .  $\square$

**Lemma 3.6.**  *$\hat{0} = x_0 \wedge z \leq x_1 \wedge z \leq \dots \leq x_n \wedge z = z$ , after we delete repeated elements, is the increasing chain in  $[\hat{0}, z]$ . Hence,  $(x_i \wedge z) \vee^z y$  is well-defined for  $y \leq z$ . Similarly,  $(x_i \vee y) \wedge_y z$  is well-defined.*

*Proof.* From the previous proof, we know that  $x_i \wedge z$  is the maximum element on the increasing chain from  $\hat{0}$  to  $z$  such that all labels on the increasing chain between  $\hat{0}$  and  $x_i \wedge z$  are in  $\{l_1, \dots, l_i\}$ . The first assertion follows easily from this.

Now apply Lemma 3.5 to the bounded poset  $[\hat{0}, z]$ . It has an obvious interpolating labelling induced from the interpolating labelling of  $P$ . Recall that our definition of the existence of  $(x_i \wedge z) \vee^z y$  only requires it to be well-defined in  $[\hat{0}, z]$ . The result follows.  $\square$






FIGURE 4

We conclude that the increasing maximal chain  $\hat{0} = x_0 \leq x_1 \leq \dots \leq x_n = \hat{1}$  of  $P$  is viable. It remains to show that it is left modular.

*Proof of Theorem 3.* Suppose that  $x_i$  is not left modular for some  $i$ . Then there exists some pair  $y \leq z$  such that  $(x_i \vee y) \wedge_y z > (x_i \wedge z) \vee^z y$ . Set  $x = x_i$ ,  $b = (x_i \wedge z) \vee^z y$  and  $c = (x_i \vee y) \wedge_y z$ . Observe that  $d := x \vee b \geq c$  while  $a := x \wedge c \leq b$ . So the picture is as shown in Figure 4.

By Lemma 3.3, the labels on the increasing chain from  $\hat{0}$  to  $a$  are less than or equal to  $l_i$ . Consider the increasing chain from  $a$  to  $c$ . Let  $w$  be the first element along the chain. If  $\gamma(a, w) \leq l_i$ , then by Lemma 3.3,  $w \leq x_i$ , contradicting the fact that  $a = x \wedge c$ . Thus the labels on the increasing chain from  $a$  to  $c$  are all greater than  $l_i$ . Dually, the labels on the increasing chain from  $b$  to  $d$  are less than or equal to  $l_i$ . But now, by Lemma 3.2, the labels on the increasing chain from  $b$  to  $c$  must be contained in the labels on the increasing chain from  $a$  to  $c$ , and also from  $b$  to  $d$ . But there are no such labels, implying a contradiction. We conclude that the  $x_i$  are all left modular.  $\square$

We have shown that if  $P$  is a bounded poset with an interpolating labelling  $\gamma$ , then the unique increasing maximal chain  $M$  is a viable left modular maximal chain. By Theorem 2,  $M$  then induces an interpolating EL-labeling of  $P$ . We now show that this labelling agrees with  $\gamma$  for a suitable choice of label set, which is a special case of the following proposition.

**Proposition 3.7.** *Let  $\gamma$  and  $\delta$  be two interpolating EL-labellings of a bounded poset  $P$ . If  $\gamma$  and  $\delta$  agree on the  $\gamma$ -increasing chain from  $\hat{0}$  to  $\hat{1}$ , then  $\gamma$  and  $\delta$  coincide.*

*Proof.* Let  $m : \hat{0} = w_0 \leq w_1 \leq \dots \leq w_r = \hat{1}$  be the maximal chain with the lexicographically first  $\gamma$  labelling among those chains for which  $\gamma$  and  $\delta$  disagree. Since  $m$  is not the  $\gamma$ -increasing chain from  $\hat{0}$  to  $\hat{1}$ , we can find an  $i$  such that  $\gamma(w_{i-1}, w_i) > \gamma(w_i, w_{i+1})$ . Let  $m'$  be the result of the basic replacement at  $i$  with respect to the labelling  $\gamma$ . Then the  $\gamma$ -label sequence of  $m'$  lexicographically precedes that of  $m$ , so  $\gamma$  and  $\delta$  agree on  $m'$ . But using the fact that  $\gamma$  and  $\delta$  are interpolating, it follows that they also agree on  $m$ . Thus they agree everywhere.  $\square$

#### 4. GENERALIZING SUPERSOLVABILITY

Suppose  $P$  is a bounded poset. For now, we consider the case of  $P$  being graded of rank  $n$ . We would like to define what it means for  $P$  to be *supersolvable*, thus generalizing Stanley's definition of lattice supersolvability. A definition of poset supersolvability with a different purpose appears in [12] but we would like a more general definition. In particular, we would like  $P$  to be supersolvable if and only if  $P$  has an  $S_n$  EL-labelling. For example,

the poset shown in Figure 3, while it doesn't satisfy Welker's definition, should satisfy our definition. We need to define, in the poset case, the equivalent of a sublattice generated by two chains.

Suppose  $P$  has a viable maximal chain  $M$ . Thus  $(x \vee y) \wedge_y z$  and  $(x \wedge z) \vee^z y$  are well-defined for  $x \in M$  and  $y \leq z$  in  $P$ . Given any chain  $c$  of  $P$ , we define  $R_M(c)$  to be the smallest subposet of  $P$  satisfying the following two conditions:

- (i)  $M$  and  $c$  are contained in  $R_M(c)$ ,
- (ii) If  $y \leq z$  in  $P$  and  $y$  and  $z$  are in  $R_M(c)$ , then so are  $(x \vee y) \wedge_y z$  and  $(x \wedge z) \vee^z y$  for any  $x$  in  $M$ .

**Definition 4.1.** We say that a bounded poset  $P$  is *supersolvable* with M-chain  $M$  if  $M$  is a viable maximal chain and  $R_M(c)$  is a distributive lattice for any chain  $c$  of  $P$ .

Since distributive lattices are graded, it is clear that a poset must be graded in order to be supersolvable. We now come to the main result of this section.

**Theorem 4.** *Let  $P$  be a bounded graded poset of rank  $n$ . Then the following are equivalent:*

- (1)  $P$  has an  $S_n$  EL-labelling,
- (2)  $P$  has a viable left modular maximal chain,
- (3)  $P$  is supersolvable.

*Proof.* Theorems 2 and 3 restricted to the graded case give us that  $(1) \Leftrightarrow (2)$ .

Our next step is to show that (1) and (2) together imply (3). Suppose  $P$  is a bounded graded poset of rank  $n$  with an  $S_n$  EL-labelling. Let  $M$  denote the increasing maximal chain  $\hat{0} = x_0 \triangleleft x_1 \triangleleft \cdots \triangleleft x_n = \hat{1}$  of  $P$ . We also know that  $M$  is viable and left modular and induces the same  $S_n$  EL-labelling. Given any maximal chain  $m$  of  $P$ , we define  $Q_M(m)$  to be the closure of  $m$  in  $P$  under basic replacements. In other words,  $Q_M(m)$  is the smallest subposet of  $P$  which contains  $M$  and  $m$  and which has the property that, if  $y$  and  $z$  are in  $Q_M(m)$  with  $y \leq z$ , then the increasing chain between  $y$  and  $z$  is also in  $Q_M(m)$ . It is shown in [8, Proof of Thm. 1] that  $Q_M(m)$  is a distributive lattice. There  $P$  is a lattice but the proof of distributivity doesn't use this fact. Now consider  $R_M(c)$ . We will show that there exists a maximal chain  $m$  of  $P$  such that  $R_M(c) = Q_M(m)$ . Let  $m$  be the maximal chain of  $P$  which contains  $c$  and which has increasing labels between successive elements of  $c \cup \{\hat{0}, \hat{1}\}$ . The only idea we need is that, for  $y \leq z$  in  $P$ , the increasing chain from  $y$  to  $z$  is given by  $y = (x_0 \vee y) \wedge_y z \leq (x_1 \vee y) \wedge_y z \leq \cdots \leq (x_n \vee y) \wedge_y z = z$ , where we delete repeated elements. This follows from Lemma 2.4 since the induced left modular chain in  $[y, z]$  has increasing labels. It now follows that  $R_M(c) = Q_M(m)$ , and hence  $R_M(c)$  is a distributive lattice.

Finally, we will show that  $(3) \Rightarrow (2)$ . We suppose that  $P$  is a bounded supersolvable poset with M-chain  $M$ . Suppose  $y \leq z$  in  $P$  and let  $c$  be the chain  $y \leq z$ . For any  $x$  in  $M$ ,  $x \vee y$  is well-defined in  $P$  (because  $M$  is assumed to be viable) and equals the usual join of  $x$  and  $y$  in the lattice  $R_M(c)$ . The same idea applies to  $x \wedge z$ ,  $(x \vee y) \wedge_y z$  and  $(x \wedge z) \vee^z y$ . Since  $R_M(c)$  is distributive, we have that

$$(x \vee y) \wedge_y z = (x \vee y) \wedge z = (x \wedge z) \vee (y \wedge z) = (x \wedge z) \vee y = (x \wedge z) \vee^z y$$

in  $R_M(c)$  and so  $M$  is left modular in  $P$ . □

*Remark 4.2.* We know from Theorem 1 that a graded lattice of rank  $n$  is supersolvable if and only if it has an  $S_n$  EL-labelling. Therefore, it follows from Theorem 4 that the definition of a supersolvable poset restricts to graded lattices to give the usual definition.

However, suppose  $P$  is a graded lattice with maximal chain  $M$ . We should note that it is not obvious, and may not even be true, that for a given chain  $c$  of  $P$ ,  $R_M(c)$  equals the sublattice of  $P$  generated by  $M$  and  $c$ .

*Remark 4.3.* The argument above for the equality of  $R_M(c)$  and  $Q_M(m)$  holds even if  $P$  is not graded. However, in the ungraded case, it is certainly not true that  $Q_M(m)$  is distributive. The search for a full generalization of Theorem 1 thus leads us to ask what can be said about  $Q_M(m)$  in the ungraded case. Is it even a lattice? Can we say anything even in the case that  $P$  is a lattice?

# ACKNOWLEDGMENTS

The authors would like to thank Andreas Blass, Bruce Sagan, Richard Stanley and Volkmar Welker for helpful comments.

# REFERENCES

- [1] G. Birkhoff, *Lattice Theory (3rd. Ed.)*, Amer. Math. Soc., Providence, RI. (1967).
- [2] A. Björner, Shellable and Cohen-Macaulay partially ordered sets, *Trans. Amer. Math. Soc.* **260** (1980), 159-183.
- [3] A. Björner and M. Wachs, Shellable non-pure complexes and posets. I., *Trans. Amer. Math. Soc.* **348** (1996), 1299-1327.
- [4] A. Björner and M. Wachs, Shellable non-pure complexes and posets. II., *Trans. Amer. Math. Soc.* **349** (1997), 3945-3975.
- [5] A. Blass and B. Sagan, Möbius functions of lattices, *Adv. Math.* **127** (1997), 94-123.
- [6] L. S.-C. Liu, Left-modular elements and edge labellings, Ph. D. thesis, Michigan State University, 1999.
- [7] S.-C. Liu and B. Sagan, Left-modular elements of lattices, *J. Combin. Theory Ser. A* **91** (2000), 369-385.
- [8] P. McNamara, EL-labelings, supersolvability and 0-Hecke algebra actions on posets, *J. Combin. Theory Ser. A* **101** (2003), 69-89.
- [9] R. Stanley, Ordered structures and partitions, *Mem. Amer. Math. Soc.* **119** (1972).
- [10] R. Stanley, Supersolvable lattices, *Algebra Universalis* **2** (1972), 197-217.
- [11] R. Stanley, "Enumerative Combinatorics," vol. 1, Wadsworth & Brooks/Cole, Monterey, CA, 1986; second printing, Cambridge University Press, Cambridge/New York, 1997.
- [12] V. Welker, Shellability in the lattice of subgroups of a finite group, in "Jerusalem Combinatorics '93", Contemp. Math. 178, Amer. Math. Soc., Providence, RI, 1994, pp. 335-360.

DEPARTMENT OF MATHEMATICS 2-342, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139, U.S.A.

*E-mail address:* mcnamara@math.mit.edu

FIELDS INSTITUTE, 222 COLLEGE ST., TORONTO, ON, M5T 3J1 CANADA

*E-mail address:* hthomas@fields.utoronto.ca