


# Environmental Science Processes & Impacts

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: C. Lihl, B. Heckel, A. Grzybkowska, A. Dybala-Defratyka, V. Ponsin, C. Torrentó, D. Hunkeler and M. Elsner, *Environ. Sci.: Processes Impacts*, 2020, DOI: 10.1039/C9EM00503J.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

# Compound-Specific Chlorine Isotope Fractionation in Biodegradation of Atrazine<sup>†</sup>

Christina Lihl<sup>a,b</sup>, Benjamin Heckel<sup>a,b</sup>, Anna Grzybkowska<sup>c</sup>, Agnieszka Dybala-Defratyka<sup>c</sup>,  
Violaine Ponsin<sup>d,e</sup>, Clara Torrentó<sup>d,f</sup>, Daniel Hunkeler<sup>d</sup>, Martin Elsner<sup>a,b,\*</sup>

<sup>a</sup> Institute of Groundwater Ecology, Helmholtz Zentrum München, Ingolstädter Landstraße 1, 85764 Neuherberg, Germany

<sup>b</sup> Chair of Analytical Chemistry and Water Chemistry, Technical University of Munich, Marchioninstraße 17, 81377 Munich, Germany

<sup>c</sup> Institute of Applied Radiation Chemistry, Faculty of Chemistry, Lodz University of Technology, Zeromskiego 116, 90-924 Lodz, Poland

<sup>d</sup> Centre of Hydrogeology and Geothermics (CHYN), University of Neuchâtel, 2000 Neuchâtel, Switzerland

<sup>e</sup> Département des sciences de la Terre et de l'atmosphère, Université du Québec à Montréal, 201 avenue du Président Kennedy, Montréal, QC, Canada

<sup>f</sup> Grup MAiMA, Departament de Mineralogia, Petrologia i Geologia Aplicada, Facultat de Ciències de la Terra, Universitat de Barcelona (UB), C/ Martí i Franquès s/n, 08028, Barcelona, Spain.

\* Corresponding Author: Phone: +49 89/2180-78231. E-mail: [m.elsner@tum.de](mailto:m.elsner@tum.de)

<sup>†</sup> Electronic Supplementary Information (ESI) available: Information concerning the HPLC temperature programs, two Figures and one Table illustrating the GC-qMS method optimization for chlorine analysis, one Table illustrating the method comparison of the GC-qMS for chlorine analysis, one Figure and one Table considering H-abstraction during chlorine CSIA, two Figures illustrating the results of HPLC concentration analysis.

KEYWORDS: hydrolysis, oxidative dealkylation, compound-specific isotope analysis, chlorine isotope effect, *Arthrobacter*, *Rhodococcus*

### Environmental Significance:

Atrazine is an important chlorinated micropollutant. Although degradable *via* different pathways (dealkylation and hydrolytic dechlorination), it is often recalcitrant and persists in groundwater. To assess and understand its degradation pathways, compound-specific carbon and nitrogen isotope analysis has been advanced, but information from chlorine isotope fractionation has been missing until today. This study explores the added benefit of chlorine isotope fractionation as indicator of natural atrazine transformation. Together with carbon and nitrogen isotope analysis, this enables a multi-element approach which can improve source identification and differentiation of microbial transformation pathways in the environment.

### ABSTRACT

Atrazine is a frequently detected groundwater contaminant. It can be microbially degraded by oxidative dealkylation or by hydrolytic dechlorination. Compound-specific isotope analysis is a powerful tool to assess its transformation. In previous work, carbon and nitrogen isotope effects were found to reflect these different transformation pathways.

1  
2  
3  
4 However, chlorine isotope fractionation could be a particularly sensitive indicator of  
5  
6 natural transformation since chlorine isotope effects are fully represented in the molecular  
7  
8 average while carbon and nitrogen isotope effects are diluted by non-reacting atoms.  
9  
10 Therefore, this study explored chlorine isotope effects during atrazine hydrolysis with  
11  
12 *Arthrobacter aurescens* TC1 and oxidative dealkylation with *Rhodococcus* sp. NI86/21.  
13  
14 Dual element isotope slopes of chlorine vs. carbon isotope fractionation ( $\Delta^{Arthro}_{Cl/C} = 1.7$   
15  
16  $\pm 0.9$  vs.  $\Delta^{Rhodo}_{Cl/C} = 0.6 \pm 0.1$ ) and chlorine vs. nitrogen isotope fractionation ( $\Delta^{Arthro}_{Cl/N} = -$   
17  
18  $1.2 \pm 0.7$  vs.  $\Delta^{Rhodo}_{Cl/N} = 0.4 \pm 0.2$ ) provided reliable indicators of different pathways.  
19  
20 Observed chlorine isotope effects in oxidative dealkylation ( $\epsilon_{Cl} = -4.3 \pm 1.8$  ‰) were  
21  
22 surprisingly large, whereas in hydrolysis ( $\epsilon_{Cl} = -1.4 \pm 0.6$  ‰) they were small, indicating  
23  
24 that C-Cl bond cleavage was not the rate-determining step. This demonstrates the  
25  
26 importance of constraining expected isotope effects of new elements before using the  
27  
28 approach in the field. Overall, the triple element isotope information brought forward here  
29  
30 enables a more reliable identification of atrazine sources and degradation pathways.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## INTRODUCTION

The herbicide atrazine has been used in agriculture to inhibit growth of broadleaf and grassy weeds<sup>1</sup>. In the U.S. atrazine was the second most commonly used herbicide in 2012 and is still in use today<sup>2</sup>. In the European Union atrazine was banned in 2004<sup>3</sup>, but together with its metabolites it is still frequently detected at high concentrations in groundwater<sup>4, 5</sup>. The massive and widespread use has led to a wide-ranging presence of atrazine in the environment, which can have harmful effects on living organisms and humans<sup>6</sup>. Therefore, the environmental fate of atrazine is of significant concern and much attention has been directed at detecting and enhancing its natural biodegradation. However, assessing microbial degradation of atrazine in the environment is challenging with conventional methods like concentration analysis. Sorption and remobilization of the parent compound and its metabolites, as well as further transformation of the metabolites inevitably lead to fluctuations in concentrations<sup>7-10</sup>, which make it difficult to assess the net extent of atrazine degradation in the field.

In recent years, compound-specific isotope analysis (CSIA) has been proposed as an alternative approach to detect and quantify the degradation of atrazine<sup>11-13</sup>.

In contrast to, and complementary to traditional methods, CSIA informs about transformation without the need to detect metabolites. The reason is that during (bio)chemical transformations molecules with heavy isotopes are typically enriched in the remaining substrate since their bonds are more stable and, therefore, usually react slower

1  
2  
3  
4 than molecules containing light isotopes (normal kinetic isotope effect). The ratios of  
5  
6 heavy to light isotopes (e.g.  $^{13}\text{C}/^{12}\text{C}$  for carbon) in the remaining substrate, therefore,  
7  
8 change during transformation. Observing such changes can be used as direct (and  
9  
10 concentration-independent) indicator of degradation<sup>14, 15</sup>.

11  
12 Isotope values are reported in the  $\delta$ -notation relative to an international reference  
13  
14 material, e.g. for carbon<sup>14, 15</sup>:

$$\delta^{13}\text{C} = \left[ \left( \frac{^{13}\text{C}}{^{12}\text{C}} \right)_{\text{Sample}} - \left( \frac{^{13}\text{C}}{^{12}\text{C}} \right)_{\text{Reference}} \right] / \left( \frac{^{13}\text{C}}{^{12}\text{C}} \right)_{\text{Reference}} \quad (1)$$


15  
16 The magnitude of the degradation-induced isotope fractionation depends on different  
17  
18 factors, which can make isotope ratios of specific elements particularly attractive to  
19  
20 observe degradation-induced isotope fractionation. To this end, first, an element needs  
21  
22 to be directly involved in the (bio)chemical reaction. For example, a carbon isotope effect  
23  
24 would be quite generally expected in organic molecules, whereas a chlorine isotope effect  
25  
26 would be primarily expected if a C-Cl bond is cleaved. Second, isotope fractionation  
27  
28 depends on the underlying kinetic isotope effect (see above), but also on the extent to  
29  
30 which this effect is represented in the molecular average isotope fractionation described  
31  
32 by the enrichment factor  $\epsilon$  (see below). Atrazine, for example, contains only one chlorine  
33  
34 atom but eight carbon and five nitrogen atoms. Hence, chlorine isotope effects at the  
35  
36 reacting position are fully represented in the molecular average, whereas position-specific  
37  
38 carbon and nitrogen isotope effects are diluted by non-reacting atoms<sup>14, 15</sup>.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Most of the publications studying the chemical mechanisms of abiotic and microbial  
5  
6 atrazine degradation have focused on the analysis of carbon ( $^{13}\text{C}/^{12}\text{C}$ ) and nitrogen  
7  
8 ( $^{15}\text{N}/^{14}\text{N}$ ) isotope fractionation. Thereby,  $\epsilon$ -values in the range of -5.4 ‰ to -1.8 ‰ for  
9  
10 carbon and -1.9 ‰ to 3.3 ‰ for nitrogen were observed<sup>9, 10, 16, 17</sup>. Chlorine isotope effects  
11  
12 for microbial atrazine degradation were so far not reported due to analytical challenges<sup>18</sup>:  
13  
14 Until recently<sup>19, 20</sup>, suitable methods were not available for chlorine isotope analysis of  
15  
16 atrazine. However, from the magnitude of chlorine isotope effects observed for  
17  
18 dechlorination of trichloroethenes (-5.7 ‰ to -3.3 ‰, where intrinsic isotope effects are  
19  
20 diluted by a factor of three<sup>21</sup>), very large chlorine enrichment factors  $\epsilon_{\text{Cl}}$  (-8 ‰ to -10 ‰ or  
21  
22 even larger) could potentially occur for a C-Cl bond cleavage in atrazine. For example,  
23  
24 enzymatic hydrolysis of the structural homologue ametryn (atrazine structure with a -  
25  
26  $\text{SCH}_3$  instead of a -Cl group) yielded a sulfur isotope enrichment factor  $\epsilon_{\text{S}}$  of -  
27  
28 14.7 ‰  $\pm$  1.0 ‰<sup>17</sup>. If the cleavage of carbon-chlorine bonds is involved in the rate-  
29  
30 determining step of a (bio)transformation, chlorine isotope effects could, therefore, enable  
31  
32 a particularly sensitive detection of natural transformation processes by compound-  
33  
34 specific (i.e., molecular average) isotope analysis.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45


46 The measurement of chlorine isotope fractionation is attractive for yet another reason –  
47  
48 multiple element isotope analysis bears potential for a better distinction of sources and  
49  
50 transformation pathways. From isotope analysis of one element alone, it is difficult to  
51  
52 distinguish sources of a particular compound, or competing transformation pathways that  
53  
54  
55  
56  
57  
58  
59  
60

may lead to metabolites of different toxicity<sup>15</sup>. For example, two different microbial transformation pathways can lead to the degradation of atrazine in the environment. Hydrolysis forms the nontoxic dehalogenated product 2-hydroxyatrazine (HAT) whereas oxidative dealkylation degrades atrazine to the still herbicidal products desethyl- (DEA) or desisopropylatrazine (DIA)<sup>22, 23</sup>. Prominent examples for microorganisms catalyzing these pathways are *Arthrobacter aurescens* TC1 and *Rhodococcus* sp. NI86/21 (see Figure 1). *A. aurescens* TC1 was directly isolated from an atrazine-contaminated soil<sup>24</sup>. By expressing the enzyme TrzN, it is capable of performing hydrolysis of atrazine<sup>24, 25</sup>. *Rhodococcus* sp. NI86/21 uses a cytochrome P450 system for catalyzing oxidative dealkylation of atrazine<sup>26</sup>.

#### *Arthrobacter aurescens* TC1 (TrzN): Hydrolysis



#### *Rhodococcus* sp. NI86/21 (Cytochrome P450): Oxidative Dealkylation



**Figure 1.** Microbial degradation of atrazine by *Arthrobacter aurescens* TC1 and *Rhodococcus* sp. NI86/21 (adapted from Meyer et al.<sup>9</sup> and Meyer & Elsner<sup>10</sup>).

1  
2  
3  
4 For these two pathways, carbon isotope fractionation was very similar, but significant  
5  
6 differences were observed in nitrogen isotope effects<sup>9, 10, 16, 17</sup>. Plotting the changes of  
7  
8 isotope ratios of these two elements relative to each other results in the regression  
9  
10 slope  $\Lambda$  for carbon and nitrogen<sup>27, 28</sup>

$$\Lambda_{C/N} = \Delta\delta^{15}\text{N} / \Delta\delta^{13}\text{C} \approx \varepsilon_{\text{N}} / \varepsilon_{\text{C}} \quad (2)$$

11 Hence, dual element (C, N) isotope trends for oxidative dealkylation of atrazine with  
12  
13 *Rhodococcus* sp. NI86/21 ( $\Lambda^{Rhodo}_{C/N} = 0.4 \pm 0.1$ )<sup>16</sup> were significantly different compared  
14  
15 to hydrolysis with *A. aurescens* TC1 ( $\Lambda^{Arthro}_{C/N} = -0.6 \pm 0.1$ )<sup>9</sup> offering an opportunity to  
16  
17 distinguish atrazine degradation pathways in the field. However, in environmental  
18  
19 assessments it is advantageous to have isotopic information of as many elements as  
20  
21 possible in order to distinguish degradation pathways and sources at the same time<sup>29-31</sup>.  
22  
23 Therefore, information from a third element, chlorine, would be highly valuable. Also on  
24  
25 the mechanistic end, information gained from a change in the chlorine isotope value could  
26  
27 lead to a more reliable differentiation of transformation pathways and contribute to a better  
28  
29 mechanistic understanding of the underlying chemical reaction<sup>31</sup>. Along these lines, triple  
30  
31 element (3D) isotope analysis was already accomplished for chlorinated alkanes<sup>31, 32</sup> and  
32  
33 alkenes<sup>33, 34</sup>.

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Until now, however, compound-specific chlorine isotope analysis has not been accessible so that chlorine isotope ratio changes for hydrolysis of atrazine have only been analyzed in abiotic systems or via computational calculations<sup>35, 36</sup>. For oxidative

1  
2  
3  
4 dealkylation, chlorine isotope effects have, so far, not been studied. Recently a GC-qMS  
5  
6 method for chlorine isotope analysis of atrazine has been brought forward<sup>20</sup> which offers  
7  
8 the opportunity to enable deeper mechanistic insights into its transformation processes.  
9  
10 Therefore, our objective was to analyze carbon, nitrogen and chlorine isotope effects  
11  
12 associated with the biodegradation of atrazine via hydrolysis with *A. aurescens* TC1 and  
13  
14 via oxidative dealkylation with *Rhodococcus* sp. NI86/21. In addition, we computationally  
15  
16 predicted the chlorine isotope effect associated with hydrolysis and oxidative dealkylation  
17  
18 for comparison. Further, we evaluated whether the additional information from chlorine  
19  
20 isotope fractionation is a particularly sensitive indicator for transformation processes and  
21  
22 whether it can confirm previously proposed mechanisms of different pathways. With this  
23  
24 study, we bring forward information about degradation-induced chlorine isotope  
25  
26 fractionation of atrazine as a basis to apply triple element (3D) isotope analysis in  
27  
28 environmental assessments.

## 38 MATERIAL & METHODS

39  
40  
41 **Bacterial strains and cultivation.** *A. aurescens* strain TC1 was grown in mineral salt  
42  
43 medium supplemented with approx. 20 mg/L of atrazine according to the protocol of  
44  
45 Meyer et al.<sup>9</sup> Likewise, *Rhodococcus* sp. strain NI86/21 was cultivated in autoclaved  
46  
47 nutrient broth (8 g/L, Difco™) with approx. 20 mg/L of atrazine according to the protocol  
48  
49 of Meyer et al.<sup>16</sup>. In the late-exponential growth phase the strains were harvested via  
50  
51 centrifugation (4000 rpm, 15 min). For the start of the degradation experiments, cell  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 pellets of each strain were transferred to 400 mL fresh media and atrazine was added to  
4  
5  
6 achieve a starting concentration of 20 mg/L. All experiments were performed in triplicate  
7  
8  
9 at 21 °C on a shaker at 150 rpm. Control experiments, which were performed without the  
10  
11 bacterial strains, did not show any degradation of atrazine.

12  
13  
14  
15 **Concentration measurements via HPLC.** The process of atrazine degradation was  
16  
17 monitored by concentration measurements. For analysis, 1 mL samples were taken and  
18  
19 filtered through a 0.22 µm filter. Atrazine and its degradation products were directly  
20  
21 analyzed using a Shimadzu UHPLC-20A system, which was equipped with an ODS  
22  
23 column 30 (Ultracarb 5 µM, 150 × 4.6 mm, Phenomenex). After sample injection (10 µL)  
24  
25 an adequate gradient program (see SI) was used for compound separation. The oven  
26  
27 temperature was set to 45 °C and the compounds were detected by their UV absorbance  
28  
29 at 222 nm. Quantitation was performed by the software “Lab Solutions” based on internal  
30  
31 calibration curves.  
32  
33  
34  
35  
36  
37  
38

39  
40 **Preparation of samples for isotope analysis.** According to the protocol of Meyer et al.<sup>9</sup>  
41  
42 between 10 and 260 mL of sample were taken for isotope analysis of atrazine at every  
43  
44 sampling event. After centrifugation (15 min, 4000 rpm) the supernatant was collected in  
45  
46 a new vial. Subsequently, samples were extracted by adding dichloromethane (5-130 mL)  
47  
48 and shaking the vial for at least 20 min. The sample extracts were dried at room  
49  
50 temperature under the fume hood. Afterwards, the dried extracts were dissolved in ethyl  
51  
52 acetate to a final atrazine concentration of approx. 200 mg/L.  
53  
54  
55  
56  
57  
58  
59  
60

**Isotope analysis of carbon and nitrogen.** The protocol for isotope analysis of carbon and nitrogen was adapted from the studies of Meyer et al.<sup>9, 16</sup>. A TRACE GC Ultra gas chromatograph hyphenated with a GC-III combustion interface and coupled to a Finnigan MAT253 isotope ratio mass spectrometer (GC-C-IRMS, all Thermo Fisher Scientific) was used. Each sample was analyzed in triplicate. Sample injection (2-3  $\mu\text{L}$ ) was performed by a Combi-PAL autosampler (CTC Analysis). The injector had a constant temperature of 220 °C, was equipped with an “A” type packed liner for large volume injections (GL Sciences) and was operated for 1 min in splitless and then in split mode (split ratio 1:10) with a flow rate of 1.4 mL/min. For peak separation, the GC oven was equipped with a DB-5 MS column (30 m  $\times$  0.25 mm, 1  $\mu\text{m}$  film thickness, Agilent). The temperature program of the oven started at 65 °C (held for 1 min), ramped at 20 °C/min to 180 °C (held for 10 min) and ramped again at 15 °C/min to 230 °C (held for 8 min). In the combustion interface, a GC Isolink II reactor (Thermo Fisher Scientific) was installed, which was operated at a temperature of 1000 °C. After combustion of the analytes to  $\text{CO}_2$  and subsequent reduction of any nitrogen oxides, the compounds were analyzed as  $\text{CO}_2$  for carbon and  $\text{N}_2$  for nitrogen isotope measurements. Three pulse of  $\text{CO}_2$  or  $\text{N}_2$ , respectively, were introduced at the beginning and at the end of each run as monitoring gas. Beforehand, these monitoring gases were calibrated against RM8563 ( $\text{CO}_2$ ) and NSVEC ( $\text{N}_2$ ), which were supplied by the International Atomic Energy Agency (IAEA). The analytical uncertainty  $2\sigma$  was  $\pm 0.5\text{‰}$  for carbon isotope values and  $\pm 1.0\text{‰}$  for nitrogen isotope values.

**Isotope analysis of chlorine.** For chlorine isotope analysis of atrazine, a 7890A gas chromatograph coupled to a 5975C quadrupole mass spectrometer (GC-qMS, both Agilent) was used. Sample injection (2  $\mu\text{L}$ ) was performed by a Pal Combi-xt autosampler (CTC Analysis). For the injector and the GC oven, the same parameters as for carbon and nitrogen isotope analysis were used with the exception that a different liner type, a “FocusLiner” (SGE), was used. The ion source had a constant temperature of 230  $^{\circ}\text{C}$  and the quadrupole of 150  $^{\circ}\text{C}$ . Prior to sample analysis, the method of Ponsin et al.<sup>20</sup> was tested and optimized for our instrument (see details in SI). Chlorine isotope ratios were evaluated by monitoring the mass-to-charge ratio  $m/z$  of 202/200. Standards and samples were measured ten times each and uncertainties were reported as standard deviation. Results were only evaluated if the peak areas of samples were inside a defined linearity range (peak area of  $1.2 \times 10^8 - 3.0 \times 10^8$  for  $m/z$  200). Inside the linearity range, the determined precision of the method is associated with a maximal deviation of  $\pm 1.1$  ‰. For analysis, the samples were diluted with ethyl acetate to a final concentration of approx. 75 mg/L and measured with a dwell time of 100 ms. Correction of the chlorine isotope values relative to Standard Mean Ocean Chloride (SMOC) was performed by an external two-point calibration with characterized standards of atrazine (Atr #4  $\delta^{37}\text{Cl} = -0.89$  ‰ and Atr #11  $\delta^{37}\text{Cl} = +3.59$  ‰)<sup>37</sup>. To this end, the standards were measured at the beginning, in between and at the end of each sequence.

**Evaluation of stable isotope fractionation.** Determination of isotope enrichment factors  $\epsilon$  was achieved by the Rayleigh equation, which describes the gradual enrichment of the residual substrate fraction  $f$  with molecules containing heavy isotopes, as expressed by isotope values according to eq. 1<sup>15, 38</sup>. For example, for chlorine:

$$\ln [ (\delta^{37}\text{Cl} + 1) / (\delta^{37}\text{Cl}_0 + 1) ] = \epsilon_{\text{Cl}} \cdot \ln f \quad (3)$$

Here  $\delta^{37}\text{Cl}_0$  refers to the chlorine isotope value at the starting point ( $t = 0$ ) of an experiment. Regression slopes  $\Lambda$  of dual element isotope plots were obtained by plotting isotope ratios of two different elements against each other, e.g. carbon vs. nitrogen (see eq. 2). The uncertainties of the calculated  $\epsilon$ -values and  $\Lambda$ -values are reported as 95 % confidence intervals (CI). Furthermore, (apparent) kinetic isotope values, (A)KIE<sub>Cl</sub>, that express the ratio of reaction rates  $^{35}\text{k}$  and  $^{37}\text{k}$  of heavy and light isotopologues, respectively,

$$\text{KIE}_{\text{Cl}} = ^{35}\text{k} / ^{37}\text{k} \quad (4)$$

were calculated according to Elsner et al.<sup>15</sup> by converting  $\epsilon_{\text{Cl}}$ -values into (A)KIE<sub>Cl</sub> and taking into account that atrazine contains only one chlorine atom ( $n = 1$ ):

$$(\text{A})\text{KIE}_{\text{Cl}} = 1 / (n \times \epsilon_{\text{Cl}} + 1) \quad (5)$$


**Prediction of chlorine kinetic isotope effects during oxidative dealkylation and hydrolysis of atrazine.** In the computational part of the study, we considered hydrogen atom transfer and hydride transfer taking place at the  $\alpha$ -position of the ethyl side chain of the atrazine

1  
2  
3  
4 molecule in the oxidative dealkylation reaction promoted by permanganate and the  
5  
6 hydronium ion, respectively. Furthermore, we considered hydrolysis under  
7  
8 acidic/enzymatic, neutral and alkaline conditions. All molecular structures and analytical  
9  
10 vibrational frequencies for involved reactant complexes and transition states were taken  
11  
12 from a previous study<sup>16</sup>. Chlorine kinetic isotope effects on dealkylation were calculated  
13  
14 using the complete Bigeleisen equation<sup>39</sup> implemented in the ISOEFF program<sup>40</sup> at  
15  
16 300 K. Additional isotope effects predictions for hydrolysis under acidic as well as neutral  
17  
18 conditions were performed following the previous computational protocol<sup>36</sup>. The tunneling  
19  
20 contributions to the overall kinetic isotope effect were omitted.  
21  
22  
23  
24  
25  
26  
27  
28

## RESULTS & DISCUSSION

29  
30  
31 **Observation of normal chlorine isotope effects in biotic hydrolysis and oxidative**  
32 **dealkylation.** Atrazine degradation by *A. aurescens* TC1 resulted in the metabolite 2-  
33  
34 hydroxyatrazine, whereas the metabolites DEA and DIA were observed for  
35  
36 *Rhodococcus* sp. NI86/21 (see Figure S4 and S5 in the SI). Detection of these expected  
37  
38 degradation products (Figure 1) demonstrates that hydrolysis and oxidative dealkylation  
39  
40 were the underlying biochemical reactions during atrazine degradation, respectively. In  
41  
42 both biodegradation experiments – biotic hydrolysis with *A. aurescens* TC1 and oxidative  
43  
44 dealkylation with *Rhodococcus* sp. NI86/21 – normal chlorine isotope fractionation was  
45  
46 observed (see Figure 2A). In the three replicates of hydrolytic degradation by  
47  
48 *A. aurescens* TC1 90 %, 90 % and 60 % transformation of atrazine was reached after  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 approx. 26 h, respectively (see SI, Figure S4). Evaluation of  $\delta^{37}\text{Cl}$  values during biotic  
5  
6 hydrolysis according to Equation 3 resulted in a small normal isotope effect of  $\epsilon_{\text{Cl}} = -1.4$   
7  
8  $\pm 0.6$  ‰. In oxidative dealkylation with *Rhodococcus* sp. NI86/21 approx. 90 %  
9  
10 degradation was reached after approx. 186 h in all three replicates (see SI, Figure S5).  
11  
12 Evaluation of changes in chlorine isotope ratios resulted in a surprisingly large normal  
13  
14 isotope effect of  $\epsilon_{\text{Cl}} = -4.3 \pm 1.8$  ‰ considering that the C-Cl bond is not broken during  
15  
16 the reaction (see Figure 1). In a next step, carbon and nitrogen isotope effects were  
17  
18 therefore analyzed to confirm whether the same reactions mechanisms are at work as  
19  
20 observed in previous studies<sup>9, 16</sup>.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Figure 2.** Isotope fractionation of (A) chlorine, (B) nitrogen and (C) carbon during microbial degradation of atrazine by *A. aurescens* TC1 (red) and *Rhodococcus* sp.


1  
2  
3  
4 NI86/21 (blue) and corresponding enrichment factors  $\epsilon$  evaluated according to eq. 3. (The  
5  
6 95 % confidence intervals are given as values and as black lines).  
7

8  
9 **Observed carbon and nitrogen isotope fractionation is consistent with previous studies.**

10  
11 Carbon and nitrogen isotope fractionation for atrazine degradation by *A. aurescens* TC1  
12 and *Rhodococcus* sp. NI86/21 was consistent with previous studies: Both experiments  
13  
14 showed significant changes in isotope ratios (see Figure 2B and C). For hydrolysis with  
15  
16 *A. aurescens* TC1, an inverse nitrogen isotope effect ( $\epsilon_N = 2.3 \pm 0.3 \text{ ‰}$ ) and a normal  
17  
18 carbon isotope effect ( $\epsilon_C = -3.7 \pm 0.4 \text{ ‰}$ ) were observed, which were slightly smaller  
19  
20 compared to the results of a former publication of Meyer et al. ( $\epsilon_N = 3.3 \pm 0.4 \text{ ‰}$ ,  $\epsilon_C = -$   
21  
22  $5.4 \pm 0.6 \text{ ‰}$ )<sup>9</sup>, but gave the same dual element isotope plot ( $\Delta^{Arthro}_{C/N} = -0.6 \pm 0.1$ )  
23  
24 confirming that the same mechanism was at work (see Figure 3A).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

1  
2  
3  
4 fractionation effect. The obtained regression slope of  $\Lambda^{Rhodo}_{C/N} = 0.7 \pm 0.1$  in this study is  
5  
6 slightly larger than the previously reported regression slopes ( $\Lambda^{Rhodo}_{C/N} = 0.4 \pm 0.1$ )<sup>10, 16</sup>  
7  
8 which may again be explained by the small difference in average nitrogen isotope effects.  
9  
10 Also here, however, the similar dual element isotope trend confirms that in this study  
11  
12 atrazine was transformed by the same mechanism as in Meyer et al.<sup>16</sup> leading to the  
13  
14 observed oxidative dealkylation products by *Rhodococcus* sp. NI86/21.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Figure 3.** Isotope effects in microbial degradation of atrazine by *A. aureescens* TC1 (red) and *Rhodococcus* sp. NI86/21 (blue) resulting in dual element isotope plots. (The 95 % confidence intervals are given as values and as black lines next to the regression slopes).

1  
2  
3  
4 (A) Regression slopes of nitrogen and carbon isotope values ( $\Lambda_{C/N}$ ). (B) Regression  
5  
6 slopes of chlorine and carbon isotope values ( $\Lambda_{Cl/C}$ ). (C) Regression slopes of chlorine  
7  
8 and nitrogen isotope values ( $\Lambda_{Cl/N}$ ).  
9

10  
11  
12 **Multi-element isotope approach.** Results of chlorine isotope analysis were combined  
13  
14 with data for carbon and nitrogen isotope measurements in dual element isotope plots  
15  
16 (see Figure 3B and C). For hydrolysis with *A. aurescens* TC1 regression slopes of  
17  
18  $\Lambda^{Arthro}_{Cl/C} = 1.7 \pm 0.9$  and  $\Lambda^{Arthro}_{Cl/N} = -1.2 \pm 0.7$  were obtained. Oxidative dealkylation by  
19  
20 *Rhodococcus* sp. NI86/21 resulted in regression slopes of  $\Lambda^{Rhodo}_{Cl/C} = 0.6 \pm 0.1$  and  
21  
22  $\Lambda^{Rhodo}_{Cl/N} = 0.4 \pm 0.2$ . Since the dual element isotope plots of chlorine and carbon and of  
23  
24 chlorine and nitrogen provide significantly different regression slopes for the respective  
25  
26 elements, they can provide an additional line of evidence to differentiate the two  
27  
28 degradation mechanisms of atrazine from each other.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39  
40 **Surprising mechanistic evidence from chlorine isotope effects.** For degradation with  
41  
42 *A. aurescens* TC1, rather small chlorine isotope fractionation was observed ( $\epsilon_{Cl} = -1.4$   
43  
44  $\pm 0.6$  ‰) despite the fact that the chlorine is cleaved off during hydrolysis (see Figure 1).  
45  
46 For oxidative dealkylation with *Rhodococcus* sp. NI86/21, the chlorine is not cleaved off  
47  
48 (see Figure 1), therefore, no or just a small chlorine isotope effect was expected.  
49  
50 However, here more pronounced chlorine isotope fractionation was observed ( $\epsilon_{Cl} = -4.3$   
51  
52  $\pm 1.8$  ‰).  
53  
54  
55  
56  
57  
58  
59  
60

The corresponding apparent kinetic isotope effects ( $AKIE_{Cl}$ , see Table 1) were compared to the  $AKIE_{Cl}$  values of other studies focusing on the same degradation mechanisms. In addition, the  $AKIE_{Cl}$  values were compared to the theoretical maximum Streitwieser Limit associated with the cleavage of a C-Cl bond ( $KIE_{Cl} = 1.02$ )<sup>41-43</sup> and to the predictions of computational calculations (Table 2).

**Table 1.**  $AKIE_{Cl}$  values associated with atrazine degradation.

Mechanism	$AKIE_{Cl}$	Study
<b>Experimental Data</b>		
abiotic alkaline hydrolysis (21 °C)	$1.0069 \pm 0.0005$	Dybala-Defratyka et al. <sup>35</sup>
abiotic alkaline hydrolysis (50 °C), microbial hydrolysis ( <i>A. aurescens</i> TC1)	$1.0009 \pm 0.0006$	Grzybkowska et al. <sup>36</sup>
microbial hydrolysis ( <i>A. aurescens</i> TC1)	$1.0014 \pm 0.0006^*$	this study
microbial dealkylation ( <i>Rhodococcus</i> sp. NI86/21)	$1.0043 \pm 0.0018^*$	this study
<b>Computational Data</b>		
abiotic acidic/enzymatic hydrolysis (transition state 1)	range of 1.0002 to 1.0011	Grzybkowska et al. <sup>36</sup>
abiotic acidic/enzymatic hydrolysis (transition state 2)	1.0099	this study
abiotic neutral hydrolysis	1.0045	this study
abiotic alkaline hydrolysis	range of 1.0003 to 1.0014	Grzybkowska et al. <sup>36</sup>
enzymatic hydrolysis	range of 0.9996 to 1.0003	Szatkowski et al. <sup>44</sup>
abiotic dealkylation (hydrogen atom transfer by $MnO_4^-$ )	0.9999	this study
abiotic dealkylation (hydride transfer by $H_3O^+$ )	0.9997	this study

\* calculated according to eq. 5

1  
2  
3  
4 For microbial hydrolysis of atrazine an experimental  $AKIE_{Cl}^{Arthro}$  value of 1.0014  
5  
6  $\pm 0.0006$  was calculated (see Table 1). Dybala-Defratyka et al.<sup>35</sup> reported a more  
7  
8 pronounced  $AKIE_{Cl}^{alk.hydro}$  value of  $1.0069 \pm 0.0005$  (see Table 1). However, that study<sup>35</sup>  
9  
10 was conducted in an abiotic alkaline solution at 21 °C so that another hydrolysis pathway  
11  
12 was involved. Newer data reported a much smaller value of  $AKIE_{Cl}^{alk.hydro} = 1.0009$   
13  
14  $\pm 0.0006$ <sup>36</sup> for the same alkaline hydrolysis at 50 °C. Later on it was confirmed that abiotic  
15  
16 alkaline hydrolysis performed earlier at 21 °C resembles rather neutral than alkaline conditions<sup>36</sup>.  
17  
18 Table 2 illustrates the different computed mechanisms that lie at the heart of the  
19  
20 computational predictions. It shows the different mechanistic routes between the alkaline  
21  
22 (substitution of Cl without protonation of the atrazine ring) and the acidic/enzymatic  
23  
24 pathway characterized in Meyer et al.<sup>9</sup> (substitution of Cl with protonation of the atrazine  
25  
26 ring) include different possible transition states. Chlorine KIEs are, among other factors<sup>45</sup>,  
27  
28 determined by the percent extension of the C-Cl bond in the transition state. As this is  
29  
30 directly related to the structure of the transition state, it can be linked to the C-Cl bond  
31  
32 orders (Table 2), which decrease in the studied hydrolysis reactions when the C-Cl bond  
33  
34 is more ruptured as compared to its length in the reactants, resulting in increased chlorine  
35  
36 KIEs. Previously performed computations<sup>36</sup> and computations of this study mimicking  
37  
38 alkaline, acidic, and neutral conditions indicated that the largest  $AKIE_{Cl}$  should be  
39  
40 expected under neutral conditions (except for transition state 2 of acidic/enzymatic  
41  
42 hydrolysis). Under neutral conditions the C-Cl bond is elongated leading to a transition  
43  
44 state geometry which differs substantially from hydrolysis reactions promoted either by  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 alkaline or acidic conditions (see Table 2). However, hydrolysis at neutral pH is too slow  
5  
6 to be of relevance. Computational calculations taking into account the transition state  
7  
8 structures at a molecular level predicted  $AKIE_{Cl}$  values ranging from 0.9996 to 1.0014 for  
9  
10 alkaline, acidic and enzymatic hydrolysis (see Table 1 and 2)<sup>36, 44</sup>. Hence, on the  
11  
12 mechanistic level, the computational studies predict that the formation of a Meisenheimer  
13  
14 complex rather than the subsequent C-Cl bond cleavage is the rate-determining step  
15  
16 during the nucleophilic aromatic substitution reaction catalyzed by TrzN<sup>36, 44</sup>. In both  
17  
18 abiotic pathways the C-Cl bond at the transition state of the rate determining step is  
19  
20 almost intact giving rise to very small  $AKIE_{Cl}$  (the computed bond orders for both alkaline  
21  
22 and acidic hydrolysis are the same and equal to 1.03, see also Table 2). In this study, we  
23  
24 therefore observed a similarly small  $AKIE^{Arthro}_{Cl}$  value for enzymatic hydrolysis in  
25  
26 *A. aurescens* TC1 which resembles acid-catalyzed hydrolysis rather than alkaline  
27  
28 hydrolysis<sup>9</sup>. Hence, the picture emerges that different hydrolytic pathways give rise to  
29  
30 experimental  $AKIE_{Cl}$  values much lower than the Streitwieser Limit of 1.02<sup>41-43</sup> indicating  
31  
32 that the chlorine isotope effect is masked in all cases and that the C-Cl bond cleavage is  
33  
34 not the rate-determining step. Interestingly, this is in contrast to ametryn hydrolysis where  
35  
36 strong sulphur isotope effects were observed in enzymatic hydrolysis by TrzN<sup>17</sup>. Further  
37  
38 experimental work, including degradation experiments with other strains, hydrolysis and  
39  
40 crude enzyme experiments, will be required to further substantiate the picture on chlorine  
41  
42 isotope effects observed in this study. For the moment, since chlorine isotope effects were  
43  
44 found to be masked, we must conclude, however, that information from chlorine isotope

1  
2  
3  
4 analysis alone would not be enough to differentiate the different reaction mechanisms.

5  
6 This illustrates the importance of analyzing more than one element for mechanistic  
7  
8 differentiation.  
9

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 2.** Mechanisms and transition states of acidic/enzymatic, neutral and alkaline hydrolysis and corresponding calculated and measured isotope effects.

Mechanism	Calculated Transition State <sup>a</sup>	Calculated Isotope Effect (position-specific and compound average AKIE values)	Measured Isotope Effect
<p><b>Acidic/Enzymatic Hydrolysis (Transition State 1)</b></p>	<p><b>C-Cl Bond Order: 1.03</b></p>	<p>Compound <b>average:</b> AKIE<sub>Cl</sub> = 1.0002<sup>a</sup></p> <p>AKIE<sub>Cl</sub> = 1.0000<sup>b</sup> 1.0037 1.0003 0.9995 1.0014 0.9973 1.0001 0.9996 1.0005 0.9902 1.0007 0.9999</p> <p>AKIE<sub>N</sub> = 0.9983<sup>a</sup> AKIE<sub>C</sub> = 1.0042<sup>a</sup></p>	<p>Compound <b>average:</b> AKIE<sub>Cl</sub> = 1.0014 ± 0.0006<sup>b</sup> AKIE<sub>N</sub> = 0.9886 ± 0.0015<sup>b</sup> AKIE<sub>C</sub> = 1.0271 ± 0.0034<sup>b</sup></p>
<p><b>Acidic/Enzymatic Hydrolysis (Transition State 2)</b></p>	<p><b>C-Cl Bond Order: 0.55</b></p>	<p>Compound <b>average:</b> AKIE<sub>Cl</sub> = 1.0099</p> <p>AKIE<sub>Cl</sub> = 1.0016<sup>b</sup> 1.0013 1.0015 0.9999 1.0005 1.0005 1.0007 1.000 0.9999 0.9995 0.9992 0.9993</p> <p>AKIE<sub>N</sub> = 1.0002 AKIE<sub>C</sub> = 1.0017</p>	-
<p><b>Neutral Hydrolysis</b></p>	<p><b>C-Cl Bond Order: 0.87</b></p>	<p>Compound <b>average:</b> AKIE<sub>Cl</sub> = 1.0045</p> <p>AKIE<sub>Cl</sub> = 1.0046<sup>b</sup> 1.0362 1.0041 0.9992 1.004 0.9984 0.9993 1.000 0.9996 0.9990 1.0013 1.0014 0.9999 1.0011 1.0014</p> <p>AKIE<sub>N</sub> = 1.0024 AKIE<sub>C</sub> = 1.0041</p>	-



1  
2  
3  
4 For oxidative dealkylation, so far, no chlorine isotope effects were reported. Regarding  
5  
6 the reaction mechanism, Meyer et al.<sup>16</sup> concluded that oxidative dealkylation of atrazine  
7  
8 with *Rhodococcus* sp. NI86/21 is initiated by hydrogen atom transfer based on the  
9  
10 observed product distribution and the carbon and nitrogen isotope effects. Hydrogen atom  
11  
12 transfer leads directly to a homolytic cleavage of the C-H bond adjacent to the nitrogen  
13  
14 atom ( $\alpha$ -position of the ethyl or isopropyl group) producing a relative unstable 1,1-  
15  
16 aminoalcohol which is then further transformed to DEA or DIA<sup>16</sup>. In parallel, two additional  
17  
18 products could be detected which were formed by oxidation of the C-H bond in the  $\beta$ -  
19  
20 position of the ethyl or isopropyl group. For this mechanistic pathway, chlorine isotope  
21  
22 effects would be expected to be rather small since the chlorine is not involved in the  
23  
24 reaction steps. The closed mass balance of the concentration analysis (see Figure S5 in  
25  
26 the SI) of this study and the results of product distribution of Meyer et al.<sup>16</sup> also indicate  
27  
28 that there is no C-Cl bond cleavage taking place since corresponding hydrolysis products  
29  
30 were not detected. Furthermore, our computations for hydrogen atom transfer at a  
31  
32 catalytic center mimicking cytochrome P450 predicted no chlorine isotope effect  
33  
34 (AKIE<sup>hydro.atom trans.</sup><sub>Cl</sub> = 0.9999, see Table 1). Hydride transfer promoted by the hydronium  
35  
36 ion also resulted in no chlorine isotope effect (AKIE<sup>hydride trans.</sup><sub>Cl</sub> = 0.9997, see Table 1). At  
37  
38 previously located transition state structures for these two reactions<sup>16</sup> the carbon-chlorine  
39  
40 bond remains intact and no stretching of this bond is involved in the reaction coordinate  
41  
42 (hydrogen transfer) mode. The observed more pronounced AKIE<sup>Rhodo</sup><sub>Cl</sub> value of 1.0043  
43  
44  $\pm 0.0018$  in this study (see Table 1) could, therefore, be indicative of isotope effects  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 caused by enzymatic interactions. Meyer et al.<sup>16</sup> proposed that for oxidative dealkylation  
5  
6 no selectivity itself is observed, however, the preferred oxidation of the  $\alpha$ -position over  
7  
8 the  $\beta$ -position could be explained by steric factors of the catalyzing enzyme which could  
9  
10 have an influence on the transformation pathway. Thus, the sensitive chlorine isotope  
11  
12 effect, which is observed even though the C-Cl bond is not cleaved during degradation,  
13  
14 can be interpreted as an indicator that non-covalent interactions between the  
15  
16 cytochrome P450 complex and the chlorine cause significant chlorine isotope  
17  
18 fractionation<sup>46</sup>.

## 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

### CONCLUSION

Since atrazine is frequently detected in groundwater systems, major efforts should be put into understanding its environmental fate. We provide an approach to 3D-isotope (C, N, Cl) analysis of atrazine and explored isotope fractionation in different transformation pathways. Together, this provides the basis to more confidently assess sources and degradation of atrazine in the environment. Specifically, we demonstrated that pronounced changes in chlorine isotope values are not an indicator of microbial hydrolysis (as one might have expected without knowledge of our experimental data), but – surprisingly – rather of oxidative dealkylation. Therefore, although trends are different than expected, they can nonetheless be used for a more confident identification of different sources and transformation pathways in field samples. Regarding the sensitivity of chlorine isotope effects, our study demonstrates the importance of performing

1  
2  
3 controlled laboratory experiments before applying the approach in the field. Specifically,  
4  
5 in other cases chlorine isotope fractionation can be much more pronounced than  
6  
7 observed for atrazine in this study. Large chlorine isotope effects were observed in proof-  
8  
9 of-principle experiments by Ponsin et al.<sup>20</sup> studying hydrolytic dechlorination of S-  
10  
11 metolachlor, an herbicide containing also only one chlorine atom. Here preliminary data  
12  
13 suggest a large chlorine isotope effect of  $\epsilon_{\text{Cl}} = -9.7 \pm 2.9 \text{ ‰}$  for abiotic alkaline hydrolysis.  
14  
15 Therefore, in the case of other substances chlorine isotope effects can be even more  
16  
17 sensitive indicators of degradation provided that the C-Cl bond cleavage occurs in the  
18  
19 rate-determining step of a reaction. Further, gaining deeper insights into these chemical  
20  
21 processes is the basis for understanding the biotic catalysis of organic micropollutant  
22  
23 degradation. This, in turn, is essential for identifying and developing optimized strategies  
24  
25 for micropollutant removal in order to make successful bioremediation possible.  
26  
27

## 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

### CONFLICT OF INTEREST

There are no conflicts to declare.

### ACKNOWLEDGMENT

This work was supported by the Swiss National Science Foundation (SNSF, Grant CRSII2\_141805), the German Israeli Foundation for Scientific Research and Development (GIF, Grant I-251-307.4-2013) and the National Science Center in Poland (Sonata BIS grant UMO-2014/14/E/ST4/00041). We thank PLGrid Infrastructure

(Poland) for computer resources and Armin Meyer for his advice regarding the microbial degradation experiments.

## REFERENCES

1. H. Gysin and E. Knusli, Chemistry and herbicidal properties of triazine derivatives, *Adv. in Pest Control Res.*, 1960, **3**, 289-353.
2. EPA, Pesticides Industry Sales and Usage: 2008–2012 Market Estimates, *US Environmental Protection Agency, Washington (DC)*, 2017.
3. European Commission, Commission decision of 10 March 2004 concerning the non-inclusion of atrazine in Annex I to Council Directive 91/414/EEC and the withdrawal of authorisations for plant protection products containing this active substance., *Official Journal of the European Union*, 2004, **78**, 53-55.
4. R. Loos, G. Locoro, S. Comero, S. Contini, D. Schwesig, F. Werres, P. Balsaa, O. Gans, S. Weiss, L. Blaha, M. Bolchi and B. M. Gawlik, Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water, *Water Res.*, 2010, **44**, 4115-4126.
5. D. Vonberg, J. Vanderborght, N. Cremer, T. Pütz, M. Herbst and H. Vereecken, 20 years of long-term atrazine monitoring in a shallow aquifer in western Germany, *Water Res.*, 2014, **50**, 294-306.
6. S. Singh, V. Kumar, A. Chauhan, S. Datta, A. B. Wani, N. Singh and J. Singh, Toxicity, degradation and analysis of the herbicide atrazine, *Environ. Chem. Lett.*, 2018, **16**, 211-237.
7. S. Kern, H. P. Singer, J. Hollender, R. P. Schwarzenbach and K. Fenner, Assessing exposure to transformation products of soil-applied organic contaminants in surface water: comparison of model predictions and field data, *Environ. Sci. Technol.*, 2011, **45**, 2833-2841.
8. C. Moreau-Kervevan and C. Mouvet, Adsorption and desorption of atrazine, deethylatrazine, and hydroxyatrazine by soil components, *J. Environ. Qual.*, 1998, **27**, 46-53.

9. A. H. Meyer, H. Penning and M. Elsner, C and N isotope fractionation suggests similar mechanisms of microbial atrazine transformation despite involvement of different Enzymes (AtzA and TrzN), *Environ. Sci. Technol.*, 2009, **43**, 8079-8085.
10. A. H. Meyer and M. Elsner,  $^{13}\text{C}/^{12}\text{C}$  and  $^{15}\text{N}/^{14}\text{N}$  Isotope Analysis To Characterize Degradation of Atrazine: Evidence from Parent and Daughter Compound Values, *Environ. Sci. Technol.*, 2013, **47**, 6884-6891.
11. A. H. Meyer, H. Penning, H. Lowag and M. Elsner, Precise and accurate compound specific carbon and nitrogen isotope analysis of atrazine: critical role of combustion oven conditions, *Environ. Sci. Technol.*, 2008, **42**, 7757-7763.
12. S. Reinnicke, D. Juchelka, S. Steinbeiss, A. H. Meyer, A. Hilker and M. Elsner, Gas chromatography-isotope ratio mass spectrometry (GC-IRMS) of recalcitrant target compounds: performance of different combustion reactors and strategies for standardization, *Rapid. Commun. Mass. Sp.*, 2012, **26**, 1053-1060.
13. K. Schreglmann, M. Hoeche, S. Steinbeiss, S. Reinnicke and M. Elsner, Carbon and nitrogen isotope analysis of atrazine and desethylatrazine at sub-microgram per liter concentrations in groundwater, *Anal. Bioanal. Chem.*, 2013, **405**, 2857-2867.
14. T. C. Schmidt, L. Zwank, M. Elsner, M. Berg, R. U. Meckenstock and S. B. Haderlein, Compound-specific stable isotope analysis of organic contaminants in natural environments: a critical review of the state of the art, prospects, and future challenges, *Anal. Bioanal. Chem.*, 2004, **378**, 283-300.
15. M. Elsner, Stable isotope fractionation to investigate natural transformation mechanisms of organic contaminants: principles, prospects and limitations, *J. Environ. Monit.*, 2010, **12**, 2005-2031.
16. A. H. Meyer, A. Dybala-Defratyka, P. J. Alaimo, I. Geronimo, A. D. Sanchez, C. J. Cramer and M. Elsner, Cytochrome P450-catalyzed dealkylation of atrazine by *Rhodococcus* sp. strain NI86/21 involves hydrogen atom transfer rather than single electron transfer, *Dalton Trans.*, 2014, **43**, 12111-12432.
17. H. K. V. Schürner, J. L. Seffernick, A. Grzybkowska, A. Dybala-Defratyka, L. P. Wackett and M. Elsner, Characteristic Isotope Fractionation Patterns in s-

- 1  
2  
3  
4 Triazine Degradation Have Their Origin in Multiple Protonation Options in the s-  
5 Triazine Hydrolase TrzN, *Environ. Sci. Technol.*, 2015, **49**, 3490-3498.
- 6  
7 18. T. B. Hofstetter and M. Berg, Assessing transformation processes of organic  
8 contaminants by compound-specific stable isotope analysis, *TrAC, Trends Anal.*  
9 *Chem.*, 2011, **30**, 618-627.
- 10  
11 19. J. Renpenning, A. Horst, M. Schmidt and M. Gehre, Online isotope analysis of  
12 <sup>37</sup>Cl/<sup>35</sup>Cl universally applied for semi-volatile organic compounds using GC-MC-  
13 ICPMS, *J. Anal. At. Spectrom.*, 2018, **33**, 314-321.
- 14  
15 20. V. Ponsin, C. Torrentó, C. Lihl, M. Elsner and D. Hunkeler, Compound-specific  
16 chlorine isotope analysis of the herbicides atrazine, acetochlor and metolachlor,  
17 *Anal. Chem.*, 2019, **91**, 14290-14298.
- 18  
19 21. C. Lihl, L. M. Douglas, S. Franke, A. Pérez-de-Mora, A. H. Meyer, M. Daubmeier,  
20 E. A. Edwards, I. Nijenhuis, B. Sherwood Lollar and M. Elsner, Mechanistic  
21 Dichotomy in Bacterial Trichloroethene Dechlorination Revealed by Carbon and  
22 Chlorine Isotope Effects, *Environ. Sci. Technol.*, 2019, **53**, 4245-4254.
- 23  
24 22. L. E. Erickson, Degradation of atrazine and related s-triazines, *Crit. Rev. Env.*  
25 *Con.*, 1989, **19**, 1-14.
- 26  
27 23. H. M. LeBaron, J. E. McFarland and O. C. Burnside, *The triazine herbicides*,  
28 Elsevier, Oxford, 1 edn., 2008.
- 29  
30 24. L. C. Strong, C. Rosendahl, G. Johnson, M. J. Sadowsky and L. P. Wackett,  
31 *Arthrobacter aurescens* TC1 metabolizes diverse s-triazine ring compounds,  
32 *Appl. Environ. Microbiol.*, 2002, **68**, 5973-5980.
- 33  
34 25. K. Sajjaphan, N. Shapir, L. P. Wackett, M. Palmer, B. Blackmon, J. Tomkins and  
35 M. J. Sadowsky, *Arthrobacter aurescens* TC1 Atrazine Catabolism Genes *trzN*,  
36 *atzB*, and *atzC* Are Linked on a 160-Kilobase Region and Are Functional in  
37 *Escherichia coli*, *Appl. Environ. Microbiol.*, 2004, **70**, 4402-4407.
- 38  
39 26. I. Nagy, F. Compennolle, K. Ghys, J. Vanderleyden and R. De Mot, A single  
40 cytochrome P-450 system is involved in degradation of the herbicides EPTC (S-  
41 ethyl dipropylthiocarbamate) and atrazine by *Rhodococcus* sp. strain NI86/21,  
42 *Appl. Environ. Microbiol.*, 1995, **61**, 2056-2060.
- 43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 27. M. Elsner, L. Zwank, D. Hunkeler and R. P. Schwarzenbach, A new concept  
5 linking observable stable isotope fractionation to transformation pathways of  
6 organic pollutants, *Environ. Sci. Technol.*, 2005, **39**, 6896-6916.
- 7  
8  
9 28. L. Zwank, M. Berg, M. Elsner, T. C. Schmidt, R. P. Schwarzenbach and S. B.  
10 Haderlein, New evaluation scheme for two-dimensional isotope analysis to  
11 decipher biodegradation processes: application to groundwater contamination by  
12 MTBE, *Environ. Sci. Technol.*, 2005, **39**, 1018-1029.
- 13  
14  
15 29. S. Reinnicke, A. Simonsen, S. R. Sørensen, J. Aamand and M. Elsner, C and N  
16 Isotope Fractionation during Biodegradation of the Pesticide Metabolite 2,6-  
17 Dichlorobenzamide (BAM): Potential for Environmental Assessments, *Environ.*  
18 *Sci. Technol.*, 2012, **46**, 1447-1454.
- 19  
20  
21 30. T. Gilevska, M. Gehre and H. H. Richnow, Multidimensional isotope analysis of  
22 carbon, hydrogen and oxygen as tool for identification of the origin of ibuprofen,  
23 *J. Pharm. Biomed. Anal.*, 2015, **115**, 410-417.
- 24  
25  
26 31. J. Palau, O. Shouakar-Stash, S. Hatijah Mortan, R. Yu, M. Rosell, E. Marco-  
27 Urrea, D. L. Freedman, R. Aravena, A. Soler and D. Hunkeler, Hydrogen Isotope  
28 Fractionation during the Biodegradation of 1,2-Dichloroethane: Potential for  
29 Pathway Identification Using a Multi-element (C, Cl, and H) Isotope Approach,  
30 *Environ. Sci. Technol.*, 2017, **51**, 10526-10535.
- 31  
32  
33 32. S. Franke, C. Lihl, J. Renpenning, M. Elsner and I. Nijenhuis, Triple-element  
34 compound-specific stable isotope analysis of 1,2-dichloroethane for  
35 characterization of the underlying dehalogenation reaction in two  
36 *Dehalococcoides mccartyi* strains, *FEMS Microbiol. Ecol.*, 2017, **93**, fix137.
- 37  
38  
39 33. T. Kuder, B. M. van Breukelen, M. Vanderford and P. Philp, 3D-CSIA: Carbon,  
40 Chlorine, and Hydrogen Isotope Fractionation in Transformation of TCE to  
41 Ethene by a *Dehalococcoides* Culture, *Environ. Sci. Technol.*, 2013, **47**, 9668-  
42 9677.
- 43  
44  
45 34. B. Heckel, K. McNeill and M. Elsner, Chlorinated Ethene Reactivity with Vitamin  
46 B12 Is Governed by Cobalamin Chloroethylcarbanions as Crossroads of  
47 Competing Pathways, *ACS Catal.*, 2018, **8**, 3054-3066.
- 48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 35. A. Dybala-Defratyka, L. Szatkowski, R. Kaminski, M. Wujec, A. Siwek and P.  
5 Paneth, Kinetic isotope effects on dehalogenations at an aromatic carbon,  
6 *Environ. Sci. Technol.*, 2008, **42**, 7744-7750.
- 7  
8  
9 36. A. Grzybkowska, R. Kaminski and A. Dybala-Defratyka, Theoretical predictions  
10 of isotope effects versus their experimental values for an example of uncatalyzed  
11 hydrolysis of atrazine, *Phys. Chem. Chem. Phys.*, 2014, **16**, 15164-15172.
- 12  
13  
14 37. C. Lihl, J. Renpenning, S. Kümmel, F. Gelman, H. K. V. Schürner, M. Daubmeier,  
15 B. Heckel, A. Melsbach, A. Bernstein, O. Shouakar-Stash, M. Gehre and M.  
16 Elsner, Toward Improved Accuracy in Chlorine Isotope Analysis: Synthesis  
17 Routes for In-House Standards and Characterization via Complementary Mass  
18 Spectrometry Methods, *Anal. Chem.*, 2019, **91**, 12290-12297.
- 19  
20  
21 38. D. Hunkeler, R. U. Meckenstock, B. Sherwood Lollar, T. Schmidt, J. Wilson, T.  
22 Schmidt and J. Wilson, *A Guide for Assessing Biodegradation and Source*  
23 *Identification of Organic Ground Water Contaminants using Compound Specific*  
24 *Isotope Analysis (CSIA)*, O. o. R. a. Development Report PA 600/R-08/148 |  
25 December 2008 | [www.epa.gov/ada](http://www.epa.gov/ada), US EPA, Oklahoma, USA, 2008.
- 26  
27  
28 39. J. Bigeleisen, The relative reaction velocities of isotopic molecules, *J. Chem.*  
29 *Phys.*, 1949, **17**, 675-679.
- 30  
31  
32 40. V. Anisimov and P. Paneth, ISOEFF98. A program for studies of isotope effects  
33 using Hessian modifications, *J. Math. Chem.*, 1999, **26**.
- 34  
35  
36 41. A. Streitwieser Jr, R. Jagow, R. Fahey and S. Suzuki, Kinetic isotope effects in  
37 the acetolyses of deuterated cyclopentyl tosylates<sup>1, 2</sup>, *J. Am. Chem. Soc.*, 1958,  
38 **80**, 2326-2332.
- 39  
40  
41 42. K. Świderek and P. Paneth, Extending limits of chlorine kinetic isotope effects, *J.*  
42 *Org. Chem.*, 2012, **77**, 5120-5124.
- 43  
44  
45 43. L. Szatkowski and A. Dybala-Defratyka, A computational study on enzymatically  
46 driven oxidative coupling of chlorophenols: An indirect dehalogenation reaction,  
47 *Chemosphere*, 2013, **91**, 258-264.
- 48  
49  
50 44. L. Szatkowski, R. N. Manna, A. Grzybkowska, R. Kamiński, A. Dybala-Defratyka  
51 and P. Paneth, in *Methods in enzymology*, Elsevier, 2017, vol. 596, pp. 179-215.
- 52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 45. A. Dybala-Defratyka, M. Rostkowski, O. Matsson, K. C. Westaway and P.  
5 Paneth, A New Interpretation of Chlorine Leaving Group Kinetic Isotope Effects;  
6 A Theoretical Approach, *J. Org. Chem.*, 2004, **69**, 4900-4905.  
7  
8  
9 46. A. Siwek, R. Omi, K. Hirotsu, K. Jitsumori, N. Esaki, T. Kurihara and P. Paneth,  
10 Binding modes of DL-2-haloacid dehalogenase revealed by crystallography,  
11 modeling and isotope effects studies, *Arch. Biochem. Biophys.*, 2013, **540**, 26-  
12 32.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Graphical Abstract:

