

OCCUPATION TIMES OF SPECTRALLY NEGATIVE LÉVY PROCESSES WITH APPLICATIONS

DAVID LANDRIAULT, JEAN-FRANÇOIS RENAUD, AND XIAOWEN ZHOU

ABSTRACT. In this paper, we compute the Laplace transform of occupation times (of the negative half-line) of spectrally negative Lévy processes. Our results are extensions of known results for standard Brownian motion and jump-diffusion processes. The results are expressed in terms of the so-called scale functions of the spectrally negative Lévy process and its Laplace exponent. Applications to insurance risk models are also presented.

1. INTRODUCTION

Let $X = (X_t)_{t \geq 0}$ be a spectrally negative Lévy process, that is a Lévy process with no positive jumps. The law of X such that $X_0 = x$ is denoted by \mathbb{P}_x and the corresponding expectation by \mathbb{E}_x . We write \mathbb{P} and \mathbb{E} when $x = 0$. As the Lévy process X has no positive jumps, its Laplace transform exists and is given by

$$\mathbb{E} \left[e^{\theta X_t} \right] = e^{t\psi(\theta)},$$

for $\theta, t \geq 0$, where

$$\psi(\theta) = \gamma\theta + \frac{1}{2}\sigma^2\theta^2 + \int_{-\infty}^0 \left(e^{\theta z} - 1 - \theta z \mathbb{I}_{(-1,0)}(z) \right) \Pi(dz),$$

for $\gamma \in \mathbb{R}$ and $\sigma \geq 0$. Also, Π is a σ -finite measure on $(-\infty, 0)$ such that

$$\int_{-\infty}^0 (1 \wedge z^2) \Pi(dz) < \infty.$$

The measure Π is called the Lévy measure of X , while (γ, σ, Π) is referred to as the Lévy triplet of X . Note that $\mathbb{E}[X_1] = \psi'(0+)$.

For an arbitrary spectrally negative Lévy process, the Laplace exponent ψ is strictly convex and $\lim_{\theta \rightarrow \infty} \psi(\theta) = \infty$. Thus, there exists a function $\Phi: [0, \infty) \rightarrow [0, \infty)$ defined by $\Phi(\theta) = \sup\{\xi \geq 0 \mid \psi(\xi) = \theta\}$ (its right-inverse) and such that

$$\psi(\Phi(\theta)) = \theta, \quad \theta \geq 0.$$

We first examine the total occupation time of the negative half-line $(-\infty, 0)$.

Theorem 1. *If $\psi'(0+) > 0$, then, for $\lambda \geq 0$,*

$$(1) \quad \mathbb{E} \left[e^{-\lambda \int_0^\infty \mathbb{I}_{\{X_s \leq 0\}} ds} \right] = \psi'(0+) \frac{\Phi(\lambda)}{\lambda},$$

Date: January 7, 2011.

Key words and phrases. Occupation time, spectrally negative Lévy processes, fluctuation theory, scale functions, ruin theory.

where $\Phi(\lambda)/\lambda$ is to be understood in the limiting sense when $\lambda = 0$.

This result generalizes the work done in [20] where the sum of a compound Poisson process and a Brownian motion is analyzed (see, e.g., Equation (4.9) in that paper). Using ruin theory terminology, they study the duration of negative surplus in such an insurance risk model; their work is itself an extension of [8] in the pure compound Poisson case. The advantage of (1) is to involve only the Laplace exponent ψ of X and its right-inverse Φ , yielding a very explicit expression.

We now recall the definition of the q -scale function $W^{(q)}$. For $q \geq 0$, the q -scale function of the process X is defined as the function with Laplace transform on $[0, \infty)$ given by

$$\int_0^\infty e^{-\theta z} W^{(q)}(z) dz = \frac{1}{\psi(\theta) - q}, \quad \text{for } \theta > \Phi(q),$$

and such that $W^{(q)}(x) = 0$ for $x < 0$. This function is unique, continuous, positive and strictly increasing. We write $W = W^{(0)}$ when $q = 0$. We have that $W^{(q)}$ is differentiable except for at most countably many points; see [17]. Moreover, $W^{(q)}$ is continuously differentiable if X has paths of unbounded variation or if the tail of the Lévy measure, i.e., the function $x \mapsto \Pi(-\infty, -x)$ on $(-\infty, 0)$, is continuous. Further, $W^{(q)}$ is twice continuously differentiable on $(0, \infty)$ if $\sigma > 0$. For more details on the smoothness properties of the q -scale function, see [3]. We will also use the functions $\{\overline{W}^{(q)}; q \geq 0\}$ and $\{Z^{(q)}; q \geq 0\}$ defined by

$$\overline{W}^{(q)}(x) = \int_0^x W^{(q)}(z) dz.$$

and

$$Z^{(q)}(x) = 1 + q \overline{W}^{(q)}(x).$$

We now examine the occupation time of $(-\infty, 0)$ until a negative level $-b$ is crossed for the first time. Let τ_{-b}^- be the first passage time below $-b$ of X :

$$\tau_{-b}^- = \inf\{t > 0: X_t < -b\}.$$

Theorem 2. *If $\psi'(0+) \geq 0$, then, for $\lambda \geq 0$,*

$$(2) \quad \mathbb{E} \left[e^{-\lambda \int_0^{\tau_{-b}^-} \mathbb{1}_{\{X_s \leq 0\}} ds} \right] = \frac{\psi'(0+) + \frac{\sigma^2}{2} \frac{A_1^{(\lambda)}(b)}{W^{(\lambda)}(b)} + \int_{-b}^{0-} A_2^{(\lambda)}(x) \int_{0+}^\infty \Pi(dx - y) dy}{\psi'(0+) + \frac{\sigma^2}{2} \frac{W^{(\lambda)'(b)}}{W^{(\lambda)}(b)} + \int_{-\infty}^{0-} A_3^{(\lambda)}(x) \int_{0+}^\infty \Pi(dx - y) dy},$$

where

$$A_1^{(\lambda)}(b) = Z^{(\lambda)}(b) W^{(\lambda)'(b)} - \lambda (W^{(\lambda)}(b))^2,$$

$$A_2^{(\lambda)}(x) = Z^{(\lambda)}(x + b) - Z^{(\lambda)}(b) \frac{W^{(\lambda)}(x + b)}{W^{(\lambda)}(b)},$$

and

$$A_3^{(\lambda)}(x) = 1 - \frac{W^{(\lambda)}(x + b)}{W^{(\lambda)}(b)}.$$

As a special case, we recover the corresponding result for standard Brownian motion; see, e.g., [11, 12].

Our proofs use fluctuation identities for spectrally negative Lévy processes and, as a consequence, our results are expressed in terms of the so-called scale functions of the spectrally negative Lévy process (see, e.g., [1, 2]) and its Laplace exponent. Only elementary arguments are needed. To the authors' knowledge, the literature seems rather scarce on the relationship between scale functions and certain occupation times of a general spectrally negative Lévy process.

The current work has been partly motivated by the study of an insurance risk model with implementation delays. Insurance risk models use stochastic processes to describe the surplus of an insurance company. In risk models of a Parisian nature, an implementation delay in the recognition of an insurer's capital insufficiency is applied. More precisely, it is assumed that ruin occurs as soon as an excursion below a critical level is longer than a deterministic time; such models have been studied only very recently in [5, 6, 16] and the idea has been borrowed from finance and Parisian barrier options (see [4]). Of more interest in our context is the work by the same authors: in [16], instead of a deterministic delay, an exponentially distributed grace period is used in the definition of the Parisian ruin. It turns out that the probability of ruin in this model is strongly related to the occupation time of the underlying process. The reader is invited to consult Section 5.2 to obtain further details on this connection.

The rest of the paper is organized as follows. In the next Section, we recall the relevant notions and results on scale functions and fluctuation identities. Then, in Section 3 and 4, we prove Theorem 2 and Theorem 1 respectively. Section 5 presents applications to the case of a Brownian motion with drift and insurance risk models.

2. SCALE FUNCTIONS AND FLUCTUATION IDENTITIES

We recall some of the properties of the q -scale function $W^{(q)}$ and its use in fluctuation theory. Let $d = \gamma - \int_{-1}^0 z\Pi(dz)$. The initial values of $W^{(q)}$ and $W^{(q) \prime}$ are known to be

$$W^{(q)}(0+) = \begin{cases} 1/d & \text{when } \sigma = 0 \text{ and } \int_{-1}^0 z\Pi(dz) < \infty, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$W^{(q) \prime}(0+) = \begin{cases} 2/\sigma^2 & \text{when } \sigma > 0, \\ (\Pi(-\infty, 0) + q)/d^2 & \text{when } \sigma = 0 \text{ and } \Pi(-\infty, 0) < \infty, \\ \infty & \text{otherwise.} \end{cases}$$

Now, define

$$\tau_0^- = \inf\{t > 0: X_t < 0\},$$

and, for $a > 0$,

$$\tau_a^+ = \inf\{t > 0: X_t > a\},$$

with the convention $\inf \emptyset = \infty$. It is well known (see, e.g., [14]) that, for $x \leq a$,

$$\mathbb{E}_x \left[e^{-q\tau_a^+}; \tau_a^+ < \tau_0^- \right] = \frac{W^{(q)}(x)}{W^{(q)}(a)},$$

$$\mathbb{E}_x \left[e^{-q\tau_0^-}; \tau_0^- < \tau_a^+ \right] = Z^{(q)}(x) - Z^{(q)}(a) \frac{W^{(q)}(x)}{W^{(q)}(a)},$$

and

$$\mathbb{E}_x \left[e^{-q\tau_a^+}; \tau_a^+ < \infty \right] = e^{-\Phi(q)(a-x)}.$$

If $\psi'(0+) \geq 0$, then $\mathbb{P}_x \{ \tau_a^+ < \infty \} = 1$ and therefore

$$\mathbb{E}_x \left[e^{-q\tau_a^+} \right] = e^{-\Phi(q)(a-x)}.$$

Also, we have that

$$\begin{aligned} \mathbb{P}_x \left\{ \tau_0^- < \infty, X_{\tau_0^-} \in dz \right\} &= \frac{\sigma^2}{2} [W'(x) - \Phi(0)W(x)] \delta_0(dz) \\ &\quad + \int_{0+}^{\infty} \Pi(dz - y) \left\{ e^{-\Phi(0)y} W(x) - W(x - y) \right\} dy, \end{aligned}$$

where δ_0 is the Dirac measure at 0. The first term of this measure corresponds to the case when $X_{\tau_0^-} = 0$, a behaviour called *creeping*. Using the above equation, together with

$$\mathbb{P}_x \{ \tau_0^- < \infty \} = \begin{cases} 1 - \psi'(0+)W(x) & \text{if } \psi'(0+) \geq 0, \\ 1 & \text{if } \psi'(0+) < 0, \end{cases}$$

if $\psi'(0+) \geq 0$ (in which case $\Phi(0) = 0$), we have

$$1 = \psi'(0+)W(x) + \frac{\sigma^2}{2} W'(x) + \int_{-\infty}^{0-} \int_{0+}^{\infty} \Pi(dz - y) \left\{ e^{-\Phi(0)y} W(x) - W(x - y) \right\} dy.$$

For more details on spectrally negative Lévy processes and fluctuation identities, the reader is referred to [14]. Further information, examples and numerical techniques related to the computation of scale functions can be found in [3, 9, 10, 15, 19].

3. PROOF OF THEOREM 2

The main idea of the proof consists in defining a quantity underestimating and overestimating the occupation time

$$\int_0^{\tau_0^-} \mathbb{I}_{\{X_s \leq 0\}} ds.$$

This respectively leads to an upper and lower bound to its Laplace transform. Subsequently, by taking an appropriate limit, we show that the two bounds converge to the expression on the right-hand side of 2.

First, we provide a lower bound to this Laplace transform by overestimating the occupation time. To this end, we consider a clock which starts at time 0 and stops when level ϵ is attained or when level $-b$ is crossed. Then, if level ϵ was attained first, every time we go below 0, we re-start the clock and subsequently stop it when we get back to ϵ (without going below $-b$); let $L^{\epsilon,b}$ be the Laplace transform of this overestimating quantity of the occupation time when the process X sits at level ϵ at time 0. Hence, by the strong Markov property of X , we have

$$\mathbb{E} \left[e^{-\lambda \int_0^{\tau_0^-} \mathbb{I}_{\{X_s \leq 0\}} ds} \right] \geq \mathbb{E} \left[e^{-\lambda \tau_\epsilon^+}; \tau_\epsilon^+ < \tau_{-b}^- \right] L^{\epsilon,b} = \frac{W^{(\lambda)}(b)}{W^{(\lambda)}(b + \epsilon)} L^{\epsilon,b}.$$

Using the strong Markov property and the spatial homogeneity of X , together with the results in Section 2, we get

$$L^{\epsilon,b} = \mathbb{P}_\epsilon\{\tau_0^- = \infty\} + \int_{-b}^{0+} \mathbb{P}_\epsilon\left\{\tau_0^- < \infty, X_{\tau_0^-} \in dx\right\} \left\{\mathbb{E}_x\left[e^{-\lambda\tau_{-b}^-}; \tau_{-b}^- < \tau_\epsilon^+\right] + \mathbb{E}_x\left[e^{-\lambda\tau_\epsilon^+}; \tau_\epsilon^+ < \tau_{-b}^-\right] L^{\epsilon,b}\right\}.$$

Consequently, using some results from Section 2,

$$L^{\epsilon,b} = \frac{\mathbb{P}_\epsilon\{\tau_0^- = \infty\} + \int_{-b}^{0+} \mathbb{P}_\epsilon\left\{\tau_0^- < \infty, X_{\tau_0^-} \in dx\right\} \mathbb{E}_x\left[e^{-\lambda\tau_{-b}^-}; \tau_{-b}^- < \tau_\epsilon^+\right]}{1 - \int_{-b}^{0+} \mathbb{P}_\epsilon\left\{\tau_0^- < \infty, X_{\tau_0^-} \in dx\right\} \mathbb{E}_x\left[e^{-\lambda\tau_\epsilon^+}; \tau_\epsilon^+ < \tau_{-b}^-\right]} = \frac{L_1^{\epsilon,b} + L_2^{\epsilon,b}}{L_3^{\epsilon,b}},$$

where

$$L_1^{\epsilon,b} = \psi'(0+)W(\epsilon) + \frac{\sigma^2}{2}W'(\epsilon) \left[Z^{(\lambda)}(b) - Z^{(\lambda)}(b+\epsilon) \frac{W^{(\lambda)}(b)}{W^{(\lambda)}(b+\epsilon)} \right],$$

$$L_2^{\epsilon,b} = \int_{-b}^{0-} \left[Z^{(\lambda)}(x+b) - Z^{(\lambda)}(b+\epsilon) \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b+\epsilon)} \right] \int_{0+}^{\infty} \Pi(dx-y) \{W(\epsilon) - W(\epsilon-y)\} dy,$$

and,

$$L_3^{\epsilon,b} = 1 - \frac{\sigma^2}{2}W'(\epsilon) \frac{W^{(\lambda)}(b)}{W^{(\lambda)}(b+\epsilon)} - \int_{-b}^{0-} \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(x+b+\epsilon)} \int_{0+}^{\infty} \Pi(dx-y) \{W(\epsilon) - W(\epsilon-y)\} dy.$$

Alternatively, we now develop a scheme to underestimate the occupation time in question. Every time we go below $-\epsilon$ ($-\epsilon > -b$), we start the clock and stop it when we get back to 0 (without going below $-b$); let $U^{\epsilon,b}$ be the Laplace transform of this underestimating quantity of the occupation time when $X_0 = 0$. Hence, by the strong Markov property of X , we also have

$$\mathbb{E} \left[e^{-\lambda \int_0^{\tau_{-b+\epsilon}^-} \mathbb{1}_{\{X_s \leq 0\}} ds} \right] \leq U^{\epsilon,b}.$$

As above, using the strong Markov property and the spatial homogeneity of X , together with some of the results in Section 2, we can write

$$U^{\epsilon,b} = \int_{-b+\epsilon}^{0+} \mathbb{P}_\epsilon\left\{\tau_0^- < \infty, X_{\tau_0^-} \in dx\right\} \left\{\mathbb{E}_x\left[e^{-\lambda\tau_{-b+\epsilon}^-}; \tau_{-b+\epsilon}^- < \tau_\epsilon^+\right] + \mathbb{E}_x\left[e^{-\lambda\tau_\epsilon^+}; \tau_\epsilon^+ < \tau_{-b+\epsilon}^-\right] U^{\epsilon,b}\right\}$$

and then

$$U^{\epsilon,b} = \frac{U_1^{\epsilon,b} + U_2^{\epsilon,b}}{U_3^{\epsilon,b}},$$

where

$$U_1^{\epsilon,b} = \frac{\sigma^2}{2}W'(\epsilon) \left[Z^{(\lambda)}(b-\epsilon) - Z^{(\lambda)}(b) \frac{W^{(\lambda)}(b-\epsilon)}{W^{(\lambda)}(b)} \right],$$

$$U_2^{\epsilon,b} = \int_{-b}^{0-} \left[Z^{(\lambda)}(x+b-\epsilon) - Z^{(\lambda)}(b) \frac{W^{(\lambda)}(x+b-\epsilon)}{W^{(\lambda)}(b)} \right] \mathbb{I}_{(-b+\epsilon,0)}(x) \\ \times \int_{0+}^{\infty} \Pi(dx-y) \{W(\epsilon) - W(\epsilon-y)\} dy,$$

and,

$$U_3^{\epsilon,b} = 1 - \frac{\sigma^2}{2} W'(\epsilon) \frac{W^{(\lambda)}(b-\epsilon)}{W^{(\lambda)}(b)} - \int_{-b}^{0-} \frac{W^{(\lambda)}(x+b-\epsilon)}{W^{(\lambda)}(b)} \mathbb{I}_{(-b+\epsilon,0)}(x) \\ \times \int_{0+}^{\infty} \Pi(dx-y) \{W(\epsilon) - W(\epsilon-y)\} dy.$$

3.1. Proof if $W^{(\lambda)'(0+)} < \infty$. First, we assume that X has a Brownian component, that is $\sigma > 0$, or that it has paths of bounded variation; in both cases $W^{(\lambda)'(0+)} < \infty$. Then,

$$\frac{L_1^{\epsilon,b}}{\epsilon} = \psi'(0+) \frac{W(\epsilon)}{\epsilon} + \frac{\sigma^2}{2} \frac{W'(\epsilon)}{W^{(\lambda)}(b+\epsilon)} \left[\frac{Z^{(\lambda)}(b)W^{(\lambda)}(b+\epsilon) - Z^{(\lambda)}(b+\epsilon)W^{(\lambda)}(b)}{\epsilon} \right] \\ \xrightarrow{\epsilon \rightarrow 0} \psi'(0+)W'(0+) + \frac{\sigma^2}{2} \frac{W'(0+)}{W^{(\lambda)}(b)} \left[Z^{(\lambda)}(b)W^{(\lambda)'(b)} - \lambda(W^{(\lambda)}(b))^2 \right],$$

$$\frac{L_2^{\epsilon,b}}{\epsilon} = \int_{-b}^{0-} \left[Z^{(\lambda)}(x+b) - Z^{(\lambda)}(b+\epsilon) \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b+\epsilon)} \right] \int_{0+}^{\infty} \Pi(dx-y) \left\{ \frac{W(\epsilon)}{\epsilon} - \frac{W(\epsilon-y)}{\epsilon} \right\} dy \\ \xrightarrow{\epsilon \rightarrow 0} W'(0+) \int_{-b}^{0-} \left[Z^{(\lambda)}(x+b) - Z^{(\lambda)}(b) \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b)} \right] \int_{0+}^{\infty} \Pi(dx-y) dy$$

and, similarly,

$$\frac{L_3^{\epsilon,b}}{\epsilon} = \psi'(0+) \frac{W(\epsilon)}{\epsilon} + \frac{\sigma^2}{2} \frac{W'(\epsilon)}{W^{(\lambda)}(b+\epsilon)} \left[\frac{W^{(\lambda)}(b+\epsilon) - W^{(\lambda)}(b)}{\epsilon} \right] \\ + \int_{-b}^{0-} \left[1 - \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b+\epsilon)} \right] \int_{0+}^{\infty} \Pi(dx-y) \left\{ \frac{W(\epsilon)}{\epsilon} - \frac{W(\epsilon-y)}{\epsilon} \right\} dy \\ + \int_{-\infty}^{-b} \int_{0+}^{\infty} \Pi(dx-y) \left\{ \frac{W(\epsilon)}{\epsilon} - \frac{W(\epsilon-y)}{\epsilon} \right\} dy \\ \xrightarrow{\epsilon \rightarrow 0} \psi'(0+)W'(0+) + \frac{\sigma^2}{2} W'(0+) \frac{W^{(\lambda)'(b)}}{W^{(\lambda)}(b)} \\ + W'(0+) \int_{-b}^{0-} \left[1 - \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b)} \right] \int_{0+}^{\infty} \Pi(dx-y) dy \\ + W'(0+) \int_{-\infty}^{-b} \int_{0+}^{\infty} \Pi(dx-y) dy.$$

Consequently,

$$\lim_{\epsilon \rightarrow 0} L^{\epsilon, b} = \frac{\psi'(0+) + \frac{\sigma^2}{2} \frac{A_1^{(\lambda)}(b)}{W^{(\lambda)}(b)} + \int_{-b}^{0-} A_2^{(\lambda)}(x) \int_{0+}^{\infty} \Pi(dx - y) dy}{\psi'(0+) + \frac{\sigma^2}{2} \frac{W^{(\lambda)'(b)}}{W^{(\lambda)}(b)} + \int_{-\infty}^{0-} A_3^{(\lambda)}(x) \int_{0+}^{\infty} \Pi(dx - y) dy}.$$

3.2. Proof if $W^{(\lambda)'(0+)} = \infty$. If X does not have a Brownian component and has paths of unbounded variation, then $W^{(\lambda)'(0+)} = \infty$. Therefore, we adapt the previous method as follows:

$$\frac{L_1^{\epsilon, b}}{W(\epsilon)} = \psi'(0+), \text{ for all } \epsilon > 0,$$

$$\begin{aligned} \frac{L_2^{\epsilon, b}}{W(\epsilon)} &= \int_{-b}^{0-} \left[Z^{(\lambda)}(x+b) - Z^{(\lambda)}(b+\epsilon) \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b+\epsilon)} \right] \int_{0+}^{\infty} \Pi(dx - y) \left\{ \frac{W(\epsilon)}{W(\epsilon)} - \frac{W(\epsilon - y)}{W(\epsilon)} \right\} dy \\ &= \int_{-b}^{0-} \left[Z^{(\lambda)}(x+b) - Z^{(\lambda)}(b+\epsilon) \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b+\epsilon)} \right] \int_{0+}^{\infty} \Pi(dx - y) dy \\ &\xrightarrow{\epsilon \rightarrow 0} \int_{-b}^{0-} \left[Z^{(\lambda)}(x+b) - Z^{(\lambda)}(b) \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b)} \right] \int_{0+}^{\infty} \Pi(dx - y) dy \end{aligned}$$

and, similarly,

$$\begin{aligned} \frac{L_3^{\epsilon, b}}{W(\epsilon)} &= \psi'(0+) + \int_{-b}^{0-} \left[1 - \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b+\epsilon)} \right] \int_{0+}^{\infty} \Pi(dx - y) \left\{ \frac{W(\epsilon)}{W(\epsilon)} - \frac{W(\epsilon - y)}{W(\epsilon)} \right\} dy \\ &\quad + \int_{-\infty}^{-b} \int_{0+}^{\infty} \Pi(dx - y) \left\{ \frac{W(\epsilon)}{W(\epsilon)} - \frac{W(\epsilon - y)}{W(\epsilon)} \right\} dy \\ &\xrightarrow{\epsilon \rightarrow 0} \psi'(0+) + \int_{-b}^{0-} \left[1 - \frac{W^{(\lambda)}(x+b)}{W^{(\lambda)}(b)} \right] \int_{0+}^{\infty} \Pi(dx - y) dy + \int_{-\infty}^{-b} \int_{0+}^{\infty} \Pi(dx - y) dy. \end{aligned}$$

3.3. Conclusion of the proof. In all cases, i.e., when X has paths of bounded variation, or of bounded variation with or without a Brownian component, the limiting results for $U^{\epsilon, b}$ can be obtained in a similar fashion. In fact, it can easily be proved that

$$\lim_{\epsilon \rightarrow 0} U_i^{\epsilon, b} = \lim_{\epsilon \rightarrow 0} L_i^{\epsilon, b},$$

for each $i = 1, 2, 3$. Therefore,

$$\lim_{\epsilon \rightarrow 0} U^{\epsilon, b} = \lim_{\epsilon \rightarrow 0} L^{\epsilon, b},$$

and, by the continuity of $W^{(\lambda)}$, this limit is equal to

$$\mathbb{E} \left[e^{-\lambda \int_0^{\tau_{-b}^-} \mathbb{1}_{\{X_s \leq 0\}} ds} \right].$$

Finally, recall that $W^{(\lambda)}(x) = 0$ if $x < 0$; therefore $A_3^{(\lambda)}(x) = 1$ if $x < -b$. Note also that $A_1^{(0)}(b) = W'(b)$ and $A_2^{(0)}(x) = A_3^{(0)}(x)$. This completes the proof of Theorem 2.

4. PROOF OF THEOREM 1

We use the same general idea as in the proof of Theorem 2. Define $L^\epsilon = L^{\epsilon, \infty}$. In this case, using the strong Markov property twice, we obtain

$$\begin{aligned} L^\epsilon &= \mathbb{P}_\epsilon \{ \tau_0^- = \infty \} + L^\epsilon \mathbb{E} \left[e^{-\lambda \tau_\epsilon^+} \right] \int_{-\infty}^{0+} \mathbb{P}_\epsilon \left\{ \tau_0^- < \infty, X_{\tau_0^-} \in dx \right\} \mathbb{E}_x \left[e^{-\lambda \tau_0^+} \right] \\ &= \psi'(0+)W(\epsilon) + L^\epsilon e^{-\Phi(\lambda)\epsilon} \int_{-\infty}^{0+} \mathbb{P}_\epsilon \left\{ \tau_0^- < \infty, X_{\tau_0^-} \in dx \right\} e^{\Phi(\lambda)x}. \end{aligned}$$

One can deduce from [7] (see also Exercise 8.7 in [14]) that

$$\begin{aligned} \int_{-\infty}^{0+} \mathbb{P}_\epsilon \left\{ \tau_0^- < \infty, X_{\tau_0^-} \in dx \right\} e^{\Phi(\lambda)x} &= \mathbb{E}_\epsilon \left[e^{\Phi(\lambda)X_{\tau_0^-}} ; \tau_0^- < \infty \right] \\ &= e^{\Phi(\lambda)\epsilon} - \lambda e^{\Phi(\lambda)\epsilon} \int_0^\epsilon e^{-\Phi(\lambda)x} W(x) dx - \frac{\lambda}{\Phi(\lambda)} W(\epsilon). \end{aligned}$$

It is easily shown that

$$\lim_{\epsilon \rightarrow 0} \frac{\int_0^\epsilon e^{-\Phi(\lambda)x} W(x) dx}{W(\epsilon)} = 0.$$

Indeed, if $W(0+) > 0$ this is trivial and if $W(0+) = 0$ one can use L'Hôpital's rule.

Consequently,

$$\lim_{\epsilon \rightarrow 0} L^\epsilon = \lim_{\epsilon \rightarrow 0} \frac{\psi'(0+)W(\epsilon)}{1 - \left[1 - \lambda \int_0^\epsilon e^{-\Phi(\lambda)x} W(x) dx - \frac{\lambda}{\Phi(\lambda)} e^{-\Phi(\lambda)\epsilon} W(\epsilon) \right]} = \psi'(0+) \frac{\Phi(\lambda)}{\lambda}.$$

Similarly, if we define $U^\epsilon = U^{\epsilon, \infty}$, one can show that

$$\lim_{\epsilon \rightarrow 0} U^\epsilon = \psi'(0+) \frac{\Phi(\lambda)}{\lambda}$$

and the result immediately follows.

5. APPLICATIONS

5.1. Brownian motion with drift. For Brownian motion with drift, i.e., if X is of the form $X_t = mt + \sigma B_t$ with $m = \psi'(0+) \geq 0$ and $\sigma > 0$, where B is a standard Brownian motion, then $\Pi \equiv 0$ and, using Theorem 2, we have

$$\mathbb{E} \left[e^{-\lambda \int_0^{\tau_0^-} \mathbb{1}_{\{X_s \leq 0\}} ds} \right] = \frac{mW^{(\lambda)}(b) + (\sigma^2/2) [Z^{(\lambda)}(b)W^{(\lambda)'}(b) - \lambda(W^{(\lambda)}(b))^2]}{mW^{(\lambda)}(b) + (\sigma^2/2)W^{(\lambda)'}(b)},$$

where, as one can easily verify from the definition,

$$W^{(\lambda)}(x) = \frac{2}{\sqrt{m^2 + 2\lambda\sigma^2}} e^{-(m/\sigma^2)x} \sinh \left((x/\sigma^2) \sqrt{m^2 + 2\lambda\sigma^2} \right),$$

for $x \geq 0$.

Then, for standard Brownian motion, in which case $m = \psi'(0+) = 0$ and $\sigma = 1$, we have, from Theorem 2, that

$$\mathbb{E} \left[e^{-\lambda \int_0^{\tau_-^b} \mathbb{I}_{\{X_s \leq 0\}} ds} \right] = \frac{Z^{(\lambda)}(b)W^{(\lambda)'(b)} - \lambda(W^{(\lambda)}(b))^2}{W^{(\lambda)'(b)},$$

where

$$W^{(\lambda)}(x) = \sqrt{\frac{2}{\lambda}} \sinh(x\sqrt{2\lambda})$$

and then

$$Z^{(\lambda)}(x) = \cosh(x\sqrt{2\lambda}).$$

Recalling that $\cosh^2(x) - \sinh^2(x) = 1$, we get that

$$\begin{aligned} \mathbb{E} \left[e^{-\lambda \int_0^{\tau_-^b} \mathbb{I}_{\{X_s \leq 0\}} ds} \right] &= \frac{2 \cosh^2(b\sqrt{2\lambda}) - \lambda(2/\lambda) \sinh^2(b\sqrt{2\lambda})}{2 \cosh(b\sqrt{2\lambda})} \\ &= \frac{1}{\cosh(b\sqrt{2\lambda})}, \end{aligned}$$

therefore recovering Proposition 4.12 of [12] (see also [11]).

5.2. Insurance risk models. Classical insurance risk models describe the surplus process of an insurance company using a compound Poisson process or a Brownian motion with drift, i.e., special cases of spectrally negative Lévy processes. In those models, it is usually assumed that the *net profit condition* holds; this condition ensures that ruin will not occur almost surely. For a Lévy insurance risk process, i.e., spectrally negative Lévy processes, it amounts to $\mathbb{E}[X_1] = \psi'(0+) > 0$ and it is also equivalent to $\lim_{t \rightarrow \infty} X_t = \infty$ almost surely. An interpretation of Lévy insurance risk models for the surplus modelling of large insurance companies is for instance given in [13].

Classically, the probability of ruin has been the most studied risk measure to evaluate the quality of a company. More recently, the analysis of the duration of the negative surplus, in other words the occupation time of the negative half-line, has also been considered in the compound Poisson case [8] and then in a jump-diffusion model [20]. Recall that Theorem 1 extends Equation (4.9) in [20] where the sum of a compound Poisson process and a Brownian motion is considered.

We now want to provide another link between Theorem 1 and insurance risk models. In [16], a *new* definition of the time to ruin is proposed. In that paper, each excursion of the surplus process X below 0 is accompanied by an independent copy of an independent (of X) and exponentially distributed random variable e_d with mean $1/d$; we will refer to it as the implementation clock. If the duration of a given excursion below 0 is less than its associated implementation clock, then ruin does not occur. More precisely, we assume that ruin occurs at the first time τ_d that an implementation clock rings before the end of its corresponding excursion below 0. It is worth pointing out that the time to ruin τ_d is easily defined when the Lévy insurance risk processes X has sample paths of bounded variation.

Therefore, in [16], the case of a surplus process of bounded variation is considered. In that model, one can show that the probability of ruin in this Parisian risk model with exponential

implementation delays can be expressed as follows:

$$\mathbb{P}\{\tau_d < \infty\} = 1 - \mathbb{E}\left[e^{-d \int_0^\infty \mathbb{I}_{\{X_s \leq 0\}} ds}\right],$$

if the exponential clock has mean $1/d$. Indeed, for each excursion of length T , the probability to survive, i.e., the probability that the exponential random variable associated with it is larger than T , is equal to $\exp\{-dT\}$. Using the independence assumption between the clocks and summing up over all the excursions (there are countably many of them), we get

$$\mathbb{E}\left[e^{-d \int_0^\infty \mathbb{I}_{\{X_s \leq 0\}} ds}\right].$$

Using Theorem 1, we recover the corresponding expression in [16], that is

$$\mathbb{P}\{\tau_d < \infty\} = 1 - \psi'(0+) \frac{\Phi(d)}{d},$$

when the net profit condition $\psi'(0+) > 0$ is verified.

More generally, using Itô's excursion theory for spectrally negative Lévy processes, this Parisian risk model with exponential implementation delays can also be defined when the underlying surplus process has paths of unbounded variation. It suffices to *mark* the Poisson point process of excursions away from zero (see [18] for a definition of the corresponding excursion process) with independent copies of the generic random variable e_d , similar to the proof of Theorem 6.16 in [14]; for an excursion away from zero starting above zero, this time spent above zero is simply ignored. As a consequence, Theorem 1 provides a generalization of the probability of ruin in a general Lévy insurance risk model with exponential implementation delays.

6. ACKNOWLEDGEMENTS

Funding in support of this work was provided by the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Institut de finance mathématique de Montréal (IFM2).

REFERENCES

- [1] J. Bertoin. *Lévy processes*. Cambridge University Press, 1996.
- [2] J. Bertoin. Exponential decay and ergodicity of completely asymmetric Lévy processes in a finite interval. *Ann. Appl. Probab.*, 7(1):156–169, 1997.
- [3] T. Chan, A. E. Kyprianou, and M. Savov. Smoothness of scale functions for spectrally negative Lévy processes. *Probability Theory and Related Fields (to appear)*.
- [4] M. Chesney, M. Jeanblanc-Picqué, and M. Yor. Brownian excursions and Parisian barrier options. *Adv. in Appl. Probab.*, 29(1):165–184, 1997.
- [5] I. Czarna and Z. Palmowski. Ruin probability with Parisian delay for a spectrally negative Lévy risk process. *arXiv:1003.4299v1 [math.PR]*, 2010.
- [6] A. Dassios and S. Wu. Parisian ruin with exponential claims. *submitted*, 2009.
- [7] R. A. Doney. Hitting probabilities for spectrally positive Lévy processes. *J. London Math. Soc. (2)*, 44(3):566–576, 1991.
- [8] A. E. dos Reis. How long is the surplus below zero? *Insurance Math. Econom.*, 12(1):23–38, 1993.
- [9] M. Egami and K. Yamazaki. On scale functions of spectrally negative Lévy processes with phase-type jumps. *arXiv:1005.0064v3 [math.PR]*, 2010.
- [10] F. Hubalek and A. E. Kyprianou. Old and new examples of scale functions for spectrally negative Lévy processes. *Sixth Seminar on Stochastic Analysis, Random Fields and Applications, eds R. Dalang, M. Dozzi, F. Russo. Progress in Probability, Birkhäuser*, 2010.

- [11] K. Itô and H. P. McKean, Jr. *Diffusion processes and their sample paths*. Springer-Verlag, Berlin, 1974.
- [12] I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus*. Springer-Verlag, New York, second edition, 1991.
- [13] C. Klüppelberg and A. E. Kyprianou. On extreme ruinous behaviour of Lévy insurance risk processes. *J. Appl. Probab.*, 43:594–598, 2006.
- [14] A. E. Kyprianou. *Introductory lectures on fluctuations of Lévy processes with applications*. Universitext. Springer-Verlag, Berlin, 2006.
- [15] A. E. Kyprianou and V. Rivero. Special, conjugate and complete scale functions for spectrally negative Lévy processes. *Electron. J. Probab.*, 13:1672–1701, 2008.
- [16] D. Landriault, J.-F. Renaud, and X. Zhou. Insurance risk models with Parisian implementation delays. *preprint*, 2010.
- [17] R. L. Loeffen. An optimal dividends problem with a terminal value for spectrally negative Lévy processes with a completely monotone jump density. *J. Appl. Probab.*, 46(1):85–98, 2009.
- [18] J. Obłój and M. Pistorius. On an explicit Skorokhod embedding for spectrally negative Lévy processes. *J. Theoret. Probab.*, 22(2):418–440, 2009.
- [19] B. A. Surya. Evaluating scale functions of spectrally negative Lévy processes. *J. Appl. Probab.*, 45(1):135–149, 2008.
- [20] C. Zhang and R. Wu. Total duration of negative surplus for the compound Poisson process that is perturbed by diffusion. *J. Appl. Probab.*, 39(3):517–532, 2002.

DEPARTMENT OF STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF WATERLOO, 200 UNIVERSITY AVENUE WEST, WATERLOO (ONTARIO) N2L 3G1, CANADA
E-mail address: `dlandria@math.uwaterloo.ca`

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL (UQAM), 201 AV. PRÉSIDENT-KENNEDY, MONTRÉAL (QUÉBEC) H2X 3Y7, CANADA
E-mail address: `renaud.jf@uqam.ca`

DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY, 1455 DE MAISONNEUVE BLVD W., MONTRÉAL (QUÉBEC) H3G 1M8, CANADA
E-mail address: `xzhou@mathstat.concordia.ca`