

Tree light capture and spatial variability of understory light increase with species mixing and tree size heterogeneity

Gauthier Ligtot^{1*}, Aitor Ameztegui^{2,4,5}, Benoît Courbaud³, Lluís Coll^{2,5,6}, Daniel Kneeshaw⁴

Abstract

Mixed and multi-layered forest ecosystems are sometimes more productive than monospecific and single-layered ones. It has been suggested that trees of different species and sizes occupy complementary positions in space which would act as a mechanism to increase canopy light interception and wood production. However, greater canopy light interception reduces the average amount and variability of transmitted radiation offering fewer opportunities for all species to regenerate and to maintain forest heterogeneity in the long-run. We investigated whether increasing overstory heterogeneity indeed results in greater canopy light interception and lower variability in transmittance. We modeled the three-dimensional structure of forest stands with 3 typical forest structures, 10 mixtures of four tree species, and 3 different basal areas. We used the forest light interception model SAMSARALIGHT and performed three-way analyses of covariance to analyze the effects of the three varied components of forest heterogeneity. We found no evidence that increasing heterogeneity increases canopy light interception. In contrast, homogeneous stands intercept more light than heterogeneous stands. Variability in transmittance increased in some cases with compositional heterogeneity and, to a lesser extent, with tree size inequalities. The advantage of heterogeneous forests is in opportunities for natural regeneration rather than in opportunities to enhance canopy light interception.

¹ Univ. de Liège, Gembloux Agro-Bio Tech, Unité de Gestion des Ressources forestières, 2, Passage des Déportés, B-5030 Gembloux, Belgique

² Forest Sciences Centre of Catalonia (CTFC-CEMFOR), Ctra. Sant Llorenç de Morunys km.2, E-25280 Solsona, Spain

³ Université de Grenoble Alpes, Irstea, Mountain Ecosystems Research Unit, 2 rue de la Papeterie, 38402 Saint Martin d'Hères, France

⁴ Centre d'étude de la Forêt, Université du Québec à Montréal, C.P. 8888, Succ. Centre-Ville, Montréal (Québec), Canada H3C 3P8

⁵ CREAM, Cerdanyola del Vallès, 08193, Spain

⁶ Department of Agriculture and Forest Engineering (EAGROF), University of Lleida, Lleida, 25198, Spain

*Corresponding author: glligt@ulg.ac.be

Contents

1	Introduction	1
2	Methods	2
2.1	Study species	2
2.2	Creating virtual stands	2
2.3	Modeling understory light	3
2.4	Statistical analyses	5
3	Results	5
3.1	Mean transmittance	5
3.2	Transmittance variability	5
4	Discussion	5
4.1	Tree light capture and overstory heterogeneity	8
4.2	Variability of understory light and overstory heterogeneity	9
5	Research perspectives	9
6	Conclusion	10
	Appendices	12

1. Introduction

It has been advocated that forests with many tree species and a great heterogeneity in tree sizes are more productive (Pretzsch and Schütze, 2009; Vallet and Pérot, 2011; Toïgo et al., 2015), resistant and resilient (Jactel et al., 2005) to future changes and harbor a greater biodiversity (Huston, 1979) than monospecific forests in which all trees have a similar size. Over the last 50 to 100 years, humans have applied forest management systems over much of the northern hemisphere that tended to simplify both the structure and composition of many forests (Bengtsson et al., 2000). This simplification resulted in increased production of the targeted product, usually timber of few species highly valued by the market at the time. However, the advantages of these management systems may have obscured long term negative impacts such as reductions of genetic and species diversity (Ledig, 1992; Noss, 1999), rigidity in the face of market and climate variations (Pretzsch et al., 2013), a decrease in forest habitat quality (Franklin et al., 2000) and soil fertility (Reich et al., 2012) and a loss in overall productivity. Many researchers and forest managers now promote the development of more heterogeneous forests

(McElhinny et al., 2005; Puettmann et al., 2009), but our understanding of the sharing and use of resources among species and vegetation strata in heterogeneous forests is limited and impedes our ability to predict the long-term dynamics of such forests. Researchers have usually hypothesized that forest productivity is greater in mixed forests than in the corresponding pure forests because species mixtures should capture more resources, and in particular, more solar radiation due to the layering of the forest canopy and the complementarity and overlapping of tree crowns (Forrester et al., 2006; Kelty, 2006; Yachi and Loreau, 2007; Pretzsch and Schütze, 2009; Bauhus and Schmerbeck, 2010; Vallet and Pérot, 2011). It is also generally assumed that the structural and compositional heterogeneity of the forest overstory directly affects the amount and variability of light available in the understory for tree regeneration (Kelty, 2006). A number of studies have found close relationships between the amount of light transmitted to the understory and canopy density (Vales and Bunnell, 1988; Comeau and Heineman, 2003; Sonohat et al., 2004; Lochhead and Comeau, 2012), stem spatial distribution (i.e. clumped, random or dispersed) (Coates et al., 2003; Drever and Lertzman, 2003; Beaudet et al., 2011; Lochhead and Comeau, 2012; Ligot et al., 2014a), vertical canopy structure or distribution of tree sizes (Drever and Lertzman, 2003; Lochhead and Comeau, 2012; Ligot et al., 2014a), and species composition (Messier et al., 1998; Lochhead and Comeau, 2012). The effects of these factors on understory light have been tested individually and sometimes in pairs, but their combined effects have been particularly difficult to assess.

Moreover, most studies have focused on the average quantity of light that reaches the forest floor but few have considered spatial variability in light conditions (Puerta-Piñero et al., 2007) even though it is also essential for the sustainability of structurally complex forests. Uneven-aged stands harboring trees of different sizes and ages can only persist if sufficient light reaches seedlings at least in scattered patches in the forest understory. Furthermore, the coexistence of different tree species can only be ensured if the variability in understory light is sufficient for all species to be able to survive, grow and reproduce (Ameztegui and Coll, 2011; Ligot et al., 2013). The exact needs in terms of variability in light availability ultimately depend on the interspecific range of shade tolerances (Kobe et al., 1995), but the greater the variability in understory light conditions, the greater the opportunities for multiple species to regenerate. However, the variability in understory light is likely positively correlated with the average quantity of light transmitted through the canopy. As canopy openness increases, both the mean and the range of understory light conditions increase (Canham et al., 1990). Since canopy transmittance is hypothesized to decrease with forest heterogeneity, variability in understory light should also decrease with forest heterogeneity. Sustainable management of heterogeneous forests should therefore optimize both the capture of solar radiation by overstory trees and by understory seedlings, i.e. it should both minimize mean light transmittance and

maximize the variability of light transmittance.

The main question we address in this paper is how does the structural and compositional heterogeneity of the forest overstory affect light interception and the amount and variability of understory light. In particular, our specific questions are: (i) How does overstory structural and compositional heterogeneity affect the total interception of light? (ii) How does structural and compositional heterogeneity of the overstory affect spatial variability of understory light? We hypothesize that stands composed of trees of multiple species and multiple sizes will intercept more light than uniform stands because canopies would be denser with reduced gap area between crowns. Since we expect the variability of light transmittance to be positively correlated with mean light transmittance, we additionally hypothesize that variability in understory light will be reduced with forest heterogeneity.

2. Methods

2.1 Study species

We selected four species for which allometric relationships required to compute canopy light transmittance with three-dimensional radiative transfer models (Ligot et al., 2014b) have been previously calibrated: sessile oak (*Quercus petraea* (Matt.) Liebl.), European beech (*Fagus sylvatica* L.), mountain pine (*Pinus uncinata* Ram ex. DC) and silver fir (*Abies alba* Mill.). In the first study from which species calibrations were determined, Ligot et al. (2014a) monitored 27 mixed stands of sessile oak and European beech in the Belgian Ardennes (50° 15'N, 5°40'E). In a second study, Ameztegui and Coll (2011) monitored 24 mixtures of mountain pine and silver fir in the Spanish Pyrenees (42° 20'N, 1°40'E). These study species provided us with a large gradient of species shade tolerance and adult tree crown properties. In particular, the four study species spanned a gradient from shade tolerant species with very dense and deep ellipsoidal crowns (European beech) to shade intolerant species with sparse and small parabolic crowns (mountain pine). By increasing order of shade tolerance, the list of the study species is pine, oak, fir and beech (Niinemets, 2006).

2.2 Creating virtual stands

To examine how overstory heterogeneity influences the mean and the variability in understory light conditions, we simulated 90 contrasted forest stands including 3 typical forest structures (single-layered, multi-layered with two or three vegetation layers and reverse J-shaped distribution), 10 mixtures of forest composition based on combinations of the four species (beech, oak, beech/oak, pine, fir, fir/pine, oak/pine, beech/fir, beech/oak/pine, beech/fir/pine), and 3 levels of basal area (15, 25 and 35 m²ha⁻¹) (Figure 1). Each virtual stand was 50 × 50 m, and since the position of the trees was randomly drawn from a uniform distribution, each combination was simulated 100 times. We chose a uniform distribution because we found no general evidence of aggregated or regular distributions of

trees in stands of the studied species (the values of the aggregation index Clark-Evans were generally close to 1, see Table 1). The resulting tree spatial distribution of most simulated plots was random (Clark-Evans' index close to 1, range 0.74 to 1.41).

Tree diameters at breast height (DBHs) in the virtual stands were drawn from statistical distributions, depending on the stand structure. To create single-layered stands, tree DBHs were drawn from a truncated normal distribution with two parameters, $N(\mu_1 = 30 \text{ cm}, \sigma_1 = 4.5 \text{ cm}, \min = 6 \text{ cm}, \max = 70 \text{ cm})$, and with all species occupying the same canopy layer. In multi-layered stands composed of two species, tree DBHs were drawn from two truncated normal distributions: $N(\mu_1 = 40 \text{ cm}, \sigma_1 = 6 \text{ cm}, \min = 6 \text{ cm}, \max = 70 \text{ cm})$, and $N(\mu_2 = 25 \text{ cm}, \sigma_2 = 3.75 \text{ cm}, \min = 6 \text{ cm}, \max = 70 \text{ cm})$. In multi-layered stands composed of three species, tree DBHs were drawn from the two previous distributions and from a third one: $N(\mu_1 = 35 \text{ cm}, \sigma_1 = 3.75 \text{ cm}, \min = 6 \text{ cm}, \max = 70 \text{ cm})$. The standard deviation of DBHs was set to correspond to a coefficient of variation of 15%. In mixed multi-layered stands, the less shade-tolerant species occupied the upper vegetation strata. To create a reverse j-shape structure, tree DBHs of all species were sampled from a single truncated exponential distribution: $\exp(k)$ with $E(\text{DBH}) = 1/k = 25 \text{ cm}, \min = 6 \text{ cm}, \max = 70 \text{ cm}$. Altogether these distribution parameters ensured that the quadratic mean diameter of all simulated stands was relatively similar (about 30 cm). In mixed stands composed of two or three species, each species was assigned 50% or 33% of the total basal area, respectively.

This experimental design crossing levels of canopy density, composition and structure would not have been possible with field experiments. It would be illusory to search for representative stands of every combination of these three factors without interference with other factors (e.g. canopy gaps, stand edges, site conditions).

The choice of the simulated levels of basal area, shapes and parameters of DBH distributions and species mixtures was critical as a different choice would have led to different results. Even though most of the simulated stands spanned a realistic range of conditions (Table 1), some of them are rather theoretical (e.g. mixtures of 3 species). However, the light model we used (see next section) is assumed to synthesize our knowledge of radiative transfer through forest canopies and to be valid for different forest structures and compositions.

Tree height and crown radius were computed using species-specific allometric relationships determined by Ameztegui and Coll (2011) and Ligot et al. (2014a) (Fig. 2). Leaf area density (LAD) was set according to previous measurements (Ligot et al., 2014a) to $0.71 \text{ m}^2\text{m}^{-3}$ for beech and $0.52 \text{ m}^2\text{m}^{-3}$ for oak. For both pine and fir, LAD was set to $0.6 \text{ m}^2\text{m}^{-3}$ based on the literature (Ligot et al., 2014b). Moreover, small variations in LAD around 0.6 have been shown to have little effect on light interception simulations (Courbaud et al., 2003; Ligot et al., 2014a).

2.3 Modeling understory light

The amount of light available in the forest understory of each simulated stand was computed with SAMSARALIGHT (Courbaud et al., 2003; Ligot et al., 2014a), a three-dimensional radiative transfer model implemented in a library of the CAPSIS platform (Dufour-Kowalski et al., 2012)¹. This model has proven to satisfactorily predict understory light in uneven-aged forests of Norway spruce in the Alps (*Picea abies* (L.) Karst) (Courbaud et al., 2003), in Belgian mixtures of oak and beech forests (Ligot et al., 2014a) and in Spanish mixtures of fir and pine in the Pyrenees (Appendix A1).

For this experiment, SAMSARALIGHT was set to compute light interception using Beer's law (Eq. 1). Beer's law describes the attenuation of a monochromatic ray within a turbid medium, i.e. a medium made up of small elements randomly scattered and presenting a homogeneous transparency (Brunner, 1998). Briefly, the probability of beam interception ($1 - \tau$) by canopy elements is a function of the canopy element density (leaf area density, LAD, in m^2m^{-3}), the path length of a ray through the canopy (l , in m), the extinction coefficient (k) and the clumping factor (Ω). k and Ω depend on canopy element inclination and spatial distribution:

$$\tau(\eta, \gamma) = \exp(-k \cdot \Omega \cdot \text{LAD} \cdot l(\eta, \gamma)) \quad (1)$$

Tree crowns of hardwood and softwood species were modeled with three-dimensional ellipsoidal and parabolic shapes, respectively, using the allometric relationships from Ligot et al. (2014a) and Ameztegui and Coll (2011). In SAMSARALIGHT, individual tree crowns were considered as homogeneous turbid media. For each month of the growing season (from April to October), 130 diffuse and 81 direct light ray directions were sampled. Diffuse ray directions were sampled over the sky hemisphere at regular zenith angles above a starting value of 10° . Similarly, direct ray directions were sampled every hour over the average monthly solar trajectory. For every direction, parallel rays were cast at the centers of each 25 m^2 ground cell. To remove edge effects, SAMSARALIGHT uses an algorithm that considers plots as being wrapped around a torus (Courbaud et al., 2003). The radiation transmitted to a cell by a light beam depended on the radiation absorption of the succession of crowns that are crossed. The amount of radiation available over a vegetation season for each cell was obtained by cumulating the radiation from light coming from every direction. The amount and angular distribution of incident diffuse and direct lights were computed assuming all sites had latitude of 50°N (i.e. the latitude of the Belgian sites). Simulating light interception at the lower latitude of the Spanish sites (42°N), did not alter our conclusions (results not shown). While latitude affects absolute values of irradiance, the transmittance (i.e. the ratio between the irradiances available above and below forest canopy) is not much affected by latitude.

¹A version of this software can freely be downloaded at <http://orbi.ulg.ac.be/handle/2268/187361> (accessed on the 15th April 2016).




Figure 1. Examples of three virtual stands with basal area of $25 \text{ m}^2 \text{ ha}^{-1}$: a single-layered stand of European beech with a mean DBH of 35 cm (a); a multi-layered stand of sessile oak and beech with mean DBHs of 15 cm for beech (green) and 35 cm for oak (purple) (b); and a reverse J-shaped stand of mountain pine (blue) and silver fir (gray) with mean DBHs of 25 cm (c). Understory light was assessed for each cell. Levels of transmittance under the trees are represented by a gradient of color from black (transmittance of 0-10%) to yellow (90-100%).




Figure 2. Allometric relationships of the four study species, based on Ligot et al. (2014a) and Ameztegui and Coll (2011).

2.4 Statistical analyses

In order to determine the relative importance of the three studied factors (stand composition, structure and basal area), two three-way ANCOVAs were carried out. The response variables were the mean and the standard deviation of light transmittance, i.e. the proportion of incident light transmitted to the understory, computed for each simulation run and each cell. The standard deviation of light transmittance was used as an indicator of the variability of light conditions in the understory. Stand composition, structure and basal area were considered as fixed factors with nine, two and one degrees of freedom, respectively. Since some interactions among factors were significant, multiple one-way ANCOVAs were carried out to further analyze the effect of each factor. To test the differences between the means of factor levels, we used Tukey's "Honest Significant Difference" method (Team, 2013). We graphically examined model residuals to verify that ANCOVA assumptions were fulfilled (i.e. normal distribution of residuals, equal variance among groups, homoscedasticity, and linear relationship). All statistical analyses were performed with R software (Team, 2013).

3. Results

3.1 Mean transmittance

Mean light transmittance (mT) depended on forest composition (62.5% of the total explained variance) and basal area (33.0%), as indicated by type II sum of squares, whereas the interactions between composition and basal area (2.9%), forest structure (1.4%) and the other interactions among factors (0.2%) had minor effects (Table 1). A gradient of mean transmittance was observed in relation to stand composition. Light transmittance was the lowest (i.e. tree light interception was the greatest) in the most shade-tolerant pure beech stands ($m_T = 4.8\%$ on average, $\pm 4.3\%$ of standard deviation) while transmittance in the least shade-tolerant pure pine stands was the greatest ($43.0 \pm 11.5\%$). As expected, mean transmittance clearly decreased with basal area. For instance, the mean transmittance in oak stands with basal area of 15 and $35 \text{ m}^2 \text{ ha}^{-1}$ were $28.7\% (\pm 2.36\%)$ and $7.6\% (\pm 1.31\%)$, respectively.

Mean transmittance of mixed stands was always intermediate between the corresponding pure stands (Fig. 3). For example, transmittance in mixtures of oak/pine ($25.9 \pm 10.8\%$) was intermediate between pure stands of oak (lower light transmitted, $16.9 \pm 9.0\%$) and pine (higher light transmitted, $43.0 \pm 11.5\%$). In the case of mixtures of three species, transmittance was intermediate between the transmittance of the corresponding two most shade-tolerant species mixtures and the transmittance of the pure stand of the most shade-intolerant species although the amount of transmitted light was usually closer to that found in mixtures of the two shade-tolerant species. Again as an example, in beech/fir/pine mixtures, light transmittance ($14.1 \pm 8.3\%$) was slightly greater than in beech/fir mixtures (the two more shade-tolerant species in the mix, $9.2 \pm 6.5\%$).

Although forest structure had a statistically significant effect on the mean of transmittance, its effect was weaker than that of stand composition or basal area (Table 1). In pure stands, transmittance increased with the number of tree layers, being the lowest in single-layered stands ($18.1 \pm 13.9\%$) and the highest in reverse j-shaped stands ($21.0 \pm 14.8\%$). In mixed stands, the mean transmittance of multi-layered stands was in some cases significantly greater than in even-sized stands but in all cases remained lower than in reverse j-shaped stands (Fig. 3).

3.2 Transmittance variability

Similar to mean transmittance, the variability in transmittance depended mostly on forest composition (71.5% of total explained variance, Table 2) and basal area (23.9%). Forest structure (1.2%) and interactions among factors (3.4%) had little effect. The variability in transmittance was also lowest in beech stands and increased according to species shade tolerance (beech < fir < oak < pine), with the exception of pine stands, for which the variability in transmittance was lower than in fir stands. The differences between the variability in transmittance of pine and fir stands were significant for basal area values of 15 and $25 \text{ m}^2 \text{ ha}^{-1}$. As expected, mean transmittance and standard deviation were positively correlated ($r = 0.759$, $p < 2.210^{-16}$). Nevertheless, the relationships between these two variables departed from a linear relationship with a maximum variability in transmittance observed at about a mean transmittance of 40%.

For some combinations of stand structure, composition and basal area, the variability in transmittance in two-species mixtures was greater than the variability in corresponding pure stands (Fig. 4). This happened, for example, in oak and beech mixtures at low basal areas, as well as in the pine and fir mixtures in high-density stands. These relationships were consistent across the three modelled stand structures. No additive effect was, however, observed when softwood species were mixed with hardwood species.

Similar to the results of mean transmittance, the variability in transmittance (σ_T) slightly increased with the number of tree layers in pure stands, being the lowest in single-layered stands ($\sigma_T = 7.5\%$ on average, $\pm 3.9\%$ of standard deviation) and the highest in stands with a reverse j-shaped structure ($8.2 \pm 3.8\%$). In mixed stands, the variability in transmittance of multi-layered stands was in some cases significantly greater than in even-aged stands but remained in all cases lower than in reverse j-shaped structure stands.

The variability in transmittance generally decreased with stand basal area (Fig. 4). Nevertheless, there was a significant interaction between basal area and stand composition (Table 2). The effect of basal area on variability in transmittance was lower in pure stands of shade intolerant or mid-tolerant species (e.g. pine and oak) than in stands of shade-tolerant species (e.g. beech and fir).

4. Discussion

Table 1. A brief description of the plots monitored in oak-beech mixtures by (Ligot et al., 2014a) and in fir-pine mixtures by (Ameztegui and Coll, 2011). The table indicates the number of studied plots (n), the average (minimum and maximum) measured quadratic mean diameter (Dg), coefficient of variation of tree diameters (CV), stand basal area, Clark-Evans aggregation index and transmittance. Regular distributions of trees have Clark-Evans values greater than 1 while aggregated distributions have a Clark-Evans value lower than 1 (Clark and Evans, 1954).

Species	n	Dg Cm	CV %	Basal area m ² ha ⁻¹	Clark-Evans -	Transmittance %
Oak-Beech ¹	27	42.4 (30.0-54.9)	43 (27-60)	18.0 (7.5-35.0)	1.1 (1.0-1.3)	22.8 (0.8-62.6)
Oak-Beech ²	2773	36.0 (6.9-87.3)	38 (0.0-150.5)	20.0 (1.2-70.4)	-	-
Fir-Pine ³	24	30.6 (19.53-42.0)	36 (15-53)	23.3 (7.6-44.5)	0.9 (0.5-1.2)	

These figures came from (1) Ligot et al. (2013), (2) the permanent inventory of forest resources in Wallonie (Lecomte, 2013) and (3) Ameztegui and Coll (2011)




Figure 3. Average light transmittance for 90 virtual forest stands as a function of forest composition, forest structure and basal area. The letters above the boxplots indicate significantly different groups ($\alpha = 0.05$).

Table 2. ANCOVA table of the mean transmittance (aggregated by simulation run). The table indicates, for each factor and interactions, the degrees of freedom (Df.), the type II sum of squares (SumSq.), the percentage of total sum of squares (Sum Sq. %), the mean square (Mean Sq.), F and P values. The R² of this model is 97%.

	Df.	Sum Sq.	Sum Sq. %	Mean Sq.	F value	P value
basal area	1	596270	32.99	596270	114564	0.00E+00
Composition	9	1129719	62.5	125524	24118	0.00E+00
Structure	2	24854	1.37	12427	2388	0.00E+00
basal area * composition	9	52436	2.9	5826	1119	0.00E+00
basal area * structure	2	891	0.05	445	86	1.51E-37
composition * structure	18	2896	0.16	161	31	3.58E-103
basal area * composition * structure	18	550	0.03	31	6	2.54E-14
Residuals	8940	46530		5		




Figure 4. Light variability in the forest understory of 90 virtual forest stands as a function of forest composition, forest structure and basal area. The letters above the boxplots indicates significantly different groups ($\alpha = 0.05$).

Table 3. ANCOVA table of the standard deviation of transmittance predictions: degree of freedom (Df.), the type II sum of squares (Sum Sq.), the percentage of total sum of squares, mean squares (Mean Sq.), F value and P value. The R2 of this model is 88%.

	Df.	Sum Sq.	Sum Sq. %	Mean Sq.	F value	P value
basal area	1	28748	23.91	28748	16042	0.00E+00
Composition	9	85970	71.51	9552	5330	0.00E+00
Structure	2	1453	1.21	726	405	3.20E-169
basal area * composition	9	3530	2.94	392	219	0.00E+00
basal area * structure	2	78	0.07	39	22	3.40E-10
composition * structure	18	312	0.26	17	10	2.86E-27
basal area * composition * structure	18	133	0.11	7	4	8.52E-09
Residuals	8940	16021		2		

4.1 Tree light capture and overstory heterogeneity

In contrast to our first hypothesis, we found no evidence that increasing compositional and structural heterogeneity reduces forest canopy light transmittance or increases tree light interception. Light transmittance in mixed stands was always intermediate between the transmittances in the corresponding pure stands. Moreover, light transmittance slightly increased with increasing forest structural heterogeneity. Stands composed of one homogeneous tree layer transmitted less light than stands with two or more tree layers or stands composed of two or more species. These results do not support the hypothesis that complementarity of light use is at the origin of the overyielding that has been observed in some mixed stands such as the beech-fir mixture (Toïgo et al., 2015) or in stands of higher structural heterogeneity (Pretzsch and Schütze, 2009; Vallet and Pérot, 2011). In contrast, our results are in line with the intermediate productivity found in mixtures such as oak-pine and oak-beech mixtures (Toïgo et al., 2015) and the higher productivity of pure stands with low size heterogeneity (Bourdier et al., 2016).

In addition to the interception of light, light use efficiency and belowground resources also influence forest growth conditions (Richards et al., 2010) and the understory (Granus et al., 2016). Despite subordinates trees receive less light than taller trees. Subordinate trees tend to use intercepted light more efficiently producing greater amount of biomass per unit of light interception than taller trees. In addition, Toïgo et al. (2015) found over-yielding growth in tree mixtures to increase when soil fertility is low (Toïgo et al., 2015). In a recent review, (Forrester, 2014) showed that net complementary effects in mixed-forests depend on the spatial and temporal variability in both above- and belowground resources. They stressed the need for further studies to quantify both the conditions (e.g. environmental gradients) and the processes driving these patterns (i.e. over-yielding, complementary effects).

A potential limit of our approach is that the optimization of light capture in our simulated forest stands is restricted by limited crown plasticity, constrained by allometries and a random spatial distribution of trees. Allometries only roughly estimate individual crown development as they assume that

the lateral growth of a tree crown is isotropic and independent of the past and present surrounding canopy. Yet, anisotropic crown growth and crown plasticity have been mentioned to be important in canopy closure (Purves et al., 2007; Vepakomma et al., 2008, 2011).

Additionally since in many regions tree growth and survival depend on light transmittance the natural spatial distribution of trees might deviate from a random distribution in order to optimize light capture. Forest management also alters the spatial distribution of trees but often tends to regularize it. Because of our approach to modeling crown dimensions and the spatial distribution of trees, we may thus underestimate the complementarity of light use in heterogeneous forests. Nevertheless, a closed layer of foliage encountered in dense monospecific even-sized stands is sufficient to capture most of the available light, as shown by the lack of ground vegetation and seedlings in dense even-sized stands (Bailey and Tappeiner, 1998).

Our findings also underline the importance of forest composition and density on light transmittance. In contrast to the literature (Messier et al., 1998; Lochhead and Comeau, 2012), stand composition had the strongest influence on understory light in our simulations. Stand density, as measured by basal area, was the second most important factor while the heterogeneity in tree size affected little (approx. 1%) understory light levels in comparison to the two other factors. The order of importance of the tested factors may however depend on the chosen study species and levels of basal area. We studied species with contrasted crown shapes and sizes. The effect of the overstory composition on understory light availability is usually explained by differences in species morphological traits related to light interception (Coll et al., 2011). In our case, leaf area density was relatively similar among the studied species (range between 0.5 and 0.7 m² m⁻³) but crown allometries differed greatly (Fig. 2) and these relationships have been shown to critically affect canopy packing (Beaudet et al., 2002; Ligot et al., 2014a). In our study, the largest differences in average transmittance were observed between the simulated softwood and hardwood stands, with softwood species (pine and fir) having narrower and smaller crowns

than hardwood species (oak and beech), thus transmitting more light (Valladares, 2003). However, this result is not generalizable as the studied hardwood and softwood species were species of different shade tolerances (Valladares and Niinemets, 2008) with the hardwood species being the most shade-tolerant species. In addition, individuals of the two studied softwood species grow at higher elevations and in colder sites than those of the two studied hardwood species, with consequences both on allometries and on leaf area density (López-Serrano et al., 2005; Astrup and Larson, 2006; Leuschner et al., 2006; Lefrançois et al., 2008).

4.2 Variability of understory light and overstory heterogeneity

In contrast to our second hypothesis, we did not find a systematic decrease in the range of understory light conditions with overstory heterogeneity. Instead, increasing tree size heterogeneity and especially species heterogeneity resulted in significantly greater variability of understory light conditions, depending on stand composition and basal area. For example, beech, a very shade tolerant species, maintained a very dark understory (low mean and variability of transmittance) even at moderate basal area (e.g. $25 \text{ m}^2 \text{ ha}^{-1}$). Adding beech in mixtures with oak generally limited the variability of understory light except in stands with very low basal area ($15 \text{ m}^2 \text{ ha}^{-1}$). In contrast, the variability of understory light in mixtures of pine and fir was greater than in the corresponding pure stands of either pine or fir, when basal area was high ($\geq 25 \text{ m}^2 \text{ ha}^{-1}$). Forest structure, i.e. the organization of trees in different vertical strata, little affected the variability in understory light conditions. In pure stands, the variability in transmittance increased with the number of tree strata, but in mixed stands, the lack of difference between single-layered and multi-layered stands was not expected. Although an earlier study in mixed hardwood stands in North-East America found little variation in light transmittance with stand age and thus canopy strata complexification (Brown and Parker, 1994). However, we acknowledge that this result may be influenced, at least in part, by the approach we used to generate mixed stands: Mixed, single-layered stands were composed of two species in one single tree layer, with trees of about the same DBH but varying height depending on specific allometries. Therefore, the comparable understory light variability observed between single-layered and multi-layered stands may be due to a lack of sufficient differences in the heterogeneity of tree sizes for these two structures.

The variability in understory light and the mean transmittance through the canopy are positively correlated and hence influenced by similar factors. Both the mean and the variability of transmittance increases as the size of canopy gap increases (Canham et al., 1990; Hardy et al., 2004). Nevertheless, our results further suggest that the relationship is not linear. As corroborated by the data of Da Silva et al. (2011), the variability in understory light is presumably maximized for a certain level of canopy openness that corresponds in

our study to an approximate mean transmittance of 40%. At low understory light conditions ($m_T < 40\%$), the frequency distribution of transmittance is typically right-skewed as most understory patches are in low light conditions and few are in high light conditions (Beaudet et al., 2011). The variability in transmittance then increases with mean transmittance, e.g. increased overstory heterogeneity or decreased overstory density. In contrast, in high light conditions ($m_T > 40\%$) a reduction in mean transmittance could lead to an increase in transmittance variability, e.g. when a shade-tolerant species was mixed in a stand with less shade-tolerant species.

5. Research perspectives

Our findings were obtained through the simulation of stands with a random spatial distribution of trees, fixed levels of basal area and constant quadratic mean diameters of 30 cm. Even though the studied stands were constructed with relatively arbitrary parameters (DBH distribution, composition, basal area), the characteristics of most simulated stands spanned typical conditions found in managed forests of the study species (Table 1) and for these stands the predictions of understory transmittance are well in the range of observed values (Fig. A2). However, further work remains to be done before generalizing our results especially in stands with varying diameters or particular spatial distributions of trees (Ngo Bieng et al., 2006), e.g. aggregation of tree species or aggregation of understory trees in overstory gaps. The former is presumably a key variable of stand structure on understory light (Bourdier et al., 2016) as understory light should increase with stand quadratic mean diameter for a given level of basal area (Lochhead and Comeau, 2012; Ligtot et al., 2014a).

In addition, further modeling efforts are required to analyze the impact of even-aged and uneven-aged silviculture on understory light conditions. Uneven-aged silviculture maintains relatively constant basal area (e.g. about $30 \text{ m}^2 \text{ ha}^{-1}$ in spruce stands with periodic removals of $5\text{-}10 \text{ m}^2 \text{ ha}^{-1}$) while in even-aged silviculture basal area varies greatly (e.g. from 0 to $60 \text{ m}^2 \text{ ha}^{-1}$) during a rotation (i.e. the full cycle from small regeneration to mature stand). A comparison of the effect of these two silviculture regimes on resource acquisition and use by forest stands therefore requires analyses that consider the change of forest structure during a complete rotation.

An investigation of the influence of individual plasticity of crown geometry such as light induced plagiotropy and leaf area density should also be performed as these traits are known to be affected by openings and stand composition (Mitchell, 1975; Umeki, 1996, 1997; Piboule, 2001; Purves et al., 2007; Strigul et al., 2008). As we repeated our simulation 100 times, this effect has probably been partly minimized. Nonetheless, the extent to which individual crowns can expand in reaction to the local environment and how this affect understory light needs to be further studied.

6. Conclusion

The advantage of heterogeneous forest stands may lie in opportunities to naturally regenerate various species in the understory rather than in opportunities to enhance light capture by the overstory. This study has taken a step in the direction of better understanding the effects of forest heterogeneity on light capture and light distribution between the canopy and regeneration layers, which is a necessary step in the current debate on increasing forest heterogeneity as a technique for making forest management more sustainable.

Acknowledgements

We are particularly grateful to François de Coligny (INRA, AMAP) for his help programming the model of forest radiative transfer (SAMSARALIGHT) and Adeline Fayolle (ULg, Gembloux Agro-Bio Tech) for her precious comments on the manuscript.

References

- Ameztegui, A. and Coll, L., 2011. Tree dynamics and coexistence in the montane-sub-alpine ecotone: the role of different light-induced strategies. *J. Veg. Sci.*, 22(6):1049–1061. doi: 10.1111/j.1654-1103.2011.01316.x.
- Astrup, R. and Larson, B. C., 2006. Regional variability of species-specific crown openness for aspen and spruce in western boreal Canada. *For. Ecol. Manage.*, 228(1-3): 241–250. doi: 10.1016/j.foreco.2006.02.048.
- Bailey, J. D. and Tappeiner, J. C., 1998. Effects of thinning on structural development in 40- to 100-year-old Douglas-fir stands in western Oregon. *For. Ecol. Manage.*, 108(1-2):99–113. doi: [http://dx.doi.org/10.1016/S0378-1127\(98\)00216-3](http://dx.doi.org/10.1016/S0378-1127(98)00216-3).
- Bauhus, J. and Schmerbeck, J., 2010. *Silvicultural options to enhance and use forest plantation biodiversity*. Ecosystem goods and services from plantation forests. Earthscan, London.
- Beaudet, M., Messier, C., and Canham, C. D., 2002. Predictions of understory light conditions in northern hardwood forests following parameterization, sensitivity analysis, and tests of the SORTIE light model. *For. Ecol. Manage.*, 165(1-3): 235–248. doi: 10.1016/S0378-1127(01)00621-1.
- Beaudet, M., Harvey, B. D., Messier, C., Coates, K. D., Poulin, J., Kneeshaw, D. D., Brais, S., and Bergeron, Y., 2011. Managing understory light conditions in boreal mixedwoods through variation in the intensity and spatial pattern of harvest: a modelling approach. *For. Ecol. Manage.*, 261(1): 84–94. doi: 10.1016/j.foreco.2010.09.033.
- Bengtsson, J., Nilsson, S. G., Franc, A., and Menozzi, P., 2000. Biodiversity, disturbances, ecosystem function and management of European forests. *For. Ecol. Manage.*, 132(1):39–50. doi: [http://dx.doi.org/10.1016/S0378-1127\(00\)00378-9](http://dx.doi.org/10.1016/S0378-1127(00)00378-9).
- Bourdier, T., Cordonnier, T., Kunstler, G., Piedallu, C., Lagarigues, G., and Courbaud, B., 2016. Size inequality reduces forest productivity: An analysis combining inventory data for ten European species and a light competition model. *PLoS one*, 11(3). doi: 10.1371/journal.pone.0151852.
- Brown, M. and Parker, G., 1994. Canopy light transmittance in a chronosequence of mixed-species deciduous forests. *Canadian Journal of Forest Research-Revue Canadienne De Recherche Forestiere*, 24(8):1694–1703.
- Brunner, A., 1998. A light model for spatially explicit forest stand models. *For. Ecol. Manage.*, 107(1-3):19–46. doi: 10.1016/S0378-1127(97)00325-3.
- Canham, C. D., Denslow, J. S., Platt, W. J., Runkle, J. R., Spies, T. A., and White, P. S., 1990. Light regimes beneath closed canopies and tree-fall gaps in temperate and tropical forests. *Can. J. For. Res.*, 20(5):620–631. doi: 10.1139/x90-084.
- Clark, P. J. and Evans, F. C., 1954. Distance to nearest neighbor as a measure of spatial relationships in populations. *Ecology*, 35(4):445–453. doi: 10.2307/1931034.
- Coates, K. D., Canham, C. D., Beaudet, M., Sachs, D. L., and Messier, C., 2003. Use of a spatially explicit individual-tree model (SORTIE) to explore the implications of patchiness in structurally complex forests. *For. Ecol. Manage.*, 186(1-3):297–310. doi: 10.1016/S0378-1127(03)00301-3.
- Coll, L., González-Olabarria, J. R., Mola-Yudego, B., Pukkala, T., and Messier, C., 2011. Predicting understory maximum shrubs cover using altitude and overstory basal area in different Mediterranean forests. *Eur. J. For. Res.*, 130(1): 55–65. doi: 10.1007/s10342-010-0395-y.
- Comeau, P. G. and Heineman, J. L., 2003. Predicting understory light microclimate from stand parameters in young paper birch (*Betula papyrifera* Marsh.) stands. *For. Ecol. Manage.*, 180(1-3):303–315. doi: 10.1016/S0378-1127(02)00581-9.
- Courbaud, B., de Coligny, F., and Cordonnier, T., 2003. Simulating radiation distribution in a heterogeneous Norway spruce forest on a slope. *Agric. For. Meteorol.*, 116(1-2): 1–18. doi: 10.1016/S0168-1923(02)00254-X.
- Da Silva, D., Balandier, P., Boudon, F., Marquier, A., and Godin, C., 2011. Modeling of light transmission under heterogeneous forest canopy: an appraisal of the effect of the precision level of crown description. *Ann. For. Sci.*, 69(2):191–193. doi: 10.1007/s13595-011-0139-2.
- Drever, C. R. and Lertzman, K. P., 2003. Effects of a wide gradient of retained tree structure on understory light in coastal Douglas-fir forests. *Can. J. For. Res.*, 33(1):137–146. doi: 10.1139/x02-167.

- Dufour-Kowalski, S., Courbaud, B., Dreyfus, P., Meredieu, C., and De Coligny, F., 2012. Capsis: an open software framework and community for forest growth modelling. *Ann. For. Sci.*, 69(2):221–233. doi: 10.1007/s13595-011-0140-9.
- Forrester, D. I., 2014. The spatial and temporal dynamics of species interactions in mixed-species forests: From pattern to process. *For. Ecol. Manage.*, 312:282–292. doi: <http://dx.doi.org/10.1016/j.foreco.2013.10.003>.
- Forrester, D. I., Bauhus, J., Cowie, A. L., and Vanclay, J. K., 2006. Mixed-species plantations of eucalyptus with nitrogen-fixing trees: A review. *For. Ecol. Manage.*, 233(2-3):211–230. doi: 10.1016/j.foreco.2006.05.012.
- Franklin, J. F., Lindenmayer, D., MacMahon, J. A., McKee, A., Magnuson, J., Perry, D. A., Waide, R., and Foster, D., 2000. Threads of continuity. *Conservation in Practice*, 1(1):8–17. doi: 10.1111/j.1526-4629.2000.tb00155.x.
- Granhus, A., Metslaid, M., and von Lüpke, N., 2016. Effects of nutrient supply on understory picea abies seedling growth in partially cut stands in south-east norway. *Forestry*. doi: 10.1093/forestry/cpv056.
- Hardy, J. P., Melloh, R., Koenig, G., Marks, D., Winstral, A., Pomeroy, J. W., and Link, T., 2004. Solar radiation transmission through conifer canopies. *Agric. For. Meteorol.*, 126(3-4):257–270. doi: 10.1016/j.agrformet.2004.06.012.
- Huston, M., 1979. A general hypothesis of species diversity. *Am. Nat.*, 113(1):81–101.
- Jactel, H., Brockerhoff, E., and Duelli, P., 2005. A test of the biodiversity-stability theory: meta-analysis of tree species diversity effects on insect pest infestations, and re-examination of responsible factors. *For. Diversity and Funct.*, 176:235–262. doi: 10.1007/3-540-26599-6_12.
- Kelty, M. J., 2006. The role of species mixtures in plantation forestry. *For. Ecol. Manage.*, 233(2-3):195–204. doi: 10.1016/j.foreco.2006.05.011.
- Kobe, R., Pacala, S., Silander Jr, J., and Canham, C., 1995. Juvenile tree survivorship as a component of shade tolerance. *Ecol. Appl.*, 5(2):517–532. doi: 10.2307/1942040.
- Lecomte, H., 2013. Etat des lieux de la forêt résineuse wallonne : Quelques chiffres de l'iprfw.
- Ledig, F. T., 1992. Human impacts on genetic diversity in forest ecosystems. *Oikos*, 63(1):87–108. doi: 10.2307/3545518.
- Lefrançois, M.-L., Beaudet, M., and Messier, C., 2008. Crown openness as influenced by tree and site characteristics for yellow birch, sugar maple, and eastern hemlock. *Can. J. For. Res.*, 38(3):488–497. doi: 10.1139/X07-177.
- Leuschner, C., Voss, S., Foetzki, A., and Clases, Y., 2006. Variation in leaf area index and stand leaf mass of european beech across gradients of soil acidity and precipitation. *Plant Ecol.*, 186(2):247–258. doi: 10.1007/s11258-006-9127-2.
- Ligot, G., Balandier, P., Coubraud, B., Jonard, M., Kneeshaw, D., and Claessens, H., 2014a. Managing understory light to maintain a mixture of species with different shade tolerance. *For. Ecol. Manage.*, 327(1):189–200. doi: 10.1016/j.foreco.2014.05.010.
- Ligot, G., Balandier, P., Courbaud, B., and Claessens, H., 2014b. Forest radiative transfer models: which approach for which application? *Can. J. For. Res.*, 44(5):385–397. doi: 10.1139/cjfr-2013-0494.
- Ligot, G., Balandier, P., Fayolle, A., Lejeune, P., and Claessens, H., 2013. Height competition between quercus petraea and fagus sylvatica natural regeneration in mixed and uneven-aged stands. *For. Ecol. Manage.*, 304:391–398. doi: 10.1016/j.foreco.2013.05.050.
- Lochhead, K. D. and Comeau, P. G., 2012. Relationships between forest structure, understorey light and regeneration in complex douglas-fir dominated stands in south-eastern british columbia. *For. Ecol. Manage.*, 284:12–22. doi: 10.1016/j.foreco.2012.07.029.
- López-Serrano, F. R., García-Morote, A., Andrés-Abellán, M., Tendero, A., and Del Cerro, A., 2005. Site and weather effects in allometries: A simple approach to climate change effect on pines. *For. Ecol. Manage.*, 215(1-3):251–270. doi: 10.1016/j.foreco.2005.05.014.
- McElhinny, C., Gibbons, P., Brack, C., and Bauhus, J., 2005. Forest and woodland stand structural complexity: Its definition and measurement. *For. Ecol. Manage.*, 218(1-3):1–24. doi: 10.1016/j.foreco.2005.08.034.
- Messier, C., Parent, S., and Bergeron, Y., 1998. Effects of overstory and understory vegetation on the understory light environment in mixed boreal forests. *J. Veg. Sci.*, 9(4): 511–520. doi: 10.2307/3237266.
- Mitchell, K. J., 1975. Dynamics and simulated yield of douglas-fir. *For. Sci.*, 21(4):1–39.
- Ngo Bieng, M. A., Ginisty, C., Goreaud, F., and Perot, T., 2006. First typology of oak and scots pine mixed stands in orléans forest (france), based on the canopy spatial structure. *N. Z. J. For. Sci.*, 36:325–346.
- Niinemets, ., 2006. The controversy over traits conferring shade-tolerance in trees: ontogenetic changes revisited. *J. Ecol.*, 94(2):464–470. doi: 10.1111/j.1365-2745.2006.01093.x.

- Noss, R. F., 1999. Assessing and monitoring forest biodiversity: A suggested framework and indicators. *For. Ecol. Manage.*, 115(2-3):135–146. doi: 10.1016/S0378-1127(98)00394-6.
- Piboule, A., 2001. *Validation et analyse de sensibilité d'un modèle de transfert radiatif en vue de son application à la cartographie de l'éclaircissement en peuplement forestier*. D.e.a.
- Pretzsch, H., Schütze, G., and Uhl, E., 2013. Resistance of european tree species to drought stress in mixed versus pure forests: Evidence of stress release by inter-specific facilitation. *Plant Biol.*, 15(3):483–495. doi: 10.1111/j.1438-8677.2012.00670.x.
- Pretzsch, H. and Schütze, G., 2009. Transgressive overyielding in mixed compared with pure stands of norway spruce and european beech in central europe: evidence on stand level and explanation on individual tree level. *Eur. J. For. Res.*, 128(2):183–204. doi: 10.1007/s10342-008-0215-9.
- Puerta-Piñero, C., Gómez, J. M., and Valladares, F., 2007. Irradiance and oak seedling survival and growth in a heterogeneous environment. *For. Ecol. Manage.*, 242:462–469. doi: 10.1016/j.foreco.2007.01.079.
- Puettmann, K. J., Coates, K. D., and Messier, C. C., 2009. *A critique of silviculture: managing for complexity*. Island Press, Washington, USA.
- Purves, D. W., Lichstein, J. W., and Pacala, S. W., 2007. Crown plasticity and competition for canopy space: A new spatially implicit model parameterized for 250 north american tree species. *PLoS one*, 2(9):1–11. doi: 10.1371/journal.pone.0000870.
- Reich, P. B., Tilman, D., Isbell, F., Mueller, K., Hobbie, S. E., Flynn, D. F. B., and Eisenhauer, N., 2012. Impacts of biodiversity loss escalate through time as redundancy fades. *Science*, 336(6081):589–592. doi: 10.1126/science.1217909.
- Richards, A. E., Forrester, D. I., Bauhus, J., and Scherer-Lorenzen, M., 2010. The influence of mixed tree plantations on the nutrition of individual species: a review. *Tree Physiol.* doi: 10.1093/treephys/tpq035.
- Sonohat, G., Balandier, P., and Ruchaud, F., 2004. Predicting solar radiation transmittance in the understory of even-aged coniferous stands in temperate forests. *Ann. For. Sci.*, 61(7):629–641. doi: 10.1051/forest:2004061.
- Strigul, N., Pristinski, D., Purves, D., Dushoff, J., and Pacala, S., 2008. Scaling from trees to forests: Tractable macroscopic equations for forest dynamics. *Ecol. Monogr.*, 78(4): 523–545. doi: 10.1890/08-0082.1.
- Team, R. C., 2013. R: A language and environment for statistical computing.
- Toïgo, M., Vallet, P., Perot, T., Bontemps, J., Piedallu, C., and Courbaud, B., 2015. Overyielding in mixed forests decreases with site productivity. *J. Ecol.*, 103(2):502–512. doi: 10.1111/1365-2745.12353.
- Umeki, K., 1996. The effects of plasticity in crown shape on size-structure dynamics of population. *Jpn. Ecol.*, 46: 87–92.
- Umeki, K., 1997. Effect of crown asymmetry on size-structure dynamics of plant populations. *Ann. Bot.*, 79(6):631–641. doi: 10.1006/anbo.1996.0388.
- Vales, D. J. and Bunnell, F. L., 1988. Relationships between transmission of solar radiation and coniferous forest stand characteristics. *Agric. For. Meteorol.*, 43(3-4):201–223. doi: 10.1016/0168-1923(88)90049-4.
- Valladares, F. and Niinemets, U., 2008. Shade tolerance, a key plant feature of complex nature and consequences. *Annual Review of Ecology, Evolution, and Systematics*, 39(1):237–257. doi: 10.1146/annurev.ecolsys.39.110707.173506.
- Valladares, F., 2003. Light heterogeneity and plants: from ecophysiology to species coexistence and biodiversity.
- Vallet, P. and Pérot, T., 2011. Silver fir stand productivity is enhanced when mixed with norway spruce: Evidence based on large-scale inventory data and a generic modelling approach. *J. Veg. Sci.*, 22(5):932–942. doi: 10.1111/j.1654-1103.2011.01288.x.
- Vepakomma, U., St-Onge, B., and Kneeshaw, D., 2011. Response of a boreal forest to canopy opening: Assessing vertical and lateral tree growth with multi-temporal lidar data. *Ecol. Appl.*, 21(1):99–121.
- Vepakomma, U., St-Onge, B., and Kneeshaw, D., 2008. Spatially explicit characterization of boreal forest gap dynamics using multi-temporal lidar data. *Remote Sens. Environ.*, 112(5):2326–2340. doi: 10.1016/j.rse.2007.10.001.
- Yachi, S. and Loreau, M., 2007. Does complementary resource use enhance ecosystem functioning? a model of light competition in plant communities. *Ecol. Lett.*, 10(1): 54–62. doi: 10.1111/j.1461-0248.2006.00994.x.

Appendices




Figure A1. Comparison of prediction and measures with hemispherical photographs of transmittance in Spanish forests composed of mountain pine and silver fir (on the left) and Belgian forests of European beech and oak (on the right). Predictions were computed with two radiative transfer models: SORTIE-ND (in red) and SAMSARALIGHT (in grey). SAMSARALIGHT model was chosen to perform the simulation of this study and its performance was further described by [Ligot et al. \(2014a\)](#).




Figure A2. Comparison of the simulated values of transmittance and values of transmittance recorded in previous studies along a gradient of stand basal area. The two plots on the left show values for mixtures of oak and beech (Ligot et al., 2013) while the two plots on the right show values for fir and pine mixtures (Ameztegui and Coll, 2011). Simulated values are depicted with boxplots while observed values are depicted with empty dots. Ligot et al. (2013) did not sample plots along the whole basal area gradient – even though such stands exist (Table 1) - because their study focused on partially open forests with natural regeneration.